

Checklist of the aphyllphoraceous fungi (*Agaricomycetes*) of the Brazilian Amazonia

ALLYNE CHRISTINA GOMES-SILVA¹ & TATIANA BAPTISTA GIBERTONI¹

allynefungi@hotmail.com *tbgibertoni@hotmail.com*
Universidade Federal de Pernambuco, Departamento de Micologia
Av. Nelson Chaves s/n, CEP 50760-420, Recife, PE, Brazil

Abstract — A literature-based checklist of the aphyllphoraceous fungi reported from the Brazilian Amazonia was compiled. Two hundred and sixteen species, 90 genera, 22 families, and 9 orders (*Agaricales*, *Auriculariales*, *Cantharellales*, *Corticiales*, *Gloeophyllales*, *Hymenochaetales*, *Polyporales*, *Russulales* and *Trechisporales*) have been reported from the area. A summary of the data in the checklist is presented here and the checklist is available at <http://www.mycotaxon.com/resources/weblast.html>.

Key words — macrofungi, neotropics

Introduction

The aphyllphoraceous fungi are currently spread throughout many orders of *Agaricomycetes* (Hibbett et al. 2007) and comprise species that function as major decomposers of plant organic matter (Alexopoulos et al. 1996).

The Amazonian Forest (00°44'–06°24'S / 58°05'–68°01'W) covers an area of 7×10^6 km² in nine South American countries. Around 63% of the forest is located in nine Brazilian States (Acre, Amazonas, Amapá, Pará, Rondônia, Roraima, Tocantins, west of Maranhão, and north of Mato Grosso) (FIG. 1). The Amazonian forest consists of a mosaic of different habitats, such as open ombrophilous, stational semi-deciduous, mountain, “terra firme,” “várzea” and “igapó” forests, and “campinaranas” (Amazonian savannahs). Six months of dry season and six months of rainy season can be observed (Museu Paraense Emílio Goeldi 2007).

Even with the high biodiversity of Amazonia and the well-documented importance of aphyllphoraceous fungi to all arboreal ecosystems, few studies have been undertaken in the Brazilian Amazonia on this group of fungi (Bononi 1981, 1992, Capelari & Maziero 1988, Gomes-Silva et al. 2008, Jesus 1996, Martin-Júnior et al. 2008, Sotão et al. 1997, 2002, 2008). This work aims to contribute to the knowledge about the diversity of the aphyllphoraceous



FIGURE 1. The nine Brazilian States (Acre - AC, Amazonas - AM, Amapá - AP, Pará - PA, Rondônia - RO, Roraima - RR, Tocantins - TO, west of Maranhão - MA, and north of Mato Grosso - MT) where the Brazilian Amazonia is found.

fungi in this high diverse biome by compiling and updating the nomenclature of published records of aphylloraceous fungi from the Brazilian Amazon.

Material and methods

All available publications with records of aphylloraceous fungi (*Agaricomycetes*) collected in the Brazilian Amazonia were examined. The reports were compiled and the nomenclature of the listed species were checked against and — when necessary— updated following the classification of Index Fungorum (<http://www.indexfungorum.org/Names/Names.asp>) and CBS (<http://www.cbs.knaw.nl/databases/>). Taxa with unresolved nomenclature according to the consulted databases and those lacking neo- or pantropical distributions were excluded. The occurrence and distribution of the species in Brazilian states were recorded. The checklist is organized in alphabetical order by families, genus, and species.

Results and discussion

Currently, 216 species of aphyllorhaceous fungi previously recorded from the Brazilian Amazonia are included in the checklist. These species are at present distributed in 90 genera, 22 families, and 9 orders (*Agaricales*, *Auriculariales*, *Cantharellales*, *Corticiales*, *Gloeophyllales*, *Hymenochaetales*, *Polyporales*, *Russulales* and *Trechisporales*). *Polyporales* has the highest number of species (144), families (7), and genera (61). Among the families, *Polyporaceae* is the most species rich (79), followed by *Hymenochaetaceae* (32), and *Meruliaceae* (29). *Botryobasidiaceae*, *Cantharellaceae*, *Corticaceae*, *Cystostereaceae*, *Meripilaceae*, *Peniophoraceae*, *Schizophyllaceae* and *Schizoporaceae* are each represented by one species. Among the genera, *Amauroderma* (16), *Phellinus* (15), *Trametes* (14), *Hymenochaete* (8), *Polystictus* (7), *Ganoderma* (7), and *Podoscypha* (7) have the highest number of species. *Phellinus gilvus* (Schwein.) Pat. and *Earliella scabrosa* (Pers.) Gilb. & Ryvarden have been recorded in all states of the Brazilian Amazonia. The State of Pará is represented by 126 species (58%), with 53 exclusive to the State. The State of Amazonas has 86 reported species (35 exclusive) and 35 species were recorded from the State of Roraima (15 exclusive). The states of Rondônia and Acre recorded 48 and 28 species respectively, each with 7 exclusive species and the states of Mato Grosso and Amapá recorded 36 and 23 species respectively, each with 6 exclusive species. Of the excluded species, all of which should have their exsiccates reviewed, 22 were excluded due to their non neo- or pantropical distribution and 65 due to nomenclature uncertainties according to the consulted databases.

Acknowledgments

The authors would like to thank Dr. Greg Mueller and Dr. Leif Ryvarden for critically reviewing of the manuscript and CNPq for the master scholarship of ACGS.

Literature cited

- Alexopoulos CJ, Mims CW, Blackwell M. 1996. *Introductory Mycology*. 4th ed., John Wiley and Sons, Inc., Nova York, 868p.
- Bononi VL. 1981. Alguns Basidiomicetos hidnóides da região amazônica. *Rickia*, 9: 17–30.
- Bononi VL. 1992. Fungos macroscópicos de Rio Branco, Acre, Brasil. *Hoehnea*, 19(1/2): 31–37.
- Capelari M, Maziero R. 1988. Fungos macroscópicos do estado de Rondônia, região dos rios Jaru e Ji-Paraná. *Hoehnea*, 15: 28–36.
- Gomes-Silva AC, Gibertoni TB, Ryvarden L. 2008. *Coltricia fragilissima* a new record to Brazil. *Mycotaxon*, 105: 469–472.
- Hibbett DS, Binder M, Bischoff JF, Blackwell M, Cannon PF, Eriksson OE, Huhndorf S, James T, Kirk PM, Lücking R, Thorsten-Lumbsch H, Lutzoni F, Matheny PB, McLaughlin DJ, Powell MJ, Redhead S, Schoch CL, Spatafora JW, Stalpers JA, Vilgalys R, Aime MC, Aptroot A, Bauer R, Begerow D, Benny GL, Castlebury LA, Crous PW, Dai YC, Gams W, Geiser DM, Griffith GW, Gueidan C, Hawksworth DL, Hestmark G, Hosaka K, Humber RA, Hyde KD, Ironside

- JE, Køljalg U, Kurtzman CP, Larsson KH, Lichtwardt R, Longcore J, Miadlikowska J, Miller A, Moncalvo JM, Mozley-Standridge S, Oberwinkler F, Parmasto E, Reeb V, Rogers JD, Roux C, Ryvarden L, Sampaio JP, Schüßler A, Sugiyama J, Thorn RG, Tibell L, Untereiner WA, Walker C, Wang Z, Weir A, Weiss M, White MM, Winka K, Yao YJ, Zhang N. 2007. A higher-level phylogenetic classification of the Fungi. *Mycological Research* 111(5): 509–547.
- Jesus MA. 1996. Contribution to the knowledge of wood-rotting fungi in Brazil. II. Checklist of fungi from Maraca Island, Roraima State. *Mycotaxon*, 57: 323–328.
- Martins-Júnior AS, Gibertoni TB, Sotão HMP. 2008. *Diplomitoporus allantoporus* (*Basidiomycetes*): a new record to Brazil. *Mycotaxon*, 106: 195–198.
- Museu Paraense Emílio Goeldi. 2007. Amazônia: Fundamentos da ecologia da maior região de florestas tropicais. http://www.museu-goeldi.br/biodiversidade/o_amazonia.asp. Acesso em 06/06/2007.
- Sotão HMP, Hennen JF, Gugliotta AM, Melo OA, Campos EL. 1997. Os fungos – *Basidiomycotina*. In: Lisboa, P.L.B. (org.). Caxiuanã. Belém, Museu Paraense Emílio Goeldi. 213–219.
- Sotão HMP, Gugliotta AM, Oliveira AP, Luz AB, Melo OA. 2002. Fungos poliporóides. In: Lisboa, P.L.B. (org.). Caxiuanã, populações tradicionais, meio físico e diversidade biológica. Belém: Museu Paraense Emílio Goeldi. p. 433–444.
- Sotão HMP, Gibertoni TB, Maziero R, Baseia I, Medeiros PS, Martins-Júnior A, Capelari M. 2008. Fungos macroscópios da Floresta nacional de Caxiuanã, Pará, Brasil: *Basidiomycota* (*Agaricomycetes*). In: Lisboa PLB. (org.). Caxiuanã: Desafios para conservação de uma Floresta Nacional na Amazônia. Belém: Museu Paraense Emílio Goeldi (in press).