

A new species of *Hohenbuehelia* from China

YU LIU & TOLGOR BAU *

Junwusuo@126.com

*Institute of Mycology, Jilin Agricultural University
Changchun 130118, China*

Abstract — *Hohenbuehelia olivacea* from China is described as new to science.

Key words — *Basidiomycota*, *Pleurotaceae*, taxonomy

Introduction

The genus *Hohenbuehelia* was established by Schulzer (Schulzer et al. 1866), and it belongs to the family *Pleurotaceae* (Kirk et al. 2001). The main characteristics of the genus are small to large basidiomata, gills that are decurrent or radiate from a point of central or lateral attachment on the under side of the cap, sessile or stipitate with a lateral pseudostipe (rarely a central stipe), a gelatinous zone often forming below the cap cuticle, monomitic and clamped hyphae, thick-walled metuloids, fusiform cheilocystidia, and commonly with an hour-glass secretory cell surrounded by a mucous droplet at the tip of a short or elongated neck (Thorn 1986, Corner 1994).

In earlier studies (Teng 1963; He 1992; Bi et al. 1993, 1997; Chang & Mao 1995; Mao 1998; Chang et al. 2001; Li & Bau 2003), eleven taxa representing *Hohenbuehelia* have been recorded in China. Recently, an additional new species was discovered during the research on the genus based on morphological examinations of collections.

Materials and methods

Specimens were examined with traditional taxonomic methods. KOH solution and Melzer's reagent were used as the mountants when examining the microstructure. Morphological characteristics of the species were described and illustrated according to the observation of the materials. Colour descriptions for

* Author for correspondence

the new species refer to Ridgway (1912). The specimen studied is now housed in the Herbarium of Mycology of Jilin Agricultural University (HMJAU).

Taxonomy

Hohenbuehelia olivacea Yu Liu & T. Bau, sp. nov.

FIG. 1

MYCOBANK MB 513041

Pileus 20–35 mm *latus*, *reniformis ad dimidiatus*, *flavirens vel fulvus*, *marginē galbinus vel olivaceus*, *strigosus*, *strigis olivaceis prope basem*. *Lamellae decurrentes, flavae, crebrae*. *Stipes nullus*. *Basidiosporae* (6–)6.5–7.5(–8) × (3–)3.5–4 μm, *ellipsoideae*. *Basidia* 25–31 × 5.5–6.5 μm, *clavata*, *4-sporigera*. *Cheilocystidia* 25–32 × 5–7 μm, *clavata*, *vel ventricosofusiformia*, *saepe capitulata*. *Metuloidae* 52–65 × 9–13 μm, *sublanceolatae vel subfusioideae*, *apice incrustatae*. *Pileipellis ex hyphis* 3–4 μm *crassis*, *repentis vel ascendentis*. *Superficies pilei sine cystidis*.

HOLOTYPE: China, Jilin Province, Lushuihe Town, 30 June 2005, Tolgor Bau, HMJAU 7290.

ETYMOLOGY: The species epithet *olivacea* refers to its pileus colour.

BASIDIOMATA small to middle-sized, reniform or dimidiate, sessile. **PILEUS** 20–35 mm broad, pale yellow-brown to olive, with dense and long tomentum, strigose and slate-“olive” colour towards the base, finely and shortly tomentose and pale “yellow green” or olive at margin when dry. **CONTEXT** thin, white to brownish. **GELATINOUS ZONE** obviously visible, light brown to pallid brown, glossed. **LAMELLAE** “pallid yellow”, narrow and moderately close, thin, decurrent. **STIPE** absent or sometimes covered with white mycelium at the base. **SPORE PRINT** white.

BASIDIOSPORES ellipsoid, (6–)6.5–7.5(–8) × (3–)3.5–4 μm, average 6.93 × 3.63 μm, Q = 1.86–2.00, average = 1.94, smooth, hyaline, colourless, thin-walled, multiguttulate. **BASIDIA** 25–31 × 5.5–6.5 μm, clavate, with four short sterigmata 1.5–2 μm long. **CHEILOCYSTIDIA** 25–32 × 5–7 μm, subclavate, fusoid-ventricose to clavate-capitate, thin-walled, hyaline, with a globular mucous droplet at the tip of a short or elongated neck; the droplet 2.5–3 × 5–7 μm, thick-walled, glossy, becoming hour-glass secretory cell when mature. **METULOIDS** sublanceolate, cyprinoid or fish-fusiform shape, 52–65 × 9–13 μm, gradually enlarged upwards and then with a sharp apex, pale yellow, thick-walled, and crystal-encrusted (about 1/5–1/4) at apex, becoming thinner at the base, implanted into hymenium. **LAMELLAR TRAMA** hyphae parallel, 3–5 μm diam., sometimes inflated to 6–8 μm diam., smooth, hyaline, branched, clamped. **PILEIPELLIS** hyphae 3–4 μm diam., thin-walled to thick-walled, smooth, clamped, forming fascicles arranged on the cuticle as a network, branches and tomentum. **GELATINOUS ZONE** composed of hyphae in 1.5–3 μm diam., colourless, lax and implicated, often with inflated nodes or clamp connections in the middle of hyphae; hyphal terminals occasionally visible, inflated, thin-walled, hyaline,




FIG. 1. *Hohenbuehelia olivacea* (HMJAU 7290) (bar = 10µm)
a: Basidiospores b: Basidia c: Cheilocystidia d: Metuloids
e: Pileipellis f: Gelatinous zone hyphae

multiseptate, cylindrical or moniliform. FLESH HYPHAE 4–6 µm diam., thin-walled, hyaline, smooth, sinuous.

HABIT—Caespitose on rotten-wood of a broad-leaf tree in mixed forest.

REMARKS— This species is similar to *H. reniformis* (G. Mey.) Singer in having a tomentose pileus and short, stout sterigmata but differs from the latter in its slate-olive tomentum on the pileus and the lack of pileocystidia. Moreover, the metuloids of the new species are sublanceolate or cyprinoid, with a sharp apex.

Acknowledgments

We thank Dr. T.H. Li of Guangdong Institute of Microbiology, China, and Dr. R.H. Petersen of Department of Biology, University of Tennessee, USA, for their presubmission reviews of the manuscript. Financial support for this work was provided by the National Natural Science Foundation of China (No. 30670049).

Literature cited

- Bi ZS, Zheng GY, Li TH. 1993. The macrofungus flora of China's Guangdong Province. The Chinese University Press: Hong Kong (China). 734 pp.
- Bi ZS, Li TH, Zhang WM, Song B. 1997. A preliminary agaric flora of Hainan Province. Guangdong Higher Education Press: Guangzhou (China). 388 pp. (in Chinese).
- Chang ST, Mao XL. 1995. Hong Kong mushrooms. The Chinese University of Hong Kong Press: Hong Kong (China). 540 pp. (in Chinese).
- Chang TZ, Chou WN, Wang YZ, Ju YM. 2001. Macrofungi of Taiwan. Nat. Agric. Count. Press: Taipei (China). 542 pp. (in Chinese).
- Corner EJH. 1994. On the agaric genera *Hohenbuehelia* and *Oudemansiella*. Part I: *Hohenbuehelia*. The Gardens' Bulletin Singapore 46: 1–47.
- He X. 1992. A new species and a new record of *Tricholomataceae* from China. Acta Mycologica Sinica 11: 18–22.
- Kirk PM, Cannon PF, David JC, Stalpers JA. 2001. Dictionary of the fungi. 9th ed. Wallingford (UK): CAB International. 655 pp.
- Li Y, Bau T. 2003. Mushrooms of Changbai Mountains. Science Press: Beijing (China). 362 pp. (in Chinese).
- Mao XL. 1998. Economic macrofungi. Beijing Science and Technology Press: Beijing (China). 762 pp. (in Chinese).
- Ridgway R. 1912. Colour standards and colour nomenclature. Published by Author: Washington, D.C. 53 pp.
- Singer R. 1951. The "Agaricales" (mushrooms) in modern taxonomy. Coll. Lilloa (Revista de Botanica). Tomo XXII. Universidad de Tucumán: Tucumán (Argentina). 832 pp.
- Schulzer von Muggenburg S, Kanitz A, Knapp JA. 1866. Die bisher bekannten Pflanzen Slavoniens. Verh. Zool.-bot. Ges. Wien 16: 1–172.
- Teng SC. 1963. Fungi of China. Science Press: Beijing (China). 896 pp. (in Chinese).
- Thorn RG, Barron GL. 1986. *Nematotonus* and the tribe *Resupinateae* in the Ontario, Canada. Mycotaxon 25: 321–453.