

MYCOTAXON

Volume 109, pp. 245–246

July–September 2009

Three nomenclatural corrections for species of *Hypocrea/Trichoderma*

ANDREW M. MINNIS¹, GARY J. SAMUELS² & PRISCILA CHAVERRI³

¹Drew.Minnis@ars.usda.gov²Gary.Samuels@ars.usda.gov

Systematic Mycology & Microbiology Laboratory

USDA-ARS, Rm. 304, B011A, 10300 Baltimore Avenue, Beltsville, MD 20705, USA

³pchaverr@umd.edu

Dept. of Plant Sciences and Landscape Architecture, University of Maryland
2112 Plant Sciences Building, College Park, MD 20742, USA

Abstract — Three hypocreoid taxa were found to be illegitimate later homonyms of previously described species. Hence, the replacement names, *Hypocrea straminella* (= *Hypocrea straminea* P. Chaverri & Samuels), *Trichoderma pseudocandidum* (= *Trichoderma candidum* P. Chaverri & Samuels), and *Trichoderma pseudonigrovirens* (= *Trichoderma nigrovirens* P. Chaverri & Samuels), are provided.

Key words — Ascomycota, Hypocreaceae, Hypocreales, nomen novum, nomenclature

Introduction

Hypocrea Fr. (anamorph *Trichoderma* Pers.) is a genus of ubiquitous fungi, some of which are exploited in commercial applications including biological control, industrial enzyme production, and bioremediation (Chaverri & Samuels 2003). In a previous systematic treatment of species with green ascospores, Chaverri & Samuels (2003) described several taxonomic novelties. Unfortunately, some of them were subsequently found to be illegitimate names according to the International Code of Botanical Nomenclature (McNeill et al. 2006), since they are later homonyms of previously described but somewhat obscure taxa. According to McNeill et al. (2006), a nomen novum is needed for each name, and we prefer to assign the following replacement names before the use of the illegitimate names has become prevalent in the literature.

Taxonomy

Hypocrea straminella P. Chaverri, Samuels & Minnis, **nom. nov.**

MYCOBANK MB 513342

= *Hypocrea straminea* P. Chaverri & Samuels, Studies in Mycology: 48: 86, 2003,
non *Hypocrea straminea* Petch, Ann. Roy. Bot. Gard. (Peradeniya) 7: 97, 1920.

The anamorph of this fungus is *Trichoderma stramineum* P. Chaverri & Samuels (Chaverri & Samuels 2003).

***Trichoderma pseudocandidum* P. Chaverri, Samuels & Minnis, nom. nov.**

MYCOBANK MB 513343

= *Trichoderma candidum* P. Chaverri & Samuels, Studies in Mycology 48: 40, 2003, non *Trichoderma candidum* Alb. & Schwein., Conspectus Fungorum in Lusatiae Superioris: 137, 1805.

The teleomorph of this fungus is *Hypocrea candida* P. Chaverri & Samuels (Chaverri & Samuels 2003).

***Trichoderma pseudonigrovirens* P. Chaverri, Samuels & Minnis, nom. nov.**

MYCOBANK MB 513344

= *Trichoderma nigrovirens* P. Chaverri & Samuels, Studies in Mycology 48: 78, 2003, non *Trichoderma nigrovirens* Goddard, Bot. Gaz. 56: 273, 1913, "nigro-virens".

The teleomorph of this fungus is *Hypocrea nigrovirens* P. Chaverri & Samuels (Chaverri & Samuels 2003).

Acknowledgments

We extend our gratitude to Dr. Walter Gams and Dr. Walter Jaklitsch for their reviews of this article.

Literature cited

- Albertini JB, Schweinitz LD von. 1805. Conspectus Fungorum in Lusatiae Superioris. Lipsiae, Sumtibus Kummerianis. 376 p.
- Chaverri P, Samuels GJ. 2003. *Hypocrea/Trichoderma* (Ascomycota, Hypocreales, Hypocreaceae): species with green ascospores. Studies in Mycology 48: 1–116.
- Goddard HN. 1913. Can fungi living in agricultural soil assimilate free nitrogen? Bot. Gaz. 56: 249–305.
- McNeill J, Barrie FR, Burdet HM, Demoulin V, Hawksworth DL, Marhold K, Nicolson DH, Prado J, Silva PC, Skog JE, Wiersema JH, Turland NJ (eds.). 2006. International Code of Botanical Nomenclature (Vienna Code): Adopted by the Seventeenth International Botanical Congress, Vienna, Austria, July 2005. A.R.G. Gantner Verlag, Ruggell. 568 p.
- Petch T. 1920. *Hypocreaceae Zeylanicae*. Ann. Roy. Bot. Gard. (Peradeniya) 7: 85–138.