

MYCOTAXON

Volume 109, pp. 301–305

July–September 2009

***Manoharachariella,* a new dematiaceous hyphomycetous genus from India**

G. BAGYANARAYANA^{1*}, N. KRISHNA RAO² & I.K. KUNWAR¹

*gbagyan@rediffmail.com

¹Department of Botany, Osmania University
Hyderabad 500 007, A.P., India

²Botany Department, PAS College
Pedanandipadu-522 235, A.P., India

Abstract — A new hitherto undescribed anamorphic dematiaceous hyphomycetous genus with monoblastic, integrated conidiogenous cells producing solitary, doliiiform, obpyriform, dictyoseptate, apiculate, conidia collected on dead twigs in a forest near Darakonda, Andhra Pradesh, India is described as *Manoharachariella* gen. nov.

Key words — macronematous, acroauxic, tiered, apiculus, *Manoharachariella lignicola*

During a survey of hyphomycetes of Andhra Pradesh a new anamorphic genus was collected on unidentified dead twigs in a forest near Darakonda, Distt. Visakhapatnam, Andhra Pradesh, India. It is characterized by the presence of macronematous, mononematous conidiophores with monoblastic, acroauxic, integrated conidiogenous cells producing solitary, doliiiform, obpyriform, dictyoseptate, apiculate conidia. A critical microscopic examination of the fungus and perusal of literature (Ellis 1971, 1976; Matsushima 1975, 1983, 1996; Carmichael et al. 1980, Castañeda 1986, Mercado-Sierra 1984, Mercado-Sierra et al. 1997, Vasanth Rao & de Hoog 1986) revealed it to be hitherto undescribed dematiaceous hyphomycetous genus.

***Manoharachariella* Bagyan., N.K. Rao & Kunwar gen. nov.**

MyCOBANK MB 512919

Coloniae effusae, tenues, pallide vel modice brunneae, villosae. Mycelio immersum. Hyphis ramosae, brunneae. Stroma, setae et hyphopodiis nullis. Conidiophora macronemata, mononemata, erecta, plerumque flexuosa, sparse septata, hyalina vel subhyalina. Cellulae conidiogenae integratae, acroauxicae, monoblasticae integratis, cylindraceus, hyalino-subhyalino. Conidiis solitaria, sicca, acrogena et acropyleurogena, doliformis vel obpyriformis, rostratis, dictyoseptatis, apiculatus, modice vel atro-brunnea, apice et basi hyalino vel pallide brunnea.

SPECIES TYPICA: *Manoharachariella lignicola* Bagyan., N.K.Rao & Kunwar

ETYMOLOGY: The new genus is named in honour of an eminent and reputed mycologist of India, Prof. C. Manoharachary, Department of Botany, Osmania University, Hyderabad, India.

Colonies effuse, thin, pale brown to mid brown, hairy, mycelium immersed, hyphae brown, smooth. Stroma none, setae and hyphopodia absent. Conidiophores macronematous, mononematous, sparsely branched, branches loosely fasciculate, arising laterally and apically from the immersed mycelium, erect, usually flexuous, septate, septa few, hyaline to sub-hyaline. Conidiogenous cells acroauxic, monoblastic, integrated, cylindrical, hyaline to sub-hyaline. Conidia solitary, dry, acrogenous and acropyleurogenous, simple, doliiform, obpyriform, ellipsoidal, apiculate, smooth, dictyoseptate, tiered, mid brown to dark brown to blackish brown, apical and basal tiers hyaline to subhyaline.

Manoharachariella lignicola Bagyan., N.K. Rao & Kunwar sp. nov.

FIGS. 1, 2

MYCOBANK MB 512919

Hyphae dense, aggregatae, subepidermalis, 3–4.5 μm lata. Conidiophora usque 35 μm longa, et 3–4.5 μm lata. Conidia 42.5–50.5 μm longa, 25–32 μm in medio saepe, atro-nigro-brunnea, septis obscures, cellulae apicalis unicellularibus, 3.2–5.5 μm longis.

HOLOTYPE: On unidentified twigs, Darakonda, Distt. Visakhapatnam, A.P., India, 3 Nov 1984, leg. N.K. Rao, IMI 296857.

ETYMOLOGY: The fungus colonized unidentified dead twigs hence lignicola.




FIG. 1. *Manoharachariella lignicola*. A. Young conidium. B. Mature conidium and conidiophore.




FIG. 2. *Manoharachariella lignicola*. A-C. Bar = 10 μm . D. Bar = 5 μm .

Colonies effuse, thin, pale brown to mid brown, hairy, spreading to few millimeters on the host substratum. Mycelium immersed, composed of groups of brown, branched, smooth, 3–4.5 μm thick hyphae present in the subepidermal region of the host substrate. Conidiophores macronematous, mononematous, sparsely branched, branches loosely fasciculate arising both laterally and apically from the immersed hyphal cells, erect, usually flexuous, septate, septa few, 1–3, hyaline, to sub-hyaline, up to 35 μm long and 3–4.5 μm thick. Conidiogenous cells monoblastic, integrated, indeterminate, terminal and lateral, cylindrical, hyaline to sub-hyaline, 2.5–3.5 μm long. Conidia solitary, dry, acrogenous and acropyleurogenous, simple, doliform, obpyriform,

ellipsoidal, apiculate, dictyoseptate, smooth, tiered, transverse septa 7–9, several longitudinal septa, young conidia uniform in colour, sub-hyaline to pale brown, mature conidia mid-brown to dark brown except 1–2 tier of cells at the apical and basal region which are sub-hyaline to pale brown, middle part of the conidium often dark blackish brown almost obscuring the septation, apiculus one celled, thick walled, hyaline to sub hyaline, 3.2–5.5 µm long, 3.5–5.5 µm wide, conidia 42.5–50.5 µm long, 25–32 µm wide at the broadest part.

The genus, *Manoharachariella* shows some resemblance with *Septosporium* Corda in general and with *Septosporium rostratum* M.B. Ellis (Ellis 1961) in particular in having dictyoseptate, beaked conidia but differs from them in the absence of setae, unbranched conidiophores and percurrent conidiogenous cells. It also shows resemblance in shape with *Xenosporium africanum* Piroz. (Deighton & Pirozynski 1966) and *X. boivinii* S. Hughes (Hughes 1978) but differs from them in producing monoblastic, acrogenous/acropleurogenous, obpyriform, apiculate, smooth and dictyoseptate conidia which are transverse and longitudinally septate. It also differs from *Xenosporium* Penz. & Sacc. in the absence of secondary conidia.

Manoharachariella is comparable with *Bioconiosporium* Bat. & J.L. Bezerra (Ellis 1976) in having solitary, dry, acropleurogenous, doliiform, dictyoseptate conidia, but differs from it in conidia being monoblastic with a single beak, whereas two protuberances are present in *Bioconiosporium*. *Manoharachariella* is slightly comparable to *Monodictys* S. Hughes (Ellis 1971) in having dictyoseptate, monoblastic, dry, acrogenous conidia but differs from it in conidia being apiculate and tiered. There is no other fungus that can accommodate the newly proposed taxon. The present fungus is unique in having effuse, thin colonies, immersed mycelium, sparsely branched conidiophores; monoblastic, acroauxic conidiogenous cells; doliiform, dictyoseptate, tiered, brown to dark brown, apiculate conidia having 1–2 tier of hyaline or sub hyaline cells at the apical and basal end. Hence it is proposed as *Manoharachariella* gen. nov.

Acknowledgements

The authors thank Prof. K.G. Mukerji, New Delhi and Dr. D.K. Agarwal, New Delhi for serving as pre-submission reviewers. The senior author (GB) expresses his grateful thanks to Dr. Uwe Braun, Martin Luther University, Germany for his kind help and to Dr. Paul Cannon and Ms. Ann Ansell, IMI, London for kindly permitting him to examine the type specimen and use the facilities at IMI. IKK is grateful to MoEF, New Delhi for financial support.

Literature cited

- Castaneda Ruiz RF. 1986. Deuteromycotina de Cuba Hyphomycetes – IV. Academia de Ciencias de Cuba. pp. 17.
- Carmichael JW, Kendrick WB, Connors IL, Sigler L. 1980. Genera of Hyphomycetes. Univ. Alberta Press. 386 pp.
- Deighton FC, Pirozynski KA. 1966. Microfungi II. Mycol. Pap. 105: 1–35.
- Ellis MB. 1961. Dematiaceous Hyphomycetes. III. Mycol. Pap. 79: 3–5.
- Ellis MB. 1971. Dematiaceous Hyphomycetes, CMI, Kew, UK. 608 pp.
- Ellis MB. 1976. More Dematiaceous Hyphomycetes. CMI, Kew, UK. 507 pp.
- Hughes SJ. 1978. New Zealand Fungi. 25. Miscellaneous species. N. Z. J. Bot. 16(3): 311–370.
- Matsushima T. 1975. Icones microfungorum, a Matsushima lectorum, Publ. by the author, Kobe, Japan. 209 pp.
- Matsushima T. 1983. Matsushima Mycological Memoirs. 3: 1–90.
- Matsushima T. 1996. Matsushima Mycological Memoirs. 9: 1–40.
- Mercado-Sierra Á. 1984. Hifomicetes demaciaceos de Sierra del Rosario, Cuba, La Habana, Cuba: Editorial Academia. 181 pp.
- Mercado-Sierra Á, Holubova-Jechova C, Mena Portales J. 1997. Hifomicetes Demaciáceos de Cuba. Enteroblásticos, Museo Regionale di Scienze Naturali, Torino. 388 pp.
- Vasantha Rao, de Hoog GS. 1986. New or critical hyphomycetes from India. Studies in Mycology. 28: 1–84.

