

MYCOTAXON

Volume 109, pp. 319–322

July–September 2009

***Ocellularia gyrostomoides* belongs to the genus *Schizoxylon* (*Stictidaceae*, *Ascomycota*)**

H. THORSTEN LUMBSCH

tlumbsch@fieldmuseum.org

Department of Botany, Field Museum of Natural History

1400 S. Lake Shore Drive, Chicago, IL 60605, USA

KHWANRUAN PAPONG

khwanruan.p@msu.ac.th

Department of Biology, Faculty of Science, Maharakham University

Kantarawichai, Maha Sarakham Province 44150, Thailand

Abstract — While revising thelotrematoid *Graphidaceae* in Australia, we studied the type specimen of *Ocellularia gyrostomoides* and found it to be a non-lichenized fungus belonging to *Stictidaceae*. Here we report that the species belongs to the genus *Schizoxylon* (*Stictidaceae*) and propose the new combination *Schizoxylon gyrostomoides*.

Key words — *Ostropales*, taxonomy, *Thelotremataceae*

Introduction

While revising thelotrematoid *Graphidaceae* in Australia (Mangold et al. 2009) we also studied the type of *Ocellularia gyrostomoides*, which is an Australian endemic that has been accepted in the recent checklist of Australian lichens (McCarthy 2006). The name has been used in herbaria for species that are currently placed in *Schizotrema* (Mangold et al. 2009, Rivas Plata et al. 2010). However, examination of the type of *O. gyrostomoides* revealed that the species does not belong to *Graphidaceae* (incl. *Thelotremataceae*) as currently circumscribed (Mangold et al. 2008), but is a non-lichenized fungus belonging to the genus *Schizoxylon* in *Stictidaceae*. The necessary combination is made here and a short description of the fungus is given below.

Material and methods

Hand sections were examined in water and Lugol's solutions using a ZEISS Axioscope 2 plus compound microscope. TLC was performed with solvent

system A and B' (Culberson 1972, Culberson & Johnson 1982). Type material from G was studied and material for comparison that is deposited in F.

The species

Schizoxylon gyrostomoides (Müll. Arg.) Lumbsch & Papong, **comb. nov.** FIG. 1

MYCOBANK MB 513489

Bas.: *Ocellularia gyrostomoides* Müll. Arg., Flora 71: 46 (1888); type:
Australia, Queensland, Daintree River, *Pentzke* (G, holotype).

APOTHECIA orbicular, at first immersed in the substrate, becoming erumpent, 0.2–0.7 mm diam.; disc urceolate, brown, plane, whitish-pruinose; split between disc and margin obvious, margin prominent, thick, dark grey, smooth, entire. Exciple cupulate, brown, interspersed with numerous large crystals, lateral paraphyses absent. Subhymenium hyaline, 15–20 μm high. Hypothecium brown, 20–25 μm high. Hymenium hyaline, 100–140 μm high, clear to sparsely interspersed with large oil droplets. Epihymenium red-brown, 10 μm high, with crystals; pigmentation dissolving in K. Paraphyses simple to sparingly branched apically, apically not thickened to slightly thickened, hyaline. Asci long-




FIGURE 1 *Schizoxylon gyrostomoides*, holotype (G) (A) Habitat; (B) Habit detail, showing urceolate ascomata; (C) Cross section through ascoma; (D) Ascus with ascospores; (E) Fusiform ascospores.

cylindrical, 80–120 µm long, 8-spored, I+ faintly blue. Ascospores hyaline, fusiform, transversally septate, with 25–40 loculi, not disarticulating, loculi angular, I–, 45–65 × 5–7.5 µm, cell walls thick. Conidiomata not seen.

CHEMISTRY. No secondary metabolites detected by TLC.

NOTES. The faintly amyloid asci and the structure of the hymenium, exciple and ascospores indicate that this fungus belongs to *Stictidaceae* (Gilenstam 1969; Wedin et al. 2005, 2006). The absence of lateral paraphyses, the presence of abundant crystals in the exciple, and the initially immersed ascomata place this species in the genus *Schizoxylon* (Sherwood 1977a, b). Within *Schizoxylon* this species is characterized by a brown exciple and non-disarticulating ascospores shorter than 100 µm. The most similar species is *S. pseudocyanosporum* Sherwood from pine wood in Pakistan, which, however, is readily distinguished by larger, elongate ascomata and slightly larger ascospores [60–75 µm long].

Acknowledgments

Philippe Clerc (Geneva) is thanked for sending the type material on loan for examination. Armin Mangold (Berlin) initially studied the material and suggested it belongs to *Stictidaceae* and not *Thelotremataceae*. John A. Elix (Canberra) and Patrick McCarthy (Canberra) are thanked for reviewing this manuscript. This study was supported by a NSF grant (DEB-0516116) to The Field Museum (PI: HTL) and a grant from Australian Biological Resources Study (ABRS) to HTL.

Literature cited

- Culberson CF. 1972. Improved conditions and new data for the identification of lichen products by a standardized thin-layer chromatographic method. *Journal of Chromatography* 72: 113–125.
- Culberson CF, Johnson A. 1982. Substitution of methyl tert.-butyl ether for diethyl ether in standardized thin-layer chromatographic method for lichen products. *Journal of Chromatography* 238: 438–487.
- Gilenstam G. 1969. Studies in the lichen genus *Conotrema*. *Arkiv för Botanik, Ser. 2*, 7: 149–179.
- Mangold A, Elix JA, Lumbsch HT. 2009. *Thelotremataceae*. *Flora of Australia* 57: 195–420.
- Mangold A, Martin MP, Lücking R, Lumbsch HT. 2008. Molecular phylogeny suggests synonymy of *Thelotremataceae* within *Graphidaceae* (*Ascomycota*: *Ostropales*). *Taxon* 57: 476–486.
- McCarthy PM. 2006. Checklist of the Lichens of Australia and its Island Territories, Australian Biological Resources Study, Canberra. Version 6. April 2006.
- Rivas Plata E, Lücking R, Sipman HJM, Mangold A, Lumbsch HT. 2010. A world-wide key to the thelotrematoid *Graphidaceae*, excluding the *Ocellularia-Myriotrema-Stegobolus* clade. *Lichenologist* 42: in press.
- Sherwood MA. 1977a. The Ostropalean Fungi. *Mycotaxon* 5: 1–277.
- Sherwood MA. 1977b. The Ostropalean Fungi II: *Schizoxylon*, with notes on *Stictis*, *Acarosporina*, *Cocopeziza*, and *Carestiella*. *Mycotaxon* 6: 215–260.
- Wedin M, Döring H, Gilenstam G. 2006. *Stictis* s. lat. (*Ostropales*, *Ascomycota*) in northern Scandinavia, with a key and notes on morphological variation in relation to lifestyle. *Mycological Research* 110: 773–789.

Wedin M, Döring H, Konberg K, Gilenstam G. 2005. Generic delimitations in the family *Stictidaceae* (*Ostropales*, *Ascomycota*): the *Stictis*-*Conotrema* problem. *Lichenologist* 37: 67–75.