

Fungi on higher plants of the upper limit of the alpine zone in Tian Shan

ANDRZEJ CHLEBICKI

A.Chlebicki@botany.pl

*Polish Academy of Sciences, W. Szafer Institute of Botany
Lubicz 46, PL-31-512 Kraków, Poland*

Abstract — Fifty-two taxa of microfungi were noted at the upper limit of closed vegetation in Zailijskij Alatau Mts. (Tian Shan) in Kazakhstan, of which only 30% were saprobic species. Among the common and rare species were seven newly named taxa, six species new to science and one new combination. The complete annotated species checklist is available at <http://www.mycotaxon.com/resources/weblists.html>.

Key words — parasite, saprotroph, distribution, taxonomy

Introduction

The current study aims to assess microfungal diversity at the upper alpine plant limit in Tian Shan. The highly variable upper limit depends on many different factors, with different limits set for individual plants, plant communities, and closed vegetation (Grabherr et al. 1995). I investigated the limit of closed vegetation. Some information on fungi of the western part of Tian Shan has already been reported by Schwartzman (1962), Akhmedova (1966), Vasyagina (1977), Korbonskaya (1951, 1954), Salieva et al. (2002), Raitviir (2004), Chlebicki (2002, 2003, 2006, 2009), Chlebicki & Aime (2006), Chlebická & Chlebicki (2007), and Gaffyorov (2005). New species discovered since my research of the Zailijskij Alatau Mts. began in 2005 are *Cyathicula brunneospora* and *Pirottaea atrofusca* (Chlebická & Chlebicki 2007), *Microbotryum adenopetalae* (Lutz et al. 2008), and *Protoventuria juniperina*, *Trichometasphaeria barriae*, and *Veronea thylacospermi* (Chlebicki 2009).

Materials and methods

STUDY AREA: The terminal glacier foreland of the Issyk valley in Zailijskij Alatau Mts. (Tian Shan) near Almaty in southern Kazakhstan was investigated. The study was conducted on the slope of a marginal moraine (inactive ground ca 300 m before the ice margin) of the uppermost small basin at the glacier front at 3436 m elev., N43°07'52.5" E77°30'25". A distinct limit of closed vegetation was present. The native plant habitats comprised initial soil partially covered by variously sized (1 cm–1 m diam) granite rocks. Material was collected from 16 permanent 2.5 × 2.5 m plots within a 10 × 10

m square (FIG. 1). Plants growing at the altitudinal limit belong to various growth forms such as cushion plants, mat forming forbs, rosette perennial plants, tussock graminoids, prostrate dwarf shrubs, and tiny bryophytes.

METHODS: Dried material was examined under a Nikon SMZ 1500 zoom stereo microscope and at magnifications of 1000× and 2000× under a Nikon Labophot 2 or Olympus BX-51 light microscope, with Nomarski contrast (DIC) occasionally used. Microscopical observations and measurements of freehand longitudinal ascocarp sections were made in water and 3% KOH mountants. Lugol's solution (IKI: 1% iodine, 3% KI in water), Melzer's reagent (MLZ), and 5% KOH were used to determine the apical ring reactions and character of setae. Gelatinous sheaths of free ascospores were observed in India ink. Materials are deposited at the W. Szafer Institute of Botany of Polish Academy of Sciences in Kraków (Poland) and the National Museum in Prague (Czech Republic).




FIG. 1. The Tian Shan 10 m × 10 m permanent survey plot with 16 2.5 × 2.5 m quadrats (1A-4A, 1B-4B, 1C-4C, 1D-4D).

Results

Microfungi included 31 ascomycete species representing *Barrmaelia*, *Cainia*, *Cistella*, *Comoclathris*, *Cyathicula*, *Davidiella*, *Didymosphaerella*, *Glomerella*, *Hysteropezizella*, *Keissleriella*, *Lachnellula*, *Lophodermium*, *Mytilinidium*, *Nectriella*, *Phaeosphaeria*, *Phomatospora*, *Pirottaea*, *Pleospora*, *Protoventuria*, *Scutellinia*, *Trichometasphaeria* and *Wettsteinina* (Pezizomycotina); three rust species in *Melampsora*, *Microbotryum*, and *Puccinia* (Pucciniomycotina); a single *Ustilago* species (*Ustilagomycotina*); three Agaricales species in *Calyptrella*, *Lagarobasidium* and *Typhula*; and 14 mitosporic species representing *Alternaria*, *Botrytis*, *Cladosporium*, *Fusarium*, *Heteropatella*, *Hymenella*, *Periconia*, *Phoma*, *Seimatosporium*, and *Veronaea*. Many host plants were abundant enough for fungus persistence. The presence of parasites on single plants (e.g., *Lachnellula arida* on *Juniperus sibirica*, *Melampsora epitea* and *Seimatosporium lichenicola* on *Salix alata*) indicates long-range dispersal. Other fungi had restricted distributions in spite of their host plants being common in the investigated

area (e.g., *Ustilago striiformis* on *Anthoxanthum alpinum*, *Microbotryum adenopetalae* on *Silene adenopetala*, *Puccinia saxifragae* on *Saxifraga cernua*, *Lagarobasidium detriticum* on *Carex griffithii*). The greatest number of fungal species was noted on tussock plants such as *Carex griffithii*, *Festuca coelestis*, and *Anthoxanthum alpinum*.

Discussion

It is very difficult to describe all fungi linked with single host plant, as competition and niche differentiation influence the composition of the fungal-plant association (Neubert et al. 2006). The upper limit of closed vegetation can be compared with Gotelli's (1991) metapopulation model, which shows the effect of propagule immigration during population size decrease. Chlebicki & Olejniczak (2007) noted that the number of fungal species on plants is directly proportional to the size of the host population. It is evident that host plants were accessible to fungus propagules originating from lower plant populations. The highest number of fungal species per host, however, were noted on the common tussock plants (TABLE 1).

TABLE 1. Number of fungi noted on investigated plants

PLANT SPECIES	NO. OF FUNGI	PLANT SPECIES	NO. OF FUNGI
1. <i>Anthoxanthum alpinum</i> Á. Löve & D. Löve	22	12. <i>Oxyria digyna</i> (L.) Hill	2
2. <i>Carex griffithii</i> Boott	28	13. <i>Pentaphylloides fruticosa</i> (L.) O. Schwarz	0
3. <i>Cerastium cerastoides</i> (L.) Britton	5	14. <i>Primula nivalis</i> Pall.	4
4. <i>Doronicum oblongifolium</i> DC.	0	15. <i>Pyrethrum karelinii</i> Krasch.	0
5. <i>Draba incurvata</i> A.N. Vassiljeva & Golosk.	3	16. <i>Salix alata</i> Kar. & Kir. ex Stscheegl.	3
6. <i>Draba oreades</i> Schrenk	5	17. <i>Saussurea</i> sp.	5
7. <i>Dryadanthè tetrandra</i> (Bunge) Juz.	0	18. <i>Saxifraga cernua</i> L.	5
8. <i>Erigeron</i> sp.	0	19. <i>Saxifraga oppositifolia</i> L.	0
9. <i>Festuca coelestis</i> (St.-Yves) V.I. Krecz. & Bobrov	18	20. <i>Silene adenopetala</i> Raikova	9
10. <i>Juniperus sibirica</i> Burgsd.	3	21. <i>Thylacospermum caespitosum</i> (Cambess.) Schischk.	11
11. <i>Leontopodium leontopodium</i> (DC.) Hand.-Mazz.	1	22. <i>Waldheimia tridactylites</i> Kar. & Kir.	7

Acknowledgments

I thank Sabine Huhndorf and Hans Otto Baral for expert review and Amy Rossman, Hans Otto Baral, Markéta Chlebická, and Zdenek Pouzar for identification assistance. I would also like to thank Lorelei Norvell and Shaun R. Pennycook for helpful comments.

Literature cited

- Akhmedova FG. 1966. Mikoflora Yugo-Zapadnykh otrogov Thian-Shanya. Autoref. dissertation kand. biol. nauk. Tashkent. 25 pp.
- Chlebicki A. 2002. Two cypericolous smut fungi (*Ustilaginomycetes*) from the Thian Shan and their biogeographic implications. *Mycotaxon* 83: 279–286.
- Chlebicki A. 2003. *Anthracoidea stenocarpae*, the orthographically correct name for a species described from Tian Shan. *Mycotaxon* 88: 469.
- Chlebicki A. 2006. Two species of smut fungi on *Polygonaceae* from Tian Shan, Kazakhstan. *Czech Mycol.* 58: 99–104.
- Chlebicki A. 2009. Fungi on higher plants of the upper limit of alpine zone: new species from Tian Shan. *Mycotaxon* 110: 443–450.
- Chlebicki A, Aime C. 2006. New distribution and host records for *Puccinia atrofusca* and other notable *Puccinia* species (*Uredinales*) from Central Asia. *Polish Bot. J.* 51: 87–91.
- Chlebická M, Chlebicki A. 2007. *Cyathicula brunneospora* and *Pirotaeta atrofusca*, two new *Helotiales* from Tian Shan (Kazakhstan). *Mycotaxon* 100: 37–50.
- Chlebicki A, Olejniczak P. 2007. Symbiotic drift as a consequence of declining host populations. *Acta Biol. Crac. Ser. Bot.* 49(1): 89–93.
- Gaffyorov YS. 2005. Micromycetes on vascular plants in Namangan Region. Thesis of the PhD of Biol. Sci. Tashkent 2005.
- Gotelli JN. 1991. Metapopulation models: The rescue effect, the propagule rain, and the core-satellite hypothesis. *Amer. Naturalist* 38: 768–776.
- Grabherr G, Gottfried M, Gruber A, Pauli H. 1995. Patterns and current changes in alpine plant diversity. In: Chapin FS III, Körner Ch (Eds). *Arctic and alpine biodiversity: patterns, causes and ecosystem consequences*. Ecological Studies 113. Springer, Berlin, New York, p. 167–181.
- Korbonskaya JaI. 1951. Species novae *Uredinarum* a Republica Tadzhičica. *Bot. Mater. Otd. Sporov. Rast. Bot. Inst. Komarova Akad. Nauk S.S.S.R.* 7: 178–181.
- Korbonskaja JaI. 1954. Rzhavichnije griby Tadzhičikistana. *Trudy Akad. Nauk Tadzhičsk. SSR* 30: 1–95.
- Lutz M, Piątek M, Kemler M, Chlebicki A, Oberwinkler F. 2008. Anther smuts of *Caryophyllaceae*: molecular analyses reveal further new species. *Mycol. Res.* 112: 1280–1296.
- Neubert K, Mendgen K, Brinkmann H, Wirsal SGR. 2006. Only a few fungal species dominate highly diverse mycofloras associated with the common reed. *Appl. Envir. Microbiol.* 72(2): 1118–1128.
- Raitviir A. 2004. Revised synopsis of the *Hyaloscyphaceae*. *Scripta Mycol.* 20: 1–133.
- Salieva YaS, Kirgizbayeva KhM, Gulyamova M. 2002. Razpredeľennye mikromitsetov po rastitel'nyh poyasam otrogov Yugo-Zapadnogo Thian-Shanya. In: Bioobrazie Yugo-Zapadnogo Thian-Shanya: okhrana i ratzional'noye izpol'zovanie. International Conference. Tashkent, p. 191–194.
- Schwartzman SR. 1962. Flora sporovykh rastenij Kazakhstana 2. Golovnevye griby (Smut fungi). IAN Kazakhskoj SSR, Alma Ata. 369 pp.
- Vasyagina MP. 1977. Additions to the *Ustilaginales* flora of Kazakhstan. *Bot. Mater. Gerb. Bot. Akad. Nauk Kazakhsk. SSR* 10: 101–105.