

***Spadicoides subsphaerica* sp. nov. from Connecticut**

DE-WEI LI

dewei.li@ct.gov

The Connecticut Agricultural Experiment Station, Valley Laboratory
153 Cook Hill Road, Windsor, CT 06095

Abstract — *Spadicoides subsphaerica*, a new dematiaceous hyphomycete species is described and illustrated from a specimen collected from Hamden, CT.

Key Words — anamorphic fungi, saprobe, taxonomy

Introduction

The genus *Spadicoides* was erected by Hughes (1958), as typified by *S. bina* (Corda) S. Hughes [as '*binum*'] with six species accepted. The genus is characterized by distinct, single, unbranched, brown conidiophores with polytretic, integrated conidiogenous cells, solitary apical and lateral conidia, and minute pores visible on conidiogenous cells where conidia have developed and been released (Hughes 1958, Ellis 1971, Sinclair & Bhat 1985, Goh & Hyde 1996). Goh and Hyde (1996) reviewed the genus and accepted 21 species. Since the review, several additional species have been proposed: *Spadicoides arengae*, *S. bambusicola*, *S. hodgkissii*, *S. mauritiana*, *S. minuta*, *S. palmicola*, and *S. versiseptatis* (Cai et al. 2004, Dulyamamode et al. 1999, Goh & Hyde 1999, Ho et al. 2002, Wong et al. 2002, Zhou et al. 1999).

A specimen collected from dead wood at the Lockwood Farm of The Connecticut Agricultural Experiment Station in Hamden, CT was found to be of an undescribed species of *Spadicoides*, which is described and illustrated in this paper.

Materials and methods

Conidiophores and conidia of the fungus were lifted with 2 × 3 mm transparent tape 600 (3M Co., St. Paul, MN) and mounted in lacto-fuchsin (0.1 g acid fuchsin, 100 ml 85% lactic acid) or 85% lactic acid (Carmichael 1955). Microscopic observations were made using bright field and Nomarski differential interference contrast optics. Photomicrographs were taken with an

Olympus Microfire digital camera (Goleta, CA). Herbarium acronyms follow Index Herbariorum (Holmgren & Holmgren 1998).

Results

Spadicoides subsphaerica D.W. Li, anam. sp. nov.

FIGURES 1–8

MYCOBANK MB515400

Conidiophora macronemata, mononematica, solitaria, determinata, erecta, simplicia, non-ramosa, 39–77(105) μm longa, (2.5–)2.7–3.1(–3.2) μm lata, 3–7-septata, laevia, brunnea; cellulae conidiogenae polytreticae, terminales et intercalares, integratae, poris manifestis praeditae. Conidia unicellularia, solitaria, subsphaerica vel ellipsoidea, brunnea, laevia, (3.3–)3.8–4.6(–5.4) × (3.2–)3.5–4.1(–4.3) μm. Teleomorphosis ignota.

TYPE: UNITED STATES. CONNECTICUT, Hamden, Lockwood Farm, superficie in ligno. Coll. 5 viii 2009, BPI 879604 (holotype).

ETYMOLOGY: referring to the subglobose conidial shape.

Conidiophores differentiated, single, determinate, erect, unbranched, straight, dark brown, smooth, 3–7-septate, thick-walled, (38–)39–77(–105) (mean = 58 ± 19, n = 20) × (2.5–)2.7–3.1(–3.2) μm (mean = 2.9 ± 0.2, n = 20), more or less uniform in width, occasionally slightly enlarged at apex, upper half fertile and paler; apical cell (4.2–)6.4–9.0(–10) (mean = 7.7 ± 1.3, n = 20) × (2.5–)2.8–3.2(–3.5) μm (mean = 3.0 ± 0.2, n = 20). Conidiogenous cells integrated, terminal and intercalary, polytretic, leaving visible minute clear pores after conidial secession. Conidia apical and lateral, unicellular, single, subglobose, globose, or broadly ellipsoidal, brown to dark brown, smooth, thick-walled, with an occasionally visible, minutely protuberant clear pore at the base (3.3–)3.8–4.6(–5.4) (mean = 4.2 ± 0.4, n = 30) × (3.2–)3.5–4.1(–4.3) (mean = 3.8 ± 0.3, n = 30) μm, Q = 1–1.2(–1.5) (mean = 1.1 ± 0.1, n = 30).

TELEOMORPH: unknown.

KNOWN GEOGRAPHICAL DISTRIBUTION: Connecticut, USA.


HABITAT: saprotrophic on dead wood of ?*Quercus* sp.

Discussion

Several *Spadicoides* species develop 1-celled conidia: *S. arengae* W.H. Ho et al. ex L. Cai et al., *S. atra* (Corda) S. Hughes, *S. cuneata* Kuthub. & Nawawi, *S. macrocontinua* Matsush., *S. minuta* L. Cai et al., *S. sphaerosperma* McKenzie,

FIGURES 1–8. *Spadicoides subsphaerica*. 1–2. Conidiophore and conidia. 3. Apical portion of a conidiophore with lateral conidia. 4. Conidiogenous pores shown by arrows. 5. Conidium at apex of a conidiophore. 6–7. Thick-walled conidia. 8. Conidia and apical portion of a conidiophore.

Scale bars: 1–2 = 10 μm, 3–8 = 5 μm.



and *S. verrucosa* V. Rao & de Hoog. *Spadicoides subsphaerica* is characterized by small conidia ($3.3\text{--}5.4 \times 3.2\text{--}4.3 \mu\text{m}$), which are brown to dark brown, subspherical or ellipsoidal, and smooth, one or more of which characters separate it from the other species. Species that develop much larger and differently shaped conidia include *S. arengae* ($11\text{--}18 \times 4\text{--}6 \mu\text{m}$, ellipsoid), *S. cuneata* ($9\text{--}12 \times 6\text{--}8 \mu\text{m}$, cuneiform), *S. macrocontinua* ($13.5\text{--}22 \times 7\text{--}9 \mu\text{m}$, obovoid), and *S. sphaerosperma* ($6\text{--}7 \mu\text{m}$, globose) (McKenzie 1982, Matsushima 1995, Goh & Hyde, 1996, Dulyamamode et al. 1999, Ho et al. 2002). *Spadicoides atra*, *S. minuta*, and *S. verrucosa* develop conidia that overlap those of *S. subsphaerica* in size. However, *S. verrucosa* has verrucose, ellipsoidal conidia $4\text{--}5.5 \times 2\text{--}3 \mu\text{m}$ (Goh & Hyde 1996), and the conidia of *S. atra* are oblong, ellipsoidal to obovoid, and $4\text{--}6.5 \times 3\text{--}4 \mu\text{m}$ (Corda 1840, Matsushima 1975). *Spadicoides minuta* has subhyaline to hyaline, ellipsoidal to broadly ellipsoidal conidia with mucronate ends that measure $3\text{--}6 \times 2.5\text{--}3.5 \mu\text{m}$.

Acknowledgments

The author expresses his gratitude to Dr. Bryce Kendrick and Dr. Rafael F. Castañeda Ruiz for their critical review of the manuscript and suggestions for revision. The author is grateful to Dr. James A. LaMondia for his pre-submission review and Dr. Shaun Pennycook for his nomenclature review.

Literature cited

- Carmichael JW. 1955. Lacto-fuchsin: a new medium for mounting fungi. *Mycologia* 47: 611.
- Corda ACJ. 1840. *Icones Fungorum hucusque Cognitorum* 4: i-iii, 1-53, plates 1-10.
- Cai L, McKenzie EHC, Hyde KD. 2004. New species of *Cordana* and *Spadicoides* from decaying bamboo culms in China. *Sydowia* 56: 222-228.
- Dulyamamode R, Kirk PM, Peeraly A. 1999. Fungi from Mauritius: three new hyphomycete species on endemic plants. *Mycotaxon* 73: 313-323.
- Ellis MB. 1971. Dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey, United Kingdom, 608 pp.
- Goh T-K, Hyde KD. 1996. *Spadicoides cordanoides* sp. nov., a new dematiaceous hyphomycete from submerged wood in Australia, with a taxonomic review of the genus. *Mycologia*, 88: 1022-1031.
- Goh T-K, Hyde KD. 1999 ("1998"). *Spadicoides palmicola* sp. nov. on *Licuala* sp. from Brunei, and a note on *Spadicoides heterocolorata* comb. nov. *Can. J. Bot.* 76: 1698-1702.
- Ho WH, Yanna, Hyde KD. 2002. Two new species of *Spadicoides* from Brunei and Hong Kong. *Mycologia* 94: 302-306.
- Holmgren PK, Holmgren NH. 1998. [continuously updated]. Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/>
- Hughes SJ. 1958. Revisiones Hyphomycetum aliquot cum appendice de nominibus rejiciendis. *Can. J. Bot.* 36: 727-836.
- Matsushima T. 1975. *Icones microfungorum a Matsushima lectorum*. Published by the author, Kobe, Japan. 209 pp. + 415 plates

- Matsushima T. 1995. Matsushima mycological memoirs no. 8. Published by the author, Kobe, Japan. 54 pp. + 120 plates.
- McKenzie EHC. 1982. New hyphomycetes on monocotyledons. *New Zealand Journal of Botany* 20: 245–252.
- Sinclair RC, Eicker A, Bhat DJ. 1985. Branching in *Spadicoides*. *Trans. Brit. Mycol. Soc.* 85: 736–738.
- Wong MKM., Goh TK, McKenzie EHC, Hyde KD. 2002. Fungi on grasses and sedges: *Paratetraploa exappendiculata* gen. et sp. nov., *Petrakia paracochinensis* sp. nov. and *Spadicoides versiseptatis* sp. nov. (dematiaceous hyphomycetes). *Cryptogamie Mycologie* 23: 195–203.
- Zhou DQ, Goh TK, Hyde KD, Vrijmoed LLP. 1999. A new species of *Spadicoides* and other hyphomycetes on bamboo from Hong Kong. *Fungal Diversity* 3: 179–185.

