

***Botryobasidium sassofratinoense* sp. nov.  
(*Cantharellales*, *Basidiomycota*) from Italy**

ANNAROSA BERNICCHIA<sup>1</sup>, GITTA LANGER<sup>2</sup> & SERGIO PÉREZ GORJÓN<sup>3</sup>

<sup>1</sup>*annarosa.bernicchia@unibo.it*

*Dipartimento di Scienze e Tecnologie Agroambientali, Patologia Vegetale  
Università degli Studi di Bologna, Via Fanin 42, 40127 Bologna, Italy*

<sup>2</sup>*gitta.langer@nw-fva.de*

*Nordwestdeutsche Forstliche Versuchsanstalt, Abt. B Waldschutz  
SG 3 Mykologie und Komplexerkrankungen  
Grätzelstr. 2, 37079 Göttingen, Germany*

<sup>3</sup>*spgorjon@usal.es*

*Departamento de Botánica, Universidad de Salamanca  
Ldo. Méndez Nieto s/n, 37007 Salamanca, Spain*

**Abstract** — *Botryobasidium sassofratinoense* is described as new cystidiate species from Riserva di Sasso Fratino, located in Emilia-Romagna Region in the northern part of Italy. A key to the cystidiate species of *Botryobasidium* is provided and the new species is compared with the most closely related species, *B. ansosum* and *B. pilosellum*.

**Key words** — taxonomy, wood-inhabiting fungi, *Abies alba*, Europe

## Introduction

During the revision of material present in HUBO herbarium, a nameless specimen fell into the hands of one of the authors (A.B.), and on its envelope was written: “This could be a nice species!” After a close look the collection turned out to represent a cystidiate *Botryobasidium* close to *B. ansosum* (H.S. Jacks. & D.P. Rogers) Parmasto and to *B. pilosellum* J. Erikss. but with some different microscopic characters. It seemed quite surprising to discover a new species in a reserve that had been so well investigated during the last 25 years (with 545 macrofungi recorded from that area; Bernicchia & Gorjón 2009), and where two other new species had been described so recently: *Fomitopsis labyrinthica* Bernicchia & Ryvar den (Bernicchia & Ryvar den 1996) and *Ceriporiopsis guidella* Bernicchia & Ryvar den (Bernicchia & Ryvar den 2003). What is even more intriguing is that all three new species were recorded along the same trail that goes from Pian del Pero to Rio Cullacce inside Sasso Fratino Natural Reserve.

The genus *Botryobasidium* Donk belongs to the fungal order *Cantharellales* (*Homobasidiomycetes*) and about 50 species are recognized (CBS 2009, Langer 1994, Parmasto et al. 2004). *Botryobasidium* is a saprobic genus with corticioid to hypochnoid resupinate basidiocarps and diagnostic basidia that are short, cylindrical or subcylindrical to suburniform with 2–8 sterigmata, and generally arranged in clusters. *Botryobasidium* shows clear relations with *Thanatephorus*, *Ceratobasidium*, and *Cejpomyces*, in which basidiospores generally grow by repetition; it is also related to *Sistotrema*, which differs in the nature of the hymenium (palisade in *Sistotrema* and clustered in *Botryobasidium*) and shape of basidia (urniform in *Sistotrema* and subcylindrical to suburniform in *Botryobasidium*). Molecular data have confirmed these relationships, placing *Botryobasidium* in the cantharelloid clade as a sister taxon of *Sistotrema*, *Cantharellus*, *Craterellus*, *Hydnum*, and *Clavulina* (Moncalvo et al. 2006).

### Material and methods

For light microscopic studies, samples were mounted in 3% potassium hydroxide (KOH) and Melzer's solution (IKI). Thirty basidiospores have been measured to calculate the mean length and width and the following abbreviations are used:  $L^*$  = mean spore length,  $W^*$  = mean spore width,  $Q^*$  = quotient of the spore length and width ( $L^*/W^*$ ). Specimens are deposited in HUBO.

### Species description

***Botryobasidium sassofratinoense* Bernicchia & G. Langer, sp. nov.** Figs. 1,2  
MYCOBANK MB 515722

*Carposomata annua, resupinata, tenuissima, effusa, levia vel hypochnoidea, subalbida vel cremeo-alba. Systema hypharum monomiticum: hyphae generativae erectae, hyalinae, fibulatae angulo recto ramosae; tenuitunicatae 5–8(–9) µm latae in subhymenio, leviter crassitunicatae, 10–12(–15) µm latae in subicolo. Cystidia rara, subcylindracea, pleraque tenuiter tunicata, 28–45(–55) µm longa et 5–7 µm lata. Basidia subcylindracea vel suburniformia, leviter constricta, cum fibula basali, (13–)18–25 µm longa et 6–8.5 µm lata, 6-sterigmatibus recurvis. Basidiosporae hyalinae, tenuiter tunicatae, leves, navicularum figura, saepe quaternibus se congregatae, (6–)7–8.5 µm longae, (3–)3.5–4.5 µm latae, inamyloideae.*

**HOLOTYPE:** *Italia, Forlì-Cesena, Riserva Integrale di Sasso Fratino, loc. Rio Cullacce, 950 m. leg. A. Bernicchia 27.09.2001, in ligno putrido Abies alba, coll. 7594 in herbario HUBO conservatus est. Isotypus in K.*

**ETYMOLOGY:** the specific epithet derives from the name of the collecting area, Riserva di Sasso Fratino.

**DESCRIPTION** — **BASIDIOMATA** annual, resupinate, effuse, very thin, smooth to arachnoid, whitish to pale ivory, 150–200 µm thick, margin indistinct and not differentiated. **HYPHAL SYSTEM** monomitic: generative hyphae clamped and smooth, thin-walled, branched at right angles, 5–8(–9) µm wide in




FIG. 1. *Botryobasidium sassofratinoense* (coll. A. Bernicchia 7594)  
Hymenial elements. Bar=10  $\mu$ m



FIG. 2. *Botryobasidium sassofratinoense* (coll. A. Bernicchia 7594)  
Basidia and basidiospores. Bar=10  $\mu$ m above, 5  $\mu$ m below

subhymenium, slightly thick-walled in subiculum, 10–12(–15)  $\mu$ m wide and slightly yellowish near to the substrate. CYSTIDIA tubular, not frequent, basally clamped, hyaline, smooth, thin-walled, apically obtuse, 28–45(–55)  $\times$  5–7  $\mu$ m, without additional septa (Figure 1). BASIDIA arranged in botryose clusters, suburniform to subcylindrical, slightly constricted, smooth, thin-walled, basally clamped, (13–)18–25  $\times$  6–8.5  $\mu$ m, usually with 6 slender sterigmata up to 2.5–3.5(–4)  $\mu$ m long (Figure 2). BASIDIOSPORES navicular (Figure 2), hyaline, smooth, thin-walled, mostly glued together in groups of four, IKI–, (6–)7–8.5  $\times$  (3–)3.5–4.5  $\mu$ m, with apiculus bent towards ventral side,  $L^* = 7.7$ ,  $W^* = 3.9$ ,  $Q^* = 1.97$ . Chlamydo spores and anamorph not found.

DISTRIBUTION AND ECOLOGY — Thus far *Botryobasidium sassofratinoense* is known only from the type locality growing on well-decayed *Abies alba* wood. The area is a 764.25 ha Natural Reserve located in the northern Apennines (43°49'59"N, 11°49'E). It has a humid oceanic to subcontinental bioclimate (Gonnelli & Bottacci 2009), where *Fagus sylvatica* is the dominant species with well preserved areas of *Abies alba*, *Acer pseudoplatanus*, *Ulmus glabra*, *Fraxinus excelsior*, *Taxus baccata*, *Quercus cerris*, *Q. petraea*, *Ostrya carpinifolia*, *Carpinus betulus*, and *Corylus avellana* the most significant.

REMARKS — *Botryobasidium sassofratinoense* belongs to the group of cystidiolate and clamped *Botryobasidium* species. Of these, *B. baicalinum* Kotir. & Ryvar den and *B. parvisetosum* Boidin & Gilles produce subglobose chlamydospores and very long, two-celled cystidia (Boidin & Gilles 1998, Kotiranta & Ryvar den 2007). *Botryobasidium grandinioides* Hallenb. has a grandinioid to aculeate basidiome with longer, densely encrusted cystidia. *Botryobasidium ansosum* is distinguished by longer, wider cystidia and basidiospores, and *B. pilosellum* has shorter basidiospores and longer encrusted cystidia. Moreover, the former two species and *B. sassofratinoense* also differ from *B. grandinioides* in the hypochnoid to arachnoid hymenophore. TABLE 1 compares some diagnostic characters of related species and *B. sassofratinoense*.

#### Key for the cystidiolate species of *Botryobasidium*

- 1a. With chlamydospores, two-celled cystidia ..... 2
- 1b. Without chlamydospores, cystidia different ..... 3
- 2a. Cystidia 50–90  $\mu\text{m}$  long, basidiospores  $7.2\text{--}9.2 \times 2.2\text{--}3 \mu\text{m}$  ..... *B. parvisetosum*
- 2b. Cystidia > 100  $\mu\text{m}$  long, basidiospores  $7\text{--}8 \times 2.7\text{--}3.4 \mu\text{m}$  ..... *B. baicalinum*
- 3a. Hyphae with clamps ..... 4
- 3b. Hyphae without clamps ..... 7
- 4a. Basidiome grandinioid to aculeate ..... *B. grandinioides*
- 4b. Basidiome smooth or hypochnoid to arachnoid ..... 5
- 5a. Hyphae/cystidia with incrustation, basidiospores  $(4.5\text{--})5\text{--}6 \times 2.5\text{--}3 \mu\text{m}$   
..... *B. pilosellum*
- 5b. Hyphae/cystidia without incrustation, basidiospores  $(6\text{--})7\text{--}10 \times (3\text{--})3.5\text{--}5 \mu\text{m}$  .. 6
- 6a. Cystidia  $50\text{--}120 \times 8\text{--}12 \mu\text{m}$ , basidiospores  $8\text{--}10 \times 4\text{--}5 \mu\text{m}$  ..... *B. ansosum*
- 6b. Cystidia  $28\text{--}45(55) \times 5\text{--}7 \mu\text{m}$ , basidiospores  $(6\text{--})7\text{--}8.5 \times (3\text{--})3.5\text{--}4.5 \mu\text{m}$   
..... *B. sassofratinoense*
- 7a. Basidia  $8\text{--}13 \times 5\text{--}6 \mu\text{m}$ , basidiospores  $4.5\text{--}7 \times 2\text{--}3 \mu\text{m}$  ..... *B. piliferum*
- 7b. Basidia and basidiospores larger ..... 8
- 8a. Basidia  $17\text{--}22 \times 9\text{--}10.5 \mu\text{m}$ , basidiospores  $10\text{--}11.5 \times 4\text{--}5.5 \mu\text{m}$  ..... *B. digitatum*
- 8b. Basidia  $15\text{--}18(20) \times 6\text{--}8 \mu\text{m}$ , basidiospores  $7.5\text{--}8.5 \times 3\text{--}4 \mu\text{m}$  ... *B. tubulicystidium*

TABLE 1. Comparison of related *Botryobasidium* species with *B. sassofratinoense* (all measures in  $\mu\text{m}$ ).

SPECIES	CLAMPS	CYSTIDIA	BASIDIA	BASIDIO- SPORES	HYMENOPHORE	ENCRUSTATION	DISTRIBUTION	SUBSTRATE
<i>ansosum</i>	constant	50–120 × 8–12	17–29 × 7–10	8–9(–10) × 4–5	hypochnoid to pilosa	none	North America	conifer wood, especially <i>Pinus, Picea</i>
<i>digitatum</i>	absent	60–125 × 6–9	17–22 × 9–10.5	10–11.5 × 4–5.5	pruinose to arachnoid	none	Panama	wood (burned)
<i>grandinioides</i>	constant	up to 200 × 9–10	15–16(–19) × 5.5–7	6–7(–9) × 3–4	hypochnoid to grandinioid	strong	Iran	wood
<i>piliferum</i>	absent	40–100 × 6–8	(8–)9–13 × 5–6	4.5–7 × 2–3	arachnoid to hypochnoid	none	Africa	wood
<i>pilosellum</i>	constant	65–100 × 4–9	15–20 × 5–6.5	(4.5–)5–6 × 2.5–3	hypochnoid	slight	Ukraine	conifer wood
<i>sassofratinoense</i>	constant	28–45(–55) × 5–7	(13–)18–25 × 6–8.5	(6–)7–8.5 × (3–)3.5–4.5	smooth to hypochnoid	none	Italy	conifer wood, <i>Abies alba</i>
<i>tubulicystidium</i>	absent	60–110 × 6–9	15–18(–20) × 6–8	7.5–8.5 × 3–4	arachnoid	none	Taiwan	wood

### Acknowledgments

We are grateful to Patrizia Tabaroni and Giovanni Consiglio (Italy) for the revision of the Latin diagnosis, the members of State Forest Corp of Pratovecchio and Badia Prataglia (Arezzo) who, over the years have guided and helped during forays in Riserva of Sasso Fratino, and Heikki Kotiranta (Finland) and Erast Parmasto (Estonia) for their critical reviews of the manuscript before submission.

### Literature cited

- Bernicchia A, Gorjón SP. 2009. La biodiversità fungina nella riserva naturale integrale di Sasso Fratino. 115–136, in Bottacci A (ed.). La Riserva naturale integrale di Sasso Fratino: 1959–2009. 50 anni di conservazione della biodiversità. CFS/UTB.
- Bernicchia A, Ryvarden L. 1996. Two new brown rot polypores from Italy. *Mycol. Helvetica* 8(2): 3–10.
- Bernicchia A, Ryvarden L. 2003. A new white-rot polypore from Italy. *Mycotaxon* 88: 219–224.
- Boidin J, Gilles G. 1998. Basidiomycètes *Aphyllophorales* de l'île de la Réunion X. Compléments aux genres traits antérieurement. *Bull. Soc. Mycol. France* 104: 59–72.
- CBS. 2009. *Aphyllophorales* database. [www.cbs.knaw.nl/databases/index.htm](http://www.cbs.knaw.nl/databases/index.htm).
- Gonnelli V, Bottacci A. 2009. Il clima di Sasso Fratino. 39–45, in Bottacci A (ed.). La Riserva naturale integrale di Sasso Fratino: 1959–2009. 50 anni di conservazione della biodiversità. CFS/UTB.
- Kotiranta H, Ryvarden L. 2007. *Botrybasidium baicalinum* sp. nova (*Aphyllophorales*, *Basidiomycetes*). *Ann. Bot. Fennici* 44: 293–297.
- Langer G. 1994. Die Gattung *Botrybasidium* Donk (*Corticaceae*, *Basidiomycetes*). *Bibliotheca Mycologica*, Band 158. J. Cramer. Berlin. Stuttgart.
- Moncalvo JM, Nilsson RH, Koster B, Dunham SM, Bernauer T, Matheny PB, Porter TM, Margaritescu S, Weiss M, Garnica S, Danell E, Langer G, Langer E, Larsson E, Larsson KH, Vilgalys R. 2007. The cantharelloid clade: dealing with incongruent gene trees and phylogenetic reconstruction methods. *Mycologia* 98: 937–948.
- Parmasto E, Nilsson RH, Larsson KH. 2004. Cortbase version 2. Extensive updates of a nomenclatural database for corticioid fungi (*Hymenomycetes*). *Phyloinformatics* 1: 5.

