

MYCOTAXON

Volume 112, pp. 143–151

April–June 2010

Three new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China

CHUNXIA LU^{1,2} & LIN GUO^{1*}

Ch.x.lu@hotmail.com & **guol@im.ac.cn*

¹*Key Laboratory of Systematic Mycology and Lichenology
Institute of Microbiology, Chinese Academy of Sciences
Beijing 100101, China*

²*Graduate University of Chinese Academy of Sciences
Beijing 100049, China*

Abstract — Three new species, *Septobasidium gaoligongense* and *S. euryae-groffii* on *Eurya groffii* associated with *Pinnaspis* spp. and *Septobasidium polygoni* on *Polygonum campanulatum* associated with *Pseudaulacaspis kuisiensis*, are described. They were collected from Gaoligong Mountains in Yunnan Province, China.

Key words — *Pucciniomycetes*, *Septobasidiales*, taxonomy

Previously, a new species of *Septobasidium* was found in Gaoligong Mountains of Yunnan province (Lu & Guo 2009b). From the same area an additional three new species are described as follows:

Septobasidium gaoligongense C.X. Lu & L. Guo, sp. nov.

FIGS. 1–6

MYCOBANK MB 516523

Basidiomata resupinata, 15–20 cm longa, 7.5–8 cm lata, cinnamomeo-brunnea, brunnea vel atrobrunnea, margine determinata, superficie laevia, maturitate fissurata, in sectione primum (260–)525–580 µm crassa, deinde 1360–5000 µm crassa. Subiculum brunneum, 30–50 µm crassum. Contextus 2–3-stratosus. Columnae hyalinae vel bruneolae, primum 190–430 µm longae, deinde 3000–4900 µm longae, 290–340 µm latae, ex hyphis 3–5 µm latis compositae. Hymenium hyalinum, 40–50 µm crassum. Basidia fusiformia, cylindrica vel leviter irregularia, recta vel leviter curvata, 4-cellularia, 17–26 × 4–7 µm, hyalina vel brunnea. Sine probasidio. Basidiosporae non visae. Haustoria ex hyphis irregulariter spiralibus constantia.

TYPE: On *Eurya groffii* Merr. (*Theaceae*): China, Yunnan, Gaoligong Mountains, Baoshan, Baihualin, alt. 1400 m, 8.VII.2009, T.G. Hou 17, HMAS 199577 (holotype), associated with *Pinnaspis* sp. (*Diaspididae*).

*corresponding author





FIG. 1. Basidia of *Septobasidium gaoligongense* (HMAS 199577, holotype).

Basidiomata on branches, resupinate, 15–20 cm long, 7.5–8 cm wide, cinnamon-brown, brown or dark brown; margin determinate; surface smooth at first, becoming cracked at maturity. In section (260–)525–580 μm thick in the young stage and 1360–5000 μm thick in the old stage. Subiculum 30–50 μm thick, brown. Pillars 190–430 μm high in the young stage, 3000–4900 μm high in the old stage, 290–340 μm wide, hyphae of pillars 3–5 μm thick, hyaline or brownish, forming 2–3 horizontal layers. Hymenium 40–50 μm thick, hyaline. Basidia arising directly from the hyphae, fusiform, cylindrical or slightly irregular, straight or slightly curved, 4-celled, 17–26 \times 4–7 μm , hyaline or brown, without a probasidial cell. Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.

REMARKS: Morphologically, *S. gaoligongense* is similar to *S. crinitum* (Fr.) Couch, but differs mainly in forming 2–3 horizontal hyphal layers, having smaller basidia (17–26 \times 4–7 μm vs 40–55 \times 8.4–10 μm), and lacking a top layer. *Septobasidium crinitum* has a thick top layer (100–200 μm high), and lacks horizontal layers.

FIGS. 2–6. *Septobasidium gaoligongense* (HMAS 199577, holotype). 2. Basidiomata on branches.
3. Basidia (arrows). 4. Haustoria. 5. Pillars. 6. Section of basidioma.



Septobasidium polygoni C.X. Lu & L. Guo, sp. nov.

FIGS. 7–13

MyCOBANK MB 516524

Basidiomata resupinata, 2–15.5 cm longa, 1–3 cm lata, alba, cinnamomeo-brunnea vel brunnea, margine determinata, superficie laevia, in vetustate separata, in sectione 390–1550 μm crassa. Subiculum hyalinum vel brunneum, 30–100 μm crassum. Columnae hyalinae vel brunneae, primum 50–80 μm altae, deinde 440 μm altae, 30–70 μm crassae vel hyphis laxe completae, interdum hyphae repullulantes, super hymenium stratum









FIG. 7. Probasidia and basidia of *Septobasidium polygoni* (HMAS 196488, holotype).

FIGS. 8–13. *Septobasidium polygoni* (HMAS 196488, holotype). 8. Basidiomata on branches.
9–10. Sections of basidiomata. 11. Probasidia. 12. Basidium (arrow). 13. Haustoria.



hypharum secundum 50–200 μm *altum formantes*. *Hymenium* 50–100 μm *crassum*, *unistratosum vel 2-stratosum*. *Probasidia subglobosa vel pyriformia*, 10–17 \times 10–15 μm , *subhyalina vel flavidobrunnea, persistentia*. *Basidia cylindrica, curvata, 4-cellularia*, 24.5–34 \times 7.5–10 μm , *hyalina or flavidobrunnea*. *Basidiosporae non visae*. *Haustoria ex hyphis irregulariter spiralibus constantia*.

TYPE: On *Polygonum campanulatum* Hook. f. (Polygonaceae): China, Yunnan, Gaoligong Mountains, Tengchong, alt. 2050 m, 5.IX.2008, S.H. He, Y.F. Zhu & L. Guo 2371, HMAS 196488 (holotype), associated with *Pseudaulacaspis kuisiuensis* (Diaspididae).

Basidiomata on stems and branches, resupinate, 2–15.5 cm long, 1–3 cm wide, white, cinnamon-brown or brown; margin determinate; surface smooth, peeling off in old stage. In section 390–1550 μm thick. Subiculum hyaline or brown, 30–100 μm thick. Pillars hyaline or brown, 50–80 μm high in young stage, up to 440 μm high in old stage, 30–70 μm wide, or loosely filled with 220–400 μm high hyphae, sometimes from hymenial layer the fungal hyphae renews growth to form a second hyphal layer, 50–200 μm high. Hymenial layer 50–100 μm thick, single or 2-stratose. Probasidia subglobose or pyriform, 10–17 \times 10–15 μm , subhyaline or pale yellowish brown; probasidial cell persistent after the formation of the basidia. Basidia cylindrical, curved, 4-celled, 24.5–34 \times 7.5–10 μm , hyaline or yellowish brown. Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.

REMARKS: Morphologically, *S. polygoni* is similar to *S. citricola* Sawada from which it differs in having tall pillars (up to 440 μm vs 84–126 μm), a thinner hymenium (50–100 μm vs 100–390 μm), and smaller basidia (24.5–34 \times 7.5–10 μm vs 50–65 \times 8.2–9.7 μm).

Septobasidium euryae-groffii C.X. Lu & L. Guo, sp. nov.

FIGS. 14–19

MyCOBANK MB 516525

Basidiomata resupinata, 5–16 cm longa, 4–11 cm lata, cinnamomeo-brunnea, brunnea vel castaneo-brunnea, margine determinata, superficie laevia et protuberantia, deinde fissurata, in sectione 1260–2620 μm crassa, 3–12-stratosa. Subiculum 40–50 μm crassum, brunneum. Columnae 40–100 altae, 50–165 μm latae, superne ramosae tunc strato hypharum 360–560 μm alto formantae, hyphae repullulantes tum duo strata horizontalia 130–180 μm alta formantes. Hymenium 50–60 μm crassum. Interdum super hymenium columnae secundae 60–110 μm altae formatae et strata hypharum 4-stratosa 810–1050 μm alta successive superposita. Hymenium denovo formatum 70–110 μm altum. Basidia cylindrica, recta vel leviter curvata, 4-cellularia, 20–45 \times 5–8 μm , hyalina or brunneola. Sterigmata conica, 2–3 μm longa. Sine probasidio. Basidiosporae non visae. Haustoria ex hyphis irregulariter spiralibus constantia.

TYPE: On *Eurya groffii* Merr. (Theaceae): China, Yunnan, Gaoligong Mountains, Baoshan, Baihualin, alt. 1400 m, 8.VII.2009, T.G. Hou 21, HMAS 199579 (holotype), associated with *Pinnaspis* sp. (Diaspididae).

Basidiomata on branches, resupinate, perennial, 5–16 cm long, 4–11 cm wide, cinnamon brown, brown or chestnut brown; margin determinate; surface

14








FIG. 14. Basidia of *Septobasidium euryae-groffii* (HMAS 199579, holotype).

smooth and protuberant, becoming cracked later. In section 1260–2620 μm thick, composed of 3–12 layers. Subiculum 40–50 μm thick, brown. Pillars 40–100 μm high, 50–165 μm wide, branched outwards to form a hyphal layer 360–560 μm high, the hyphae renewing to form two horizontal layers 130–180 μm high, forming a hymenial layer 50–60 μm thick at the upper, with closely packed parallel upright threads. Sometimes from the hymenium successively forming pillars 60–110 μm high, and 4 hyphal layers 810–1050 μm high. Hymenial layer renewing, up to 70–110 μm high. Basidia arising directly from the hyphae without a probasidial cell, cylindrical, straight or slightly curved, 4-celled, 20–45 \times 5–8 μm , hyaline or brownish. Sterigmata coniform, 2–3 μm long. Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.

REMARKS: *Septobasidium euryae-groffii* is similar to *S. henningsii* Pat., from which it differs in producing shorter pillars (40–110 μm vs 300–1100 μm) and shorter sterigmata (3–5 μm vs 14–34 μm). In addition, the basidioma surface of *S. euryae-groffii* is bumpy whereas that of *S. henningsii* is smooth. Another similar species, *S. thwaitesii* (Berk. & Broome) Pat., has curved basidia and probasidial cells.



Excluded species

Septobasidium parlatoriae Sawada, Rep. Dept. Agric. Govt. Res. Inst. Formosa. 51: 57, 1931.

A study of the type specimen of *S. parlatoriae*, borrowed from TAI, showed that no scale insects are present beneath the fungal hyphae. It is an anamorphic fungus.

To date, 23 species of *Septobasidium* have been reported in China (Sawada 1933, Couch 1938, Teng 1963, Tai 1979, Kirschner & Chen 2007, Lu & Guo 2009a, b,c, Lu et al. 2010), including the three species reported in this paper.

Acknowledgements

The authors would like to express their deep thanks to Drs Eric H.C. McKenzie (Auckland, New Zealand) and Shuanghui He (Beijing Forestry University) for serving as pre-submission reviewers, to Dr. Shaun Pennycook (Auckland, New Zealand) for nomenclatural review, to Prof. Jianyun Zhuang (Institute of Microbiology, Chinese Academy of Sciences) for Latin corrections, to Prof. Zhenyu Li (Institute of Botany, Chinese Academy of Sciences) for identifying the host plants, to Prof. Sanan Wu (Beijing Forestry University) for identifying the scale insects, to Mrs. Xiangfei Zhu for inking in line drawings, and to the Curator of Taiwan University Herbarium (TAI) for loan of a specimen. This study was supported by the National Natural Science Foundation of China (No. 30499340 and No. 30670005).

Literature cited

- Couch JN. 1938. The Genus *Septobasidium*. Univ. of North Carolina Press, Chapel Hill. 480 p.
- Kirschner R, Chen CJ. 2007. New reports of two hypophylloous *Septobasidium* species from Taiwan. Fung. Sci. 22(1,2): 39–46.
- Lu CX, Guo L. 2009a. *Septobasidium maesae* sp. nov. (*Septobasidiaceae*) from China. Mycotaxon 109: 103–106.
- Lu CX, Guo L. 2009b. Two new species of *Septobasidium* (*Septobasidiaceae*) from China. Mycotaxon 109: 477–482.
- Lu CX, Guo L. 2009c. *Septobasidium annulatum* sp. nov. (*Septobasidiaceae*) and *Septobasidium kameii* new to China. Mycotaxon 110: 239–245.
- Lu CX, Guo L, Wei JG, Li JB. 2010. Two new species of *Septobasidium* (*Septobasidiaceae*) from southern China. Mycotaxon 111: 269–271.
- Sawada K. 1933. Descriptive catalogue of the Formosan fungi. Part VI. Rep. Dept. Agric. Govt. Res. Inst. Formosa 61: 1–99.
- Tai FL. 1979. Sylloge Fungorum Sinicorum. Science Press, Beijing. 1527 p.
- Teng SC. 1963. Fungi of China. Science Press, Beijing. 808 p.

Figs. 15–19. *Septobasidium euryae-groffii* (HMAS 199579, holotype). 15. Basidiomata on branches. 16. Basidia (arrow). 17, 19. Sections of basidiomata. 18. Haustoria.

