

Symphaster ximeniae sp. nov.: a rare asterinaceous fungus from Brazil

JOSÉ LUIZ BEZERRA¹, ELISANDRO RICARDO DRECHSLER-SANTOS²,
JADERGUDSON PEREIRA^{1*} & LEONOR COSTA MAIA²

*jader@uesc.br

¹Deptº. de Ciências Agrárias e Ambientais, Universidade Estadual de Santa Cruz – UESC
Rod. Ilhéus-Itabuna, km 16, Ilhéus, BA, 45662-000, Brazil

²Deptº. de Micologia, Universidade Federal de Pernambuco – UFPE
Av. Prof. Nelson Chaves s/n, Recife, PE, 50670-420, Brazil

Abstract — A new asterinaceous fungus collected on *Ximenia americana* is described from Northeastern Brazil and named *Symphaster ximeniae*.

Key words — *Asterinaceae*, *Olcaceae*, systematics

Introduction

Among the 46 genera of *Asterinaceae* Hansf. recently listed (Kirk et al. 2008), *Symphaster* Theiss. & Syd. (Theissen & Sydow 1915: 217) has the smallest number of species. It comprises only two species: the type species *S. gesneriaceae* (Henn.) Theiss. & Syd. (basonym *Cocconia gesneriaceae* Henn.), and *S. areolata* (Doidge) Arx (basonym *Isipinga areolata* Doidge). The first species was observed in Rio de Janeiro, Brazil, by Hennings (1904: 91) on leaves of an unknown *Gesneriaceae* plant, and since then no other registers of specimens of this fungus have been made, indicating its rare condition. The second species was found on *Euclea natalensis* A. DC. (*Ebenaceae*) in South Africa (Doidge 1921: 15).

Many epiphytic fungi have been described on leaves of *Ximenia americana* L. (*Olcaceae*), mainly *Meliolales* (Viégas 1961, Silva & Minter 1995, Mendes et al. 1998), but no *Symphaster* species has been registered. Similarly, new *Asterina* species have been recorded in recent years (Hosagoudar et al. 2001a; Hofmann & Piepenbring 2008; Song 2003; Song & Li 2002, 2004; Song et al. 2003a,b, 2004), but no new *Symphaster* species.

During the past few decades, Müller & Arx (1962) and Arx & Müller (1975) added new information about *Symphaster* and in this century Hosagoudar et al.

(2001b) and Bezerra (2004) made new contributions. The family *Asterinaceae* has been well characterized by Müller & Arx (1962), Lutrell (1973), Arx & Müller (1975), Barr (1987), Hosagoudar et al. (2001b), and Bezerra (2004).

As occurs with other biotrophic pathogens in *Asterinaceae*, *Symphaster* species are apparently host specific. In this case, not only morphological characters but also the host plant may be useful to separate species. Considering the low number of records of the genus, however, host specificity should be confirmed. For Hofmann & Piepenbring (2008), induction of plant infection and DNA sequence data may help elucidate this question for this family.

During a survey of *Asterinaceae* in a tropical forest in Brazil, a fungus with characteristics of *Symphaster* was found and is now described as a new species.

Materials and methods

Leaves of *Ximenia americana* (local name: Limão; Ameixeira-do-Brasil) showing superficial black stromata of an asterinaceous fungus were collected in October 2006 in the “Reserva Ecológica de Dois Irmãos”, a remnant of Atlantic Rain Forest, in the municipality of Recife, State of Pernambuco, Brazil. The aspect of the colonies on the leaf was observed on a stereomicroscope and the adhesive transparent tape method was used to visualize hyphae and hyphopodia. Free hand sections and squash mounts stained with lactophenol cotton blue were used to study the morphology of the fungus under the light microscope. The structures were measured in water. An exsiccatum of the material was deposited in the mycological collection of URM Herbarium and Mycobank number for new species was cited.

Taxonomy

Symphaster ximeniae J.L. Bezerra, Drechsler-Santos & Jad. Pereira, *sp. nov.*


MYCOBANK MB512160

FIGS. 1–7

Coloniae epyphyllae vel amphigenae, densae, dispersae vel confluentes, 1–4 mm diam. *Hyphae* flexuosae, brunneae, septatae, hyphopodiatae, ramosae, cellulis 16–20 × 5–6 µm. *Hyphopodia* unicellularia, brunnea obovata vel oblonga, recta vel incurvata, alternata vel opposita, integra, cellulis, 8–13 × 5.5–7 µm. *Haustoriae* intraepidermale, hyalinae. *Thyriothecia* ad 60–200 µm diam, confluentes, multilocullata, rotunda vel irregulariter, stellato dehiscentes ad centre, margine crenata; asci 38–52 × 23–32 µm, octosporae, globosae, bitunicatae, sessilia; paraphysoides mucosae praeditae; ascosporae 21–25 × (6–)7–10(–14) µm, ellipsoideae, brunneae, bicellulatae, fortiter constrictae, submedianae septae, parietus glabrae vel leniter espinescentis.

TYPE: BRAZIL: Pernambuco, Recife, Reserva Ecológica de Dois Irmãos (08°00'39.1"S and 34°56'38.8"W, 10m alt.), 12.X.2006, leg. J.L. Bezerra and E.R. Drechsler-Santos, on living leaves of *Ximenia americana* (HOLOTYPE, URM 79224).

ETYMOLOGY: derived from the host genus *Ximenia*.



FIGS. 1–7. *Symphaster ximeniae*. 1–2. A leaf of *Ximenia americana* showing epiphyllous colonies. 3. Young ascomata with hyphopodiate hyphae (arrows) 4. Vertical section of ascoma. 5. Young bitunicate ascus. 6. Ascus with mature ascospores. 7. Ascospore with septum below the middle.

Scale bars: 1 = 5 mm; 2 = 1 mm; 3, 4 = 50 μ m; 5 = 20 μ m; 6 = 10 μ m; 7 = 5 μ m.

Colonies dull black, amphigenous, mostly epiphyllous, crustose, subcircular to irregular, isolate or confluent, scattered, 1–4 mm diam. Mycelium superficial of flexuous, brown, septate, hyphopodiate, oppositely or unilaterally branched, teleomorphic hyphae 16–20 \times 5–6 μ m. Hyphopodia unicellular, brown concolorous with the hyphae, obovoid to oblong or cylindrical, straight or curved, opposite or alternate, entire, 8–13 \times 5.5–7 μ m. Haustoria coralloid, hyaline, intra-epidermal. Ascomata dark brown, round to irregular, scutate, confluent, 60–200 μ m diam, forming stromatic multilocular crusts; upper wall,

opaque dark brown, 8–17 μm thick, formed of radiating rectangular cells, 6–12 \times 3–5 μm diam, opening by stellate dehiscence. Basal wall, 10–17 μm thick, formed by hyaline, thin walled hyphal cells. Paraphysoids numerous, in gelatinous mass, hyaline, filiform, septate, 2–3 μm diam. Asci 8-spored, globose to subglobose, sessile, thick walled, bitunicate, not bluing in Melzer's reagent, 38–52 \times 23–32 μm . Ascospores 1-septate below the middle, constricted in the septum, oblong, with rotund ends, brown at maturity, smooth to slightly rough, 21–25 \times (6–)7–10(–14) μm , with a larger apical cell.

NOTES: *Symphaster ximeniae* differs from *S. gesneriaceae* and *S. areolata* by possessing globose to subglobose asci and smaller ascospores, which are septate below the middle. *Symphaster areolata* and *S. gesneriaceae* differ from each other in ascospore size and type of hyphopodia. Authentic material of *S. areolata* (URM 23061 = PRE 22362) was examined, but no ascoma was seen. Each of the three *Symphaster* species occurs on a different host family.

Acknowledgments

The authors thank Maria de Fátima de Araújo and Prof. Marccus Alves (Departamento de Botânica/UFPE) for plant identification and gratefully acknowledge James W. Kimbrough and Francisco Das Chagas Oliveira Freire for pre-submission reviews. Thanks are also due to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação de Amparo à Pesquisa do Estado da Bahia (FAPESB) for financial support.

References

- Arx JA von, Müller E. 1975. A re-evaluation of the bitunicate ascomycetes with keys to families and genera. *Stud Mycol* 9: 1–160.
- Barr ME. 1987. *Prodromus to Class Loculoascomycetes*. Lubrecht & Cramer, Forestburg
- Bezerra JL. 2004. Taxonomia de Ascomycetos. Ordem Asteroales. *Revisão Anual de Patologia de Plantas* 11: 15–28.
- Doidge EM. 1921. South African ascomycetes in the National Herbarium I. *Bothalia* 1: 5–32.
- Hennings P. 1904. Fungi fluminenses a cl. E. Ule collecti. *Hedwigia* 43: 78–95.
- Hofmann TA, Piepenbring M. 2008. New species and records of *Asterina* from Panama. *Mycol. Progress* 7: 87–98.
- Hosagoudar VB, Abraham TK, Biju, CK, Hyde KD. 2001a. Fungi from palms. XLVII. A new species of *Asterina* on palms in India. *Fungal Diversity* 6: 69–73.
- Hosagoudar VB, Abraham TK, Biju CK. 2001b. Re-evaluation of the family *Asterinaceae*. *Journal of Mycopathological Research* 39: 61–63.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. *Ainsworth and Bisby's dictionary of the fungi*. 10th ed. CABI International. Wallingford, UK.
- Luttrell ES. 1973. *Loculoascomycetes*. In: Ainsworth GC, Sparrow FK, Sussman AS (eds.). *The Fungi*. vol. IV A. Academic Press, New York, pp 135–219.
- Mendes MAS, Silva VL, Dianese JC, et al. 1998. *Fungos em plantas no Brasil*. Embrapa-SPI/Embrapa-Cenargen, Brasília.

- Müller E, Arx JA von. 1962. Die Gattungen der didymosporen Pyrenomyceten. Beitr. Kryptogfl. Schwz 11(2). 922 p.
- Silva MS, Minter D. 1995. Fungi from Brazil recorded by Batista and co-workers. CAB International, Mycological Papers 169.
- Song B. 2003. New species of the genus *Asterina* from China III. Mycotaxon 85: 319–324.
- Song B, Li TH. 2002. New species of the genus *Asterina* from China, Mycotaxon 84: 407–412.
- Song B, Li TH. 2004. New species of *Asterina* in HMAS, China. Mycotaxon 89: 193–199.
- Song B, Li TH, Hosagoudar VB. 2003a. Four new *Asterina* species from Yunnan, China. Fungal Diversity 14: 157–164.
- Song B, Li TH, Shen YH. 2003b. Two new *Asterina* species from Hainan, China. Mycotaxon 87: 417 – 419.
- Song B, Li TH, Shen YH. 2004. New species of *Asterina* from Guangdong, China. Mycotaxon 90: 29–34.
- Theissen F, Sydow H. 1915. Die *Dothideales*. Kritisch-systematische Originaluntersuchungen, Ann. Mycol. 13: 149–746.
- Viégas AP. 1961. Índice de fungos da América do Sul. Instituto Agrônômico, Campinas.

