

MYCOTAXON

DOI: 10.5248/113.87

Volume 113, pp. 87–93

July–September 2010

**Two new species of *Septobasidium* (*Septobasidiaceae*)
and *S. pallidum* new to China**CHUNXIA LU^{1,2} & LIN GUO^{1*}

Ch.x.lu@hotmail.com & *guol@im.ac.cn

¹Key Laboratory of Systematic Mycology and Lichenology
Institute of Microbiology, Chinese Academy of Sciences
Beijing 100101, China²Graduate University of Chinese Academy of Sciences
Beijing 100049, China

Abstract — Two new species, *Septobasidium meridionale* on *Litsea cubeba* associated with *Aulacaspis* sp. and *S. aulacaspidis* on an unidentified tree associated with *Aulacaspis* sp., are described. *Septobasidium pallidum* on *Zanthoxylum bungeanum*, *Z. simulans*, and *Pyrus phaeocarpa* is new to China.

Key words — *Pucciniomycetes*, *Septobasidiales*, taxonomy

Previously, a new species of *Septobasidium* was found in Hainan province (Lu & Guo 2009a). In December 2009, many specimens of *Septobasidium* were collected from the same area. Among them, an additional two new species are described as follows:

***Septobasidium meridionale* C.X. Lu & L. Guo, sp. nov.**

Figs. 1–7

MYCOBANK MB 518060

Basidiomata resupinata, 4–8.5 cm longa, 1–6 cm lata, alba vel brunnea, margine determinata, superficie laevia vel vlutina, in sectione 840–1000 µm crassa. Subiculum brunneum vel hyalinum, 20–50 µm crassum. Columnae hyalinae vel brunneolae, 40–130 µm altae, 40–340 µm latae, ex hyphis 3–4 µm latis compositae, superne ramosae tunc strato hypharum 580–780 µm alto formatae, interdum strata horizontalia formantes. Hymenium 40–90 µm crassum, hyalinum vel brunneum. Basidia cylindrica, recta vel curvata, 4-cellularia, 27–36 × 7–9.5 µm, hyalina vel brunneola. Sine probasidio. Basidiosporae non visae. Haustoria ex hyphis irregulariter spiralibus constantia.

TYPE: On *Litsea cubeba* Pers. (*Lauraceae*): China, Hainan, Bawangling, Nanchahe, alt. 600 m, 11.XII.2009, Y.F. Zhu & L. Guo 128, HMAS 240076 (**holotype**), associated with *Aulacaspis* sp. (*Diaspididae*).

*corresponding author




FIG. 1. Basidia of *Septobasidium meridionale* (HMAS 240076, holotype).

Basidiomata on trunks and branches, resupinate, perennial, 4–8.5 cm long, 1–6 cm wide, white or brown; margin determinate; surface smooth or velutinous. In section 840–1000 μm thick. Subiculum 20–50 μm thick, brown or hyaline. Pillars hyaline or brownish, 40–130 μm high, 40–340 μm wide, hyphae of pillars 3–4 μm thick. Hyphal layer 580–780 μm high, sometimes forming a horizontal layer. Hymenium 40–90 μm thick, hyaline or brown. Basidia arising directly from the hyphae, cylindrical, straight or curved, 4-celled, 27–36 \times 7–9.5 μm , hyaline or brownish, without a probasidial cell. Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.

REMARKS: Morphologically, *Septobasidium meridionale* is similar to *S. septobasidioides* (Henn.) Höhn. & Litsch., from which it differs in having short pillars (40–130 μm vs 350–450 μm) and smaller basidia (27–36 \times 7–9.5 μm vs 40–55 \times 8.4–10 μm), and sometimes forming a horizontal layer.






***Septobasidium aulacaspidis* C.X. Lu & L. Guo, sp. nov.**

FIGS. 8–12

MYCOBANK MB 518061

Basidiomata resupinata, 0.2–7 cm longa, 0.1–5 cm lata, alba vel cinnamomeo-brunnea, margine indeterminata, superficie laevia vel velutina, in sectione 360–550 μm crassa. Subiculum brunneum, 30–50 μm crassum. Columnae hyalinae vel brunneolae, 200–260 μm altae, 20–120 μm latae. Hymenium 50–80 μm crassum. Basidia cylindrica, recta vel curvata, 4-cellularia, 28–50 \times 5–10 μm , hyalina vel brunnea. Basidiosporae ovoideae vel reniformes, 10–16 \times 4–5.5 μm . Sine probasidio. Haustoria ex hyphis irregulariter spiralibus constantia.

FIGS. 2–7. *Septobasidium meridionale* (HMAS 240076, holotype). 2. Basidiomata on trunk. 3–4. Sections of basidiomata. 5–6. Basidia (arrows). 7. Haustoria.






FIG. 8. Basidia of *Septobasidium aulacspidis* (HMAS 240074, holotype).








TYPE: On unidentified tree [probably *Neolitsea* sp. (*Lauraceae*)]: China, Hainan, Jianfengling, alt. 900 m, 11.XII.2009, S.H. He 2803, HMAS 240074 (**holotype**), associated with *Aulacaspis* sp. (*Diaspididae*).

Basidiomata on trunks, resupinate, subcircular or irregular, often confluent 0.2–7 cm long, 0.1–5 cm wide, white or cinnamon-brown; margin indeterminate; surface smooth or velvety. In section 360–550 μm thick. Subiculum 30–50 μm thick, brown. Pillars 200–260 μm high, 20–120 μm wide, hyaline or brownish. Hymenium 50–80 μm thick, with irregularly arranged upright hymenial hyphae. Basidia arising directly from the hyphae, cylindrical, straight or curved, 4-celled, 28–50 \times 5–10 μm , hyaline or brown, without a probasidial cell. Sterigmata conical, 5–13 \times 2 μm . Basidiospores ovoid or reniform, 10–16 \times 4–5.5 μm , pale yellowish brown. Haustoria consisting of irregularly coiled hyphae.

REMARKS: *Septobasidium aulacspidis* is similar to *S. pallidum*, but differs mainly in having indeterminate margin, smooth and velvety surfaces of basidiomata, and tall pillars (200–260 μm vs 84 μm). *Septobasidium pallidum* has determinate margin, non-velvety surface of basidioma, and short pillars.

Recently, several specimens of a *Septobasidium* sp. on *Zanthoxylum bungeanum* and *Pyrus phaeocarpa* were collected in Sichuan province. They were identical to a specimen of *Septobasidium* sp. on *Zanthoxylum simulans* previously deposited in our herbarium. No basidia were found in the specimen. The fungus is identified as *S. pallidum*, a species unrecorded previously in China:

FIGS. 9–12. *Septobasidium aulacspidis* (HMAS 240074, holotype). 9. Basidiomata on trunk. 10. Section of basidioma. 11. Basidia (arrows). 12. Haustoria. FIGS. 13–16. *Septobasidium pallidum* (HMAS 199578). 13. Basidiomata on trunk. 14. Section of basidioma. 15. Basidium (arrow). 16. Haustoria.






FIG. 17. Basidia of *Septobasidium pallidum* (HMAS 199578).

Septobasidium pallidum Couch ex L.D. Gómez & Henk, Lankesteriana 4(1): 88, 2004. FIGS. 13–17

Basidiomata on trunks and branches, resupinate, subcircular, 0.2–7 cm long, 0.2–3 cm wide, patches frequently confluent, tiller buff, yellowish-brown or brown; surface often smooth, with mounds and wrinkles, sometimes cracked; margin determinate, white. In section 220–510(–720) μm thick. Subiculum 20–60 μm thick. Pillars 40–90 μm high, 20–140 μm thick. Hyphal layer 100–400(–600) μm high. Hymenium 50–100 μm thick. Basidia arising directly from the hyphae without a probasidial cell, cylindrical, straight or slightly curved, 4-celled, 17–38(–42) \times 6–12 μm , hyaline or brown. Sterigmata 12–27 μm long, 2–3 μm wide. Hymenial hyphae irregularly arranged. Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.

SPECIMENS EXAMINED: On *Zanthoxylum bungeanum* Maxim. (*Rutaceae*): China, Sichuan, Jinyang, alt. 1100 m, 15.VIII.2009, S.H. He, C.X. Lu, Y.F. Zhu & L. Guo 2680, HMAS 199578; Jinyang, alt. 600 m, 15.VIII.2009, S.H. He, C.X. Lu, Y.F. Zhu & L. Guo 2686, HMAS 196491; Jinyang, Wuke, alt. 1100 m, 15.VIII.2009, S.H. He, C.X. Lu, Y.F. Zhu & L. Guo 2681, HMAS 199582; Jinyang, Mufu, alt. 1600 m, 15.VIII.2009, S.H. He, C.X. Lu, Y.F. Zhu & L. Guo 2685, HMAS 196487; Mianning, Manshuiwan, Ganghe, alt. 1740 m, 24.VIII.2009, S.H. He, C.X. Lu, Y.F. Zhu & L. Guo 2789, HMAS 196493; Xide, Mianshan, Dengxiangying, alt. 2300 m, 23.VIII.2009, S.H. He, C.X. Lu, Y.F. Zhu & L. Guo 2781, HMAS 196492.

On *Zanthoxylum simulans* Hance (*Rutaceae*): China, Sichuan, Hanyuan, 14.XII.1937, Y. Hu, HMAS 10165.

On *Pyrus phaeocarpa* Rehder (*Rosaceae*): China, Sichuan, Xide, Tanshan, alt. 1860 m, 23.VIII.2009, S.H. He, C.X. Lu, Y.F. Zhu & L. Guo 2784, HMAS 199628.

To date, 26 species of *Septobasidium* have been reported in China (Sawada 1933, Couch 1938, Teng 1963, Tai 1979, Kirschner & Chen 2007, Lu & Guo 2009a,b,c, 2010, Lu et al. 2010) including the three species reported in this paper.

Acknowledgements

The authors would like to express their deep thanks to Dr Eric H.C. McKenzie (Auckland, New Zealand) for serving as pre-submission reviewer, to Dr Shuanghui He (Beijing Forestry University) for serving as pre-submission reviewer and sending a specimen, to Prof. Jianyun Zhuang (Institute of Microbiology, Chinese Academy of Sciences) for Latin corrections, to Mr Ziyu Cao (Institute of Botany, Chinese Academy of Sciences) for identifying the host plants, to Prof. Sanan Wu (Beijing Forestry University) for identifying the scale insects, and to Mrs Xiangfei Zhu for inking in line drawings. This study was supported by the National Natural Science Foundation of China (No. 30499340 and No. 30870016) and the Ministry of Science and Technology of the People's Republic of China (No. 2006FY110500–5).

Literature cited

- Couch JN. 1938. The Genus *Septobasidium*. Univ. of North Carolina Press, Chapel Hill. 480 p.
- Kirschner R, Chen CJ. 2007. New reports of two hypophyllous *Septobasidium* species from Taiwan. *Fung. Sci.* 22: 39–46.
- Lu CX, Guo L. 2009a. *Septobasidium maesae* sp. nov. (*Septobasidiaceae*) from China. *Mycotaxon* 109: 103–106.
- Lu CX, Guo L. 2009b. Two new species of *Septobasidium* (*Septobasidiaceae*) from China. *Mycotaxon* 109: 477–482.
- Lu CX, Guo L. 2009c. *Septobasidium annulatum* sp. nov. (*Septobasidiaceae*) and *S. kameii* new to China. *Mycotaxon* 110: 239–245.
- Lu CX, Guo L. 2010. Three new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. *Mycotaxon* 112: 143–151.
- Lu CX, Guo L, Wei JG, Li JB. 2010. Two new species of *Septobasidium* (*Septobasidiaceae*) from southern China. *Mycotaxon* 111: 269–274.
- Sawada K. 1933. Descriptive catalogue of the Formosan fungi. Part VI. Rep. Dept. Agric. Govt. Res. Inst. Formosa 61: 1–99.
- Tai FL. 1979. *Sylloge Fungorum Sinicorum*. Science Press, Beijing. 1527 p.
- Teng SC. 1963. *Fungi of China*. Science Press, Beijing. 808 p.

