

MYCOTAXON

DOI: 10.5248/114.423

Volume 114, pp. 423–428

October–December 2010

New records of *Corynesporopsis* from ChinaJIAN MA, SHOU-CAI REN, LI-GUO MA,
YI-DONG ZHANG & XIU-GUO ZHANG*

zhxg@sdau.edu.cn, sdau613@163.com

Department of Plant Pathology, Shandong Agricultural University
Taian, 271018, China

Abstract — Three species of *Corynesporopsis* — *C. uniseptata*, *C. quercicola*, and *C. indica* — are recorded for the first time from China. They are described and illustrated from specimens collected on dead branches of unidentified plants. The specimens are deposited in Herbarium of Shandong Agricultural University, Plant Pathology (HSAUP) and Mycological Herbarium, Institute of Microbiology, Chinese Academy of Sciences (HMAS).

Key words — hyphomycetes, taxonomy

Introduction

Kirk (1981a) established the genus *Corynesporopsis* for the single species previously known as *Corynespora quercicola*. It is characterized by single, differentiated, sometimes percurrent, conidiophores and integrated, terminal, monotretic conidiogenous cells that produce short acropetal chains of ellipsoid to cylindrical euseptate conidia. These characters separate *Corynesporopsis* P.M. Kirk from other similar genera including *Corynespora* Güssow (Güssow 1906), *Corynesporella* Munjal & H.S. Gill (Munjal & Gill 1961), *Hemicorynespora* M.B. Ellis (Ellis 1972), and *Solicorynespora* R.F. Castañeda & W.B. Kendr. (Castañeda & Kendrick 1990). Nine species are currently included in this genus, of which *Corynesporopsis quercicola* and *C. biseptata* (M.B. Ellis) Morgan-Jones were transferred from *Corynespora* (Kirk 1981a,b, 1983, Holubová-Jechová & Mercado 1986, Holubová-Jechová 1987, Morgan-Jones 1988, Sutton 1989, Castañeda Ruíz & Kendrick 1990, Matsushima 1993). Only *Corynesporopsis isabelicae* Hol.-Jech. has previously been reported from China (Lu et al. 2000). Most species are reported to survive saprophytically on dead branches, twigs,

*Corresponding author

and decaying leaves of various plants. During a continuing survey of tropical microfungi from the forests of Hainan Province of southern China, three species of *Corynesporopsis* were collected on dead branches. They are introduced as new records for China.

Taxonomy

Corynesporopsis uniseptata P.M. Kirk, Trans. Br. Mycol. Soc. 77(3):
463 (1981)

FIG. 1

SPECIMEN EXAMINED: on dead branches of unidentified plant, tropical forest of Bawangling, Hainan Province, China. 12 Dec 2009, J. Ma, HSAUP H5137 (duplicate HMAS 146082).




FIG. 1. *Corynesporopsis uniseptata*. Conidiophores and conidia.

ANAMORPHIC FUNGI. Colonies effuse, blackish brown to black, hairy. Mycelium partly superficial, partly immersed in the substratum, composed of branched, septate, pale brown to brown, smooth-walled hyphae, 2–4.5 μm wide. Conidiophores differentiated, arising single or in groups on the hyphae, erect, straight or flexuous, unbranched, brown, smooth, septate, up to 160 μm long, 3.5–5 μm wide. Conidiogenous cells monotretic, integrated, terminal, determinate, cylindrical, brown, smooth, 16–35 μm long, 4.5–6 μm wide. Conidia acrogenous, dry, in acropetal chains of up to 10, ellipsoid to cylindrical, 1-euseptate, constricted at the septum, smooth, brown, often with darker pigmentation at the septum, 14–21 μm long, 6–8 μm wide in the widest part.

NOTES: *Corynesporopsis uniseptata* is reported for the first time from China. Compared with the morphology of the type specimen described by Kirk (1981b), the conidia of our collection are longer (14–21 μm vs. 12–16 μm) and the conidiophores are also longer (up to 160 μm vs. 60–100 μm), but we believe they are basically the same species. *Corynesporopsis uniseptata* most closely resembles *C. cylindrica* B. Sutton (Sutton 1989) in conidial shape and size range but differs in having didymospores with a median constriction at the septum. Moreover, the conidia of *C. cylindrica* are guttulate while *C. uniseptata* conidia are not.

Corynesporopsis quercicola (Borowska) P.M. Kirk, Trans. Br. Mycol. Soc. 77(2):284 (1981) FIG. 2

= *Corynespora quercicola* Borowska, Acta Mycol. 11(1): 60 (1975)

SPECIMEN EXAMINED: on dead branches of unidentified plant, tropical forest of Bawangling, Hainan Province, China. 10 Dec 2009, J. Ma, HSAUP H5082 (duplicate HMAS 146083).

ANAMORPHIC FUNGI. Colonies effuse, blackish brown to black, hairy. Mycelium partly superficial, partly immersed in the substratum, composed of branched, septate, pale brown, smooth-walled hyphae, 2–4.5 μm wide. Conidiophores differentiated, arising single or in groups, erect, straight or flexuous, unbranched, brown to dark brown, smooth, septate, 45–114 μm long, 3–4 μm wide, sometimes once or twice percurrent. Conidiogenous cells monotretic, integrated, terminal, cylindrical, brown, smooth. Conidia acrogenous, dry, in short acropetal chains, broadly ellipsoid to oblong, mainly 2-euseptate, rarely 1-euseptate, sometimes slightly constricted at the septum, smooth, polar cells pale brown, middle cell brown to dark brown, 13–21 μm long, 6–7 μm wide in the widest part.

NOTES: This is the first report of this species in China. The conidia of the specimen examined are somewhat longer (13–21 μm vs. 12–18 μm) than those




FIG. 2. *Corynesporopsis quercicola*. Conidiophores and conidia.

of the type specimen described by Kirk (1981a). This species has been recorded from Russia, Poland, United Kingdom, and Cuba. *Corynesporopsis quercicola* somewhat resembles *C. biseptata* (Morgan-Jones 1988) in conidial shape and septation but has smaller ($13\text{--}21 \times 6\text{--}7 \mu\text{m}$ vs. $18\text{--}33 \times 7\text{--}9 \mu\text{m}$) versicolored conidia.

Corynesporopsis indica P.M. Kirk, Mycotaxon 17: 405 (1983)

FIG. 3

SPECIMEN EXAMINED: on dead branches of unidentified plant, tropical forest of Baomeiling, Hainan Province, China. 9 Dec 2009, J. Ma, HSAUP H5274-1 (duplicate HMAS 146084).




FIG. 3. *Corynesporopsis indica*. A. Conidiophores and conidia. B. Conidia.

ANAMORPHIC FUNGI. Colonies effuse, blackish brown to black, hairy. Mycelium mostly immersed in the substratum, composed of branched, septate, pale brown to brown, smooth-walled hyphae, 1.5–3 µm wide. Conidiophores differentiated, single or in groups, erect, straight or slightly flexuous, unbranched, brown to dark brown, smooth, septate, 67–172 µm long, 3–5.5 µm wide, sometimes swollen at the base. Conidiogenous cells integrated, terminal, monotretic, cylindrical, brown, smooth, sometimes with percurrent proliferation. Conidia acrogenous, dry, solitary or in acropetal chains of 2 or 3, ellipsoid to broadly obovoid, sometimes somewhat biconic, i with one indistinct median euseptic,

the septum usually obscured by a darkly pigmented band, smooth, dark brown to blackish brown, 16–27 µm long, 8–13.5 µm wide in the widest part.

NOTES: This species has not been previously recorded in China. The size range of conidia and conidiophores in our specimen overlaps well with that of the type specimen described by Kirk (1983), and other features of this taxon also match those of the original species. *Corynesporopsis indica* is unique within the genus in its ellipsoid to obovoid, 1-septate conidia with the septum usually obscured by a band of pigment.

Acknowledgments

The authors express gratitude to Dr W.B. Kendrick and Dr R.F. Castañeda Ruiz for serving as pre-submission reviewers and for their valuable comments and suggestions. This project was supported by the National Natural Science Foundation of China (Nos. 30499340, 30770015) and the Ministry of Science and Technology of the People's Republic of China (Nos. 2006FY120100, 2006FY110500–5).

Literature cited

- Castañeda Ruiz RF, Kendrick B. 1990. Conidial fungi from Cuba: II. Univ Waterloo Biol Ser 33: 1–61.
- Ellis MB. 1972. Dematiaceous hyphomycetes. XI. Mycological Papers 131: 1–25.
- Güssow HT. 1906. Über eine neue Krankheit an Gurken in England (*Corynespora mazei*, Güssow gen. et sp. nov.). Zeitschrift für Pflanzenkrankheiten 16: 10–13.
- Holubová-Jechová V. 1987. Studies on hyphomycetes from Cuba VI. New and rare species with tretic and phialidic conidiogenous cells. Česká Mykologie 41(2): 107–114.
- Holubová-Jechová V, Mercado SA. 1986. Studies on hyphomycetes from Cuba IV. Dematiaceous hyphomycetes from the Province Pinar del Rio. Česká Mykologie 40(3): 142–164.
- Kirk PM. 1981a. New or interesting microfungi II. Dematiaceous hyphomycetes from Esher Common, Surrey. Transactions of the British Mycological Society 77(2): 279–297. doi:10.1016/S0007-1536(81)80031-9.
- Kirk PM. 1981b. New or interesting microfungi III. A preliminary account of microfungi colonizing *Laurus nobilis* leaf litter. Transactions of the British Mycological Society 77(3): 457–473. doi:10.1016/S0007-1536(81)80093-9.
- Kirk PM. 1983. New or interesting microfungi VIII. *Corynesporopsis indica* sp. nov.. Mycotaxon 17: 405–408.
- Lu BH, Hyde KD, Ho WH, Tsui K., Taylor JE, Wong KM, Yanna, Zhou DQ. 2000. Checklist of Hong Kong Fungi. Fungal Diversity Research Series 5: 1–207.
- Matsushima T. 1993. Matsushima Mycological Memoirs No. 7. Published by the author, Kobe, Japan.
- Morgan-Jones G. 1988. Notes on hyphomycetes. LVII. *Corynespora biseptata*, reclassified in *Corynesporopsis*. Mycotaxon 31(2): 511–515.
- Munjal RL, Gill HS. 1961. *Corynesporella*: a new genus of hyphomycetes. Indian Phytopathology 14(1): 6–9.
- Sutton BC. 1989. Notes on deuteromycetes. II. Sydowia 41: 330–343.