

MYCOTAXON

DOI: 10.5248/114.459

Volume 114, pp. 459–462

October–December 2010

Two new species of *Stachybotrys* from soil

YUE-MING WU & TIAN-YU ZHANG*

tyzhang1937@yahoo.com.cn

Department of Plant Pathology, Shandong Agricultural University
Taian, 271018, China

Abstract — Two new species are described and illustrated: *Stachybotrys jiangziensis* and *S. xigazensis*, both from soil in China. The type specimens (dried cultures) and living cultures are deposited in the Herbarium of Shandong Agricultural University Plant Pathology (HSAUP). Isotypes are kept in the Herbarium of Institute of Microbiology, Academia Sinica (HMAS).

Key words — taxonomy, soil fungi, dematiaceous hyphomycetes

Introduction

Stachybotrys Corda was erected in 1837, and since then 96 epithets have been proposed in the genus (Index Fungorum 2010). This genus is characterized by distinct, mononematous conidiophores bearing an apical cluster of several swollen phialides producing unicellular phialoconidia that become aggregated in globose masses. In the course of a survey of soil dematiaceous hyphomycetes in China, several unusual species of *Stachybotrys* were collected. Two of them are described as new species, *S. jiangziensis* and *S. xigazensis*.

Taxonomy***Stachybotrys jiangziensis* Y.M. Wu & T.Y. Zhang, sp. nov.**

FIG. 1

MYCOBANK MB 518786

Coloniae in *CMA effusae, atrogriseae vel nigrae. Hyphis ramosis, septatis, laevibus, hyalinis vel subhyalinis, 1.5–3 µm crassis. Conidiophora erecta, 2–4-septata, basim versus subhyalina, supra griseo-brunnea, levia, 60–80 µm longa, ad basim 4–5 µm diam. Phialides 6–8 ad apicem conidiophori productae, pallide brunneae, leves, 8–10 × 5–7 µm. Conidia tuberculata, globosa vel subglobosa, brunnea vel atrobrunnea, 6–9 µm diam.*

HOLOTYPE: China. Tibet, Jiangzi, from a grassland soil, altitude 4050 m, 9 Sept. 2007, Y.M. Wu, HSAUPII₀₇0881, **holotype**; HMAS 196256, **isotype**.

ETYMOLOGY: in reference to the type locality.

*Corresponding author: Tian-Yu Zhang




FIG. 1. *Stachybotrys jiangziensis* (ex holotype). Conidia, conidiophores, and conidiogenous cells. Above: photomicrographs. Below: drawings. (Bars = 25 μ m).

Colonies on CMA (cornmeal agar) at 25°C for 21 days 4–6 cm diam., effuse, darkish grey to black. Mycelium mostly superficial, partly immersed. Hyphae branched, septate, smooth, hyaline to subhyaline, 1.5–3 μ m wide. Conidiophores

erect, branched, 2–4-septate, subhyaline near the base, greylish brown above, smooth, 60–80 μm long, 4–5 μm wide near base. Phialides borne in groups of 6–8 at the apices of conidiophores, pale brown, smooth, 8–10 \times 5–7 μm . Conidia globose to subglobose, tuberculate, brown to dark brown, 6–9 μm in diameter.

In conidial morphology this fungus somewhat resembles *Stachybotrys nilagirica* Subram. (Subramanian 1957) and *S. sphaerospora* Morgan-Jones & R.C. Sinclair (Morgan-Jones & Sinclair 1980). However *S. nilagirica* has larger conidia (16–20 μm diam.) and *S. sphaerospora* larger (11–12 μm diam.), ridged conidia than *S. jiangziensis*.

Stachybotrys xigazensis Y.M. Wu & T.Y. Zhang, sp. nov.

FIG. 2

MYCOBANK MB 518787

Coloniae in CMA effusae, atrogriseae vel nigrae. Hyphis ramosis, septatis, laevibus, hyalinis vel subhyalinis, 2–3 μm crassis. Conidiophora erecta, 1–4-septata, basim versus subhyalina, supra griseo-brunnea, verrucosa, interdum granulis magnis tecta, 60–100 μm longa, ad basim 4–6 μm diam. Phialides 6–8 ad apicem conidiophori productae, pallide brunneae, leves, 7–10 \times 5–8 μm . Conidia ovoidea, ellipsoidea vel oblonga, tuberculata, brunnea vel atrobrunnea, 9–12.5 \times 7.5–10 μm .



FIG. 2. *Stachybotrys xigazensis* (ex holotype). Conidia, conidiophores, and conidiogenous cells. Left: photomicrographs. Right: drawings. (Bars = 25 μm).

HOLOTYPE: China, Tibet, Xigazen, from a grassland soil, altitude 4150 m, 19 Sept. 2007, Y.M. Wu, HSAUPII₀₇ 1450, **holotype**; HMAS 196257, **isotype**.

ETYMOLOGY: in reference to the type locality.

Colonies on CMA at 25°C for 21 days 5–8 cm diam., effuse, darkish grey to black. Mycelium mostly superficial, partly immersed. Hyphae branched, septate, smooth, hyaline to subhyaline, 2–3 µm wide. Conidiophores erect, sympodially branched, 1–4-septate, subhyaline near the base, greyish brown above, verrucose, sometime covered with large granules, 60–100 µm long, 4–6 µm wide near base. Phialides borne in groups of 6–8 at the apices of conidiophores, pale brown, smooth, 7–10 × 5–8 µm. Conidia ovoid, ellipsoid or oblong, tuberculate, brown to dark brown, 9–12.5 × 7.5–10 µm.

This fungus somewhat resembles *Stachybotrys chartarum* (Ehrenb.) S. Hughes (Hughes 1958) and *S. microspora* (B.L. Mathur & Sankhla) S.C. Jong & E.E. Davis (Jong & Davis 1976) in conidial colour and size, but *S. xigazensis* has more obviously tuberculate conidia. In addition, the conidia of *S. xigazensis* are larger than those of *S. microspora* (6–8 × 4–5 µm) and wider than those of *S. chartarum* (7–12 × 4–6 µm).

Acknowledgments

The authors are grateful for pre-submission comments and suggestions provided by Dr. Eric McKenzie, Prof. Y.L. Guo, and Dr. Shaun Pennycook. This project was supported by the National Science Foundation of China (no. 30670014 & 30499340).

Literature cited

- Hughes SJ. 1958. Revisioes Hyphomycetum aliquot cum appendice de nominibus rejiciendis. Canadian Journal of Botany 36: 727–836. doi: 10.1139/b58-067.
- Index Fungorum. 2010. <http://www.indexfungorum.org/Names/Names.asp>; accessed 28 August 2010.
- Jong SC, Davis EE. 1976. Contribution to the knowledge of *Stachybotrys* and *Memnoniella* in culture. Mycotaxon 3: 409–485.
- Morgan-Jones G, Sinclair RC. 1980. Notes on Hyphomycetes. XXXIII. *Stachybotrys sphaerospora* sp. nov. from South Africa. Mycotaxon 10: 372–374.
- Subramanian CV. 1957. Hyphomycetes–IV. Proceedings of the Indian Academy of Sciences B. 46: 330–331.