

## MYCOTAXON

Volume 115, pp. 1–4

January–March 2011

DOI: 10.5248/115.1

**A new species of *Ophiocordyceps* (Clavicipitales, Ascomycota) from southwestern China**JI-YUE CHEN<sup>1\*</sup>, YONG-QIANG CAO<sup>1</sup>, DA-RONG YANG<sup>1\*</sup> & MING-HUA LI<sup>2</sup><sup>1</sup>Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Kunming 650223, Yunnan, China<sup>2</sup>Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, Yunnan, China\*CORRESPONDENCE TO: [chenjy@xtbg.ac.cn](mailto:chenjy@xtbg.ac.cn), [yangdr@xtbg.ac.cn](mailto:yangdr@xtbg.ac.cn)

ABSTRACT — A new species of caterpillar fungus, *Ophiocordyceps laojunshanensis*, is described based on specimens collected from southwestern China. This species is characterized by slender stromata, sparse perithecia, and the unique habitat of growing in mosses. Its habitat, gross morphology, and microscopic features are illustrated and relationships to similar species discussed.

KEY WORDS — *Cordyceps*, *Ophiocordyceps sinensis*, *Ophiocordycipitaceae*, taxonomy

**Introduction**

The genus *Cordyceps* includes over 400 species, of which 120 species have been reported in China (Liang 2007). Although the genus was formerly classified in the *Clavicipitaceae*, based on recent results of the multi-gene phylogeny of *Cordyceps* sensu lato, the taxonomy of both *Cordyceps* and the *Clavicipitaceae* has been revised.

A new family *Ophiocordycipitaceae* has been proposed based on the genus *Ophiocordyceps* Petch, which has been emended by Sung et al. (2007). *Ophiocordyceps* is now characterized by stromata that are darkly (or rarely brightly coloured) pigmented, tough, fibrous, pliant to wiry and rarely fleshy, often with aperithecial apices or lateral pads, perithecia that are superficial to completely immersed and arranged ordinaly or obliquely (Sung et al. 2007).

In the past three years, the authors undertook numerous mycological explorations to mountainous Northern Yunnan, where they collected many specimens of caterpillar fungi that grow among mosses and have morphological features that do not correspond to any previously described taxa. One new species, *Ophiocordyceps laojunshanensis*, is described and illustrated in this paper.

## Materials & methods

Field notes were taken for gross morphology of fresh ascomata and habitat. Detailed macroscopic features were examined under a stereomicroscope (Carl Zeiss Discovery 12). Microscopic examination of ascomata and measurements of microscopic structures were made from freehand sections under a microscope (Olympus 520). The sections were mounted in 5% aqueous KOH and then Cotton–blue lactophenol. The holotype of *Ophiocordyceps laojunshanensis* was deposited in the Cryptogamic Herbarium, Kunming Institute of Botany, Chinese Academy of Sciences (KUN–HKAS 57066). Isolates from fresh ascomata of *O. laojunshanensis* were maintained on potato dextrose agar with 1% peptone (PPDA) using techniques described by Liu et al (1989), all isolates were kept in Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences (No. lao1–lao20).

## Taxonomy

*Ophiocordyceps laojunshanensis* J.Y. Chen, Y.Q. Cao & D.R. Yang, **sp. nov.** FIG. 1

MYCOBANK MB 517206

*Stromatibus clavatis, singularibus, 47.0–93.0 mm longis, 1.0–3.9 mm crassis. Capitulis cylindricis, 8.5–17.0 mm longis, prophyro–brunneis vel atro–brunneis, apic acerosis sterilibus 3.5–8.0 mm longis. Perithecia globosa, 200–300 × 200–350 µm; Stipitibus, 25.0–72.0 mm longis; Asci clavuli, cylindraci, 165–275 × 11.5–14.5 µm. Ascosporis hyalinis, filiformis, multiseptatis, 130–250 × 5.0–6.0 µm.*

TYPE: China: Yunnan Province, Lijiang Naxi Autonomous Prefecture, Laojunshan mountain, 26°37'33.76N 99°42'58.07E, alt. 3974 m; under *Rhododendron mariae* Hance; associated with *Oncophorus wahlenbergii* Brid. and *Pleurozium schreberi* (Brid.) Mitt., 8 May 2009, Ji–yue Chen 09032. (HOLOTYPE: KUN–HKAS 57066).

ETYMOLOGY: The epithet refers to the type locality, Laojunshan Mountain

Stromata clavate, slender, simple, rarely 2 or 3 from host head, rarely 1 from host head and another from the terminal region, 47.0–93.0 mm long, 1.0–3.9 mm in diam. Ascogenous portion almost 1/5 to 1/3 of the stromata length, cylindrical, 8.5–17.0 mm long, purplish to dark brown. Apex sterile acuminate, 3.5–8.0 mm long. Perithecia, globoid, 200–300 × 200–350 µm. arranged loosely in irregular lateral cushions. Stipe 25.0–72.0 mm long. Asci clavate, 165.0–275.0 × 11.5–14.5 µm. Ascospores hyaline, filiform, septate, 130.0–250.0 × 5.0–6.0 µm (FIG. 1a–f).

Colonies (FIG. 1g–l) grew very slowly, only up to 6–10 mm in diameter after 2 months on PPDA at 16°C. They were white at first, then became brown–yellow or black brown, and sparse white mycelia emerged 3 month later, with conidiophore and conidia, reverse blackish brown. Conidiophore, hyaline, branching or non–branching; conidiogenous cell phialidic, hyaline, with verrucose, acerate, 15–39(–50) µm long; Conidia, hyaline, long elliptic, 6.0–13.5 × 3.0–4.0 µm, one or two encased in a mucus drop.

HOST: Larvae of *Thitarodes yunnanensis* Nielsen et al. (*Hepialus yunnanensis* Yang et al.)




FIG 1. *Ophiocordyceps laojunshanensis*—a. habitat; b. fresh fruiting body; c. dry fruiting bodies; d. ascogenous portion, bar = 2 mm; e. ascus with ascospore, bar = 20  $\mu\text{m}$ ; f. ascus tip, bar = 20  $\mu\text{m}$ ; g–h. culture (bar = 5 mm): g. surface, h. reverse; i. colonies, bar = 20 mm; j. conidiophore and conidia, bar = 20  $\mu\text{m}$ ; k. conidiophore (detail), bar = 25  $\mu\text{m}$ ; l. conidia (detail), bar = 10  $\mu\text{m}$ .

ADDITIONAL SPECIMENS EXAMINED: CHINA: YUNNAN PROVINCE, Lijiang Naxi Autonomous Prefecture, Laojunshan mountain, 26°37'03.98N 99°42'18.02E to 26°37'82.71N 99°42'97.86E, alt. 3874–4075 m. Under *Abies delavayi* Franch and *Rhododendron mariae* Hance. Associated with *Oncophorus wahlenbergii* Brid., *Dicranum muelhenbeckii* Bruch & Schimp. or *Pleurozium schreberi* (Brid.) Mitt., 8 May 2009, Ji–yue Chen 09033–09035.

DISCUSSION: *Ophiocordyceps laojunshanensis* is characterized by slender stromata, sparse perithecia, and its unique habitat of growing among mosses. It strongly resembles the most precious caterpillar fungus, *O. sinensis* (Berk.) G.H. Sung et al., and is often sold together with that fungus. The two species could be living in the same host larvae, except that *O. laojunshanensis* is found most often growing among mosses, while *O. sinensis* appears to grow more frequently in grasslands. Morphologically, *O. laojunshanensis* differs from *O. sinensis* by having a slimmer stroma with a shorter ascogenous portion and relatively sparse irregular perithecia compared to the close ordered perithecia found in *O. sinensis*.

### **Acknowledgments**

The authors would like to thank Prof. Mu Zang (Kunming Institute of Botany, Chinese Academy of Sciences) for his valuable suggestions and discussion. They also give thanks to Dr. Yun Wang (Plant & Food Research, New Zealand) and Mr. Daniel Winkler (MushRoaming, USA) for serving as pre-submission reviewers, Dr. Nicholas Cummings (Plant & Food Research, New Zealand) for reviewing English, Prof. X.J. Li (Kunming Institute of Botany, Chinese Academy of Sciences) for identifying the moss species. This study was supported by the Knowledge Innovation Program of the Chinese Academy of Sciences Project (No. KSCX2-YW-G-024), the Natural Science Foundation of Yunnan (No. 2007C105M), and West doctor Project of West Light Foundation of The Chinese Academy of Sciences.

### **Literature cited**

- Liang ZQ. 2007. *Cordyceps*, Flora Fungorum Sinicorum. Science Press, Beijing. 32: 1–190.
- Liu XJ, Guo YL, Yu YX, Zeng W. 1989. Isolation and identification of the anamorphic stage of *Cordyceps sinensis* (Berk.) Sacc. *Acata Mycologica Sinica* 8: 35–40.
- Sung GH, Hywel-Jones NL, Sung JM, Luangsa-aard J, Shrestha B, Spatafora JW. 2007. Phylogenetic classification of *Cordyceps* and the clavicipitaceous fungi. *Studies in Mycology* 57: 5–59. doi:10.3114/sim.2007.57.01