

---

# MYCOTAXON

<http://dx.doi.org/10.5248/117.359>

Volume 117, pp. 359–363

July–September 2011

---

## A new species of *Lepiota* (Agaricaceae) from southwestern China

JUN F. LIANG<sup>1\*</sup> & ZHU L. YANG<sup>2</sup>

<sup>1</sup>Research Institute of Tropical Forestry, Chinese Academy of Forestry,  
Guangzhou 510520, P. R. China

<sup>2</sup>Key Laboratory of Biodiversity and Biogeography, Kunming Institute of Botany,  
Chinese Academy of Sciences, Kunming 650204, P. R. China

\* CORRESPONDENCE TO: [jfliang2000@163.com](mailto:jfliang2000@163.com)

**ABSTRACT** — A new species, *Lepiota pallidiochracea*, is described. It is characterized by the ochraceous-red squamules on pileus, large ellipsoid to oblong spores, polymorphic cheilocystidia, and a trichodermium of elongate pileus covering, which is apically attenuate.

**KEW WORDS** — Agaricales, lepiotaceous fungi, taxonomy

### Introduction

The eastern Himalaya is one of the twenty-five hotspots for biodiversity (Myers et al 2000), and macrofungi are very rich in the area (Yang 2005). Many new agarics and aphylloporoid fungi have recently been described from there (Yang et al. 2004; Dai et al. 2007; Yuan & Dai 2008; Li et al. 2009; Dai 2010, 2011; Li et al. 2011). This paper is another contribution on the lepiotaceous fungi occurring in the region.

The genus *Lepiota* (Pers.) Gray contains more than 400 described species (Kirk et al 2008), and it is reasonable to expect many representatives of this cosmopolitan genus in China. However, rather few taxa have been originally described from China (Chiu 1948, Bi et al 1986, Yang 1994, Tolgor & Li 2004, Wang & Yang 2005a,b). In our research on *Lepiota* in China, in addition to several recently published new species (Liang et al 2009, 2010, 2011, Liang & Yang 2011), several interesting and undescribed taxa still await documentation. The present paper describes one new *Lepiota* species from eastern Himalaya.

### Materials & methods

Macro-morphological features were recorded in the field. Material was dried using an electric drier and deposited in the Herbarium of Cryptogams, Kunming Institute of Botany, Chinese Academy of Sciences (HKAS) and Research Institute of Tropical

Forestry, Chinese Academy of Forestry (RITF). Terminology for descriptive terms follows Vellinga & Noordeloos (2001). Color designations are from Kornerup and Wanscher (1981).

For microscopic observations, sections of fruitbodies were made by hand and mounted in 5% KOH, Congo red, and Melzer's reagent. Basidiospores were mounted in cresyl blue to test for a metachromatic reaction (Singer 1986). Size ranges were determined for basidia, basidiospores, cheilocystidia, and elements of the pileipellis, based on ocular micrometer measurements of at least 20 elements of each character. The abbreviation [n/m/p] indicates that measurements were made on n basidiospores in m basidiomata from p collections. Dimensions of basidiospores are given using a notation of the form (a-)b-c(-d). The range b-c contains a minimum of 90% of the measured values. Extreme values are given in parentheses. The following abbreviations are used: Q refers to the length/breadth ratio of basidiospores; Q refers to the average Q of all basidiospores  $\pm$  sample standard deviation.

## Taxonomy

*Lepiota pallidiochracea* J.F. Liang & Zhu L. Yang, sp. nov.

FIG. 1-4

MYCOBANK 561241

*Pileus sordide albus, squamulis densibus, rubro-ochraceis, rubro-brunneis vel brunneis. Stipes subcylindricus, squamulis brunneis. Basidiosporae 9.0-12.0  $\times$  6.0-9.0  $\mu$ m, ellipsoideae vel lato fusiformes. Cheilocystidia diversa, clavata, fusiformia vel hamata. Pleurocystidia nulla. Squamulae pilei ex trichodermiis attenuatis terminalibus compositae. Fibulae praesentes.*

TYPE: China, Tibet, Changdu Co., near Qiongka Bridge, 27 July 2004, Zhu L. Yang 4194 (HKAS 45579 holotype).

ETYMOLOGY: *pallidiochracea* (Lat.), referring to pale ochraceous squamules on the pileus.

Basidiomata (FIG. 1) small-sized. Pileus 1.5-4 cm diam., campanulate with small umbo when young, plano-convex with obtuse umbo at maturity, cream to whitish, at centre brownish red (10D6-8) to dark brown (9E4-8), around centre densely covered with concentric squamules; squamules more or less uplifted, minute, ochraceous-red (8C7-8), brown (7D8) to reddish brown (9C7-8). Lamellae free, subventricose, moderately crowded with lamellulae, dirty white to cream. Stipe 2-4  $\times$  0.2-0.3 cm, subcylindrical, hollow; surface glabrous and white at the apex, light ochraceous (7C6-7) with scattered ochraceous-red (8C7-8), brown (7D8) to reddish brown (9C7-8) squamules at the low part. Annulus whitish, membranous, evanescent. Odor none. Taste not recorded.

Basidiospores (FIG. 2) [63/3/2] 9-12(-14)  $\times$  6-8(-9)  $\mu$ m [Q = 1.29-1.85, Q = 1.56  $\pm$  0.18], ellipsoid to oblong, sometimes ovoid with small hilar appendage in side view, with neither suprahilar depression nor germ pore, ellipsoid in frontal view; hyaline, smooth, slightly thick-walled, dextrinoid, weakly congophilic, not metachromatic in Cresyl Blue. Basidia 25-37  $\times$  9-14  $\mu$ m, clavate, 4-spored, rarely 2-spored. Lamella edge sterile. Cheilocystidia




FIG. 1–4 *Lepiota pallidiochracea* (All from holotype, HKAS 45579).  
1. Basidiomata. 2. Basidiospores. 3. Cheilocystidia. 4. Pileus covering.

(FIG. 3) 15–40 × 5–10 μm, variable in shape, subcylindrical to narrowly clavate, or fusiform, lageniform to utriform; walls smooth, thin, hyaline in KOH, congophilic. Pleurocystidia absent. Pileus covering (FIG. 4) a trichodermium of elongate, apically attenuate terminal elements 60–256 × 6.5–11 μm, often with some short narrowly clavate elements interspersed, with pale yellowish brown parietal and intracellular pigment in upper part; fine incrustation sometimes in basal part of elements. Clamp connections common.

ADDITIONAL SPECIMEN EXAMINED: CHINA: YUNNAN PROV., SHANGRI-LA COUNTY, Daxiagu, alt. 3060 m, 20 July 2008, Jun F. Liang 799 (RITF 660).

HABITAT & DISTRIBUTION: in small groups, gregarious, saprotrophic and terrestrial on slope, in summer. Known only from southwestern China.

DISCUSSION: *Lepiota pallidiochracea* is characterized by its ochraceous-red squamules on the pileus, which is covered with a trichodermium of long, elongate, apically attenuate and short narrowly clavate elements, large ellipsoid to oblong spores, and polymorphic cheilocystidia.

Based on the trichodermium type of the pileus squamules and the ellipsoid basidiospores, *L. pallidiochracea* may tentatively be placed in *L.* sect. *Ovisporae*

(J.E. Lange) Kühner (Singer 1986, Vellinga 2001), although species in the section usually have spores smaller than 10  $\mu\text{m}$  (Singer 1986). Recent molecular phylogenetic studies indicate that section *Ovisporae* is not monophyletic, so that re-evaluation of this section is needed (Vellinga 2003, Liang et al 2011). Phylogenetically, the new species may be referred to section *Lepiota* or clade 1 of *Lepiota* s.l. (Vellinga 2003) until a new taxonomic system is proposed.

Several *Lepiota* species have similarly shaped spores. The polymorphic cheilocystidia coupled with the larger spores help distinguish *L. pallidiochracea* from *L. brunneolilacea* Bon & Boiffard (also a member of *L. sect. Ovisporae*), *L. helveola* Bres. and *L. ochraceoaurantiaca* Dennis (Bon 1996, Candusso & Lanzoni 1990, Dennis 1952). The larger spores are reminiscent of *L. oreadiformis* Velen. (in sect. *Lepiota*), which, however, has much larger, fusiform to amygdaliform spores and narrowly clavate or utriform cheilocystidia (Vellinga 2001).

#### Acknowledgments

We are grateful to Dr. Xiang-Hua Wang and Dr. Yan-Chun Li for offering advice, suggestions, and analytical assistance. We are greatly indebted to Dr. Yu-Cheng Dai and Dr. Ping Zhang for their critical reviewing the manuscript. This study is supported by the National Natural Science Foundation of China (No. 31070014), the Joint Fund of the National Natural Science Foundation of China and Yunnan Province (No. U0836604), the National Basic Research Program of China (No. 2009CB522300), the Foundation of RITF (RITFKYYW 2010-10), and the Commonweal Industry-specific Foundation of State Forestry Administration (No. 201104057).

#### Literature cited

- Bi ZS, Li TH, Zheng GY. 1986. New and rare species of *Agaricales*. Acta Mycol Sinica Suppl. 1: 288–296.
- Bon M. 1996. Die Großpilzflora von Europa 3 *Lepiotaceae*. IHW-Verlag: Eching (Germany). 141 p.
- Candusso M, Lanzoni G. 1990. Fungi Europaei 4. *Lepiota* s.l. Giovanna Biella: Saronno. 743 p.
- Chiu WF. 1948. The *Amanitaceae* of Yunnan. Sci Rept Natl Tsing Hua Univ Ser B, Biol, Psychol Sci 3(3): 165–178.
- Dai YC, 2010. *Hymenochaetaceae (Basidiomycota)* in China. Fungal Diversity 45: 131–343.
- Dai YC, 2011. A revised checklist of corticioid and hydroid fungi in China for 2010. Mycoscience 52: 69–79.
- Dai YC, Yu CJ, Wang HC, 2007. Polypores from eastern Xizang (Tibet), western China. Ann Bot Fenn 44: 135–145.
- Dennis RW. 1952. *Lepiota* and allied genera in Trinidad, British West Indies. Kew Bull 7: 459–499.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. Ainsworth & Bisby's dictionary of the fungi. 10th Edition. CABI Publishing, Wallingford.
- Kornerup A, Wanscher JH. 1981. Taschenlexikon der Farben. 3. Aufl. Muster-Schmidt Verlag, Göttingen.
- Li YC, Yang ZL, Tolgor B, 2009. Phylogenetic and biogeographic relationships of *Chroogomphus* species as inferred from molecular and morphological data. Fungal Divers 38: 85–104.

- Li YC, Feng B, Yang ZL, 2011. *Zangia*, a new genus of *Boletaceae* supported by molecular and morphological evidence. *Fungal Divers* 49: 125–143.  
<http://dx.doi.org/10.1007/s13225-011-0096-y>.
- Liang JF, Yang ZL. 2011. Two taxa close to *Lepiota cristata* from China. *Mycotaxon* 116: 387–394.  
<http://dx.doi.org/10.5248/116.387>.
- Liang JF, Xu J, Yang ZL. 2009. Divergence, dispersal and recombination in *Lepiota cristata* from China. *Fungal Divers* 38: 105–124.
- Liang JF, Yang ZL, Xu J, Ge ZW. 2010. Two new unusual *Leucoagaricus* species (*Agaricaceae*) from tropical China with blue-green staining reactions. *Mycologia* 102: 1141–1152.  
<http://dx.doi.org/10.3852/09-021>.
- Liang JF, Yang ZL, Xu DP. 2011. A new species of *Lepiota* from China. *Mycologia* 103: 820–830.  
<http://dx.doi.org/10.3852/10-216>.
- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.
- Singer R. 1986. *The Agaricales in modern taxonomy*. 4th ed. Koeltz Scientific Books, Koenigstein.
- Tolgor B, Li Y. 2004. *Lepiota squamulosa*, a new species from China. *J Fung Res* 2 (3): 49–50.
- Vellinga EC. 2001. *Lepiota* (Pers.: Fr.) S.F. Gray. 109–151, in: ME Noordeloos et al. (eds). *Flora Agaricina Neerlandica*, Vol. 5. A.A. Balkema, Rotterdam.
- Vellinga EC. 2003. Phylogeny of *Lepiota* (*Agaricaceae*) – evidence from nrITS and nrLSU sequences. *Mycol Prog* 2: 305–322.
- Vellinga EC, Noordeloos ME. 2001. Glossary. 6–11, in: ME Noordeloos et al. (eds). *Flora Agaricina Neerlandica*, Vol. 5. A.A. Balkema, Rotterdam.
- Wang HC, Yang ZL. 2005a. Notes on *Lepiota shixingensis* and an allied new species (*Basidiomycetes*). *Nova Hedwigia* 81(3–4): 463–469.
- Wang HC, Yang ZL. 2005b. A new species of *Lepiota* (*Agaricaceae*, *Basidiomycetes*) from China. *Mycotaxon* 91: 51–54.
- Yang ZL. 1994. *Clarkeinda*, *Lepiota*, *Leucoagaricus*, *Leucocoprinus* and *Macrolepiota*. 122–131, in: Ying JZ, Zang M (eds). *Economic macrofungi of southwestern China*. Science Press: Beijing.
- Yang ZL. 2005. Diversity and biogeography of higher fungi in China. 35–62, in: Xu J (ed). *Evolutionary genetics of fungi*. Norfolk (UK): Horizon Bioscience.
- Yang ZL, Weiß M, Oberwinkler F. 2004. New species of *Amanita* from the eastern Himalaya and adjacent regions. *Mycologia* 96: 636–646.
- Yuan HS, Dai YC, 2008. Polypores from northern and central Yunnan Province, southwestern China. *Sydowia* 60: 147–159.