

MYCOTAXON

<http://dx.doi.org/10.5248/118.283>

Volume 118, pp. 283–288

October–December 2011

***Septobasidium saurauiae* sp. nov. (Septobasidiaceae)
and *S. pseudopedicellatum* new to China**SUZHEN CHEN^{1,2} & LIN GUO^{1*}¹Key Laboratory of Systematic Mycology and Lichenology, Institute of Microbiology,
Chinese Academy of Sciences, Beijing 100101, China²Ocean University of China, Qingdao 266003, ChinaCORRESPONDENCE TO *: Chensz2009@gmail.com & *guol@im.ac.cn

ABSTRACT—A new species, *Septobasidium saurauiae* on *Saurauia tristyla* associated with *Chionaspis* sp., and a new Chinese record, *Septobasidium pseudopedicellatum* on *Schefflera octophylla* and *Psychotria serpens*, are described. They were collected from Hainan Province.

KEY WORDS—Pucciniomycetes, Septobasidiales, taxonomy

Hainan Island is located in the South China Sea. It is a tropical area with a rich fungal diversity. To date, six new species and one new Chinese record of *Septobasidium* have been discovered from this Province (Chen & Guo 2011b, Lu & Guo 2009a, 2010b,c). An additional new species and a new Chinese record of *Septobasidium* are reported as follows:

***Septobasidium saurauiae* S.Z. Chen & L. Guo, sp. nov.**

FIGS. 1–7

MYCOBANK MB 561871

Basidiomata resupinata, discontinue crescentia, 6–7 cm longa, 3–4 cm lata, cinnamomeo-brunnea vel griseo-brunnea, margine determinata, superficie laevia, saepe protuberationibus rotundis praedita, in sectione 900–1330 µm crassa. Subiculum brunneum, 30–50 µm crassum, ab subiculo stratum hypharum vel curtam columnam formans. Columnae brunneae, 20–50 µm altae, 20–70 µm latae. Strata hyphararum 780–970 µm alta. Ab strato hymenii hyphae saepe repullulantes tum stratum hypharum et hymenium secundum formantes. Stratum hypharum secundum 200–240 µm altum. Hymenium hyalinum, 40–50 µm crassum. Basidia cylindrica, recta vel leviter curvata, 4-cellularia, 42–50 × 6–8 µm, hyalina. Sine probasidio. Basidiosporae 10–25 × 4–6 µm, hyalinae. Haustoria hyalina, ex hyphis irregulariter spiralibus constantia.

TYPE: China, Hainan Province, Limu Mountain, alt. 529 m, on *Saurauia tristyla* DC. (*Actinidiaceae*), associated with *Chionaspis* sp. (*Diaspididae*), 23.XI.2010, Y.F. Zhu & F. He 520 (holotype, HMAS 263145).










FIG. 1. Basidia and basidiospores of *Septobasidium saurauiae* (HMAS 263145, holotype).

Basidiomata on branches, resupinate, growing discontinuously, 6–7 cm long, 3–4 cm wide, cinnamon-brown or grey-brown; margin determinate, surface smooth, frequently with round protuberance. In section 900–1330 µm thick. Subiculum brown, 30–50 µm thick. Forming hyphal layer or short pillars. Pillars brown, 20–50 µm high, 20–70 µm wide. Hyphal layer 780–970 µm high. From hymenial layer the fungal hyphae often renew growth to form the second hyphal layer and hymenium. Second hyphal layer 200–240 µm high. Hymenium hyaline, 40–50 µm thick. Basidia arising directly from the hyphae without a probasidial cell, at first obovoid, later cylindrical, straight or slightly curved, 4-celled, 42–50 × 6–8 µm, hyaline. Basidiospores 10–25 × 4–6 µm, hyaline. Haustoria hyaline, consisting of irregularly coiled hyphae.

REMARKS: Morphologically, *Septobasidium saurauiae* is similar to *S. meizhouense* C.X. Lu et al., which differs in numerous cracks in the basidioma surface and a thinner section (400–650 µm high) and hyphal layer (220–440 µm high) (Lu et al. 2010).

FIGS. 2–7 (right). *Septobasidium saurauiae* (HMAS 263145, holotype). 2. Basidiomata on branch. 3–4. Sections of basidiomata. 5–6. Basidia (arrows). 7. Haustoria.






FIG. 8. Basidia and probasidia of *Septobasidium pseudopedicellatum* (HMAS 242745).






Septobasidium pseudopedicellatum Burt, Ann. Mo. bot. Gdn 3: 327, 1916.

FIGS. 8–14

Basidiomata on trunks, branches and leaves, resupinate, 0.3–19 cm long, 0.2–9 cm wide, chestnut brown, brown or smoke grey; margin white, determinate, surface smooth. The pillars near the margin are visible with naked eyes. In section 850–1700 μm thick. Subiculum white or brown, 30–50 μm thick. Pillars brown, 500–700 μm high, 50–170 μm wide. The pillars branch out to form hyphal layer, 400–650 μm high. From hymenial layer the fungal hyphae often renew growth to form the second hyphal layer and hymenium, up to 3-strata. Hymenium brown, 40–60 μm thick, with closely packed paraphyses that are parallel, upright and curved at top. Probasidia hyaline or brownish, obovoid or ellipsoidal, 16–20 \times 10–16 μm . Basidia cylindrical, straight or slightly curved, 4-celled, 52–62 \times 5–7.5 μm , hyaline, with persisting probasidial cell. Sterigmata conical, 2.5–8 \times 2.5–3.5 μm . Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.

SPECIMENS EXAMINED: CHINA, HAINAN PROVINCE, Bawangling Natural Reserve, Nanchahe, alt. 600 m, on *Schefflera octophylla* (Lour.) Harms (*Araliaceae*), 14.IV.2011, L.Guo 11611, HMAS 242745; on *Psychotria serpens* L. (*Rubiaceae*), 14.IV.2011, L.Guo 11612, HMAS 251216.

FIGS. 9–14 (right). *Septobasidium pseudopedicellatum* (HMAS 242745). 9. Basidiomata on trunk. 10–11. Sections of basidiomata. 12. Probasidia (arrows). 13. Basidium (arrow). 14. Haustoria.



Including the two species reported in this paper, 35 *Septobasidium* species have now been reported in China (Sawada 1933, Couch 1938, Teng 1963, Tai 1979, Kirschner & Chen 2007, Lu & Guo 2009a,b,c, 2010a,b,c, 2011, Lu et al. 2010, Chen & Guo 2011a,b).

Acknowledgements

The authors would like to express their deep thanks to Drs. Eric H.C. McKenzie (Auckland, New Zealand) and Shuanghui He (Beijing Forestry University) for serving as pre-submission reviewers, to Dr. Shaun Pennycook (Auckland, New Zealand) for nomenclatural review, to Prof. Jian-Yun Zhuang (Institute of Microbiology, Chinese Academy of Sciences) for Latin corrections, to Prof. Zhenyu Li and Mr. Ziyu Cao (Institute of Botany, Chinese Academy of Sciences) and Mr. Qing Chen (Bawangling Natural Reserve, Hainan Province) for identifying the host plants, to Prof. Sanan Wu (Beijing Forestry University) for identifying the scale insect, and to Mrs. Xiangfei Zhu for inking in line drawings. This study was supported by the Foundation of Ministry of Science and Technology of the People's Republic of China (No. 2006FY110500-5).

Literature cited

- Chen SZ, Guo L. 2011a. *Septobasidium sichuanense* sp. nov. (*Septobasidiaceae*) from China. Mycotaxon 115: 481–484. <http://dx.doi.org/10.5248/115.481>
- Chen SZ, Guo L. 2011b. *Septobasidium atalantiae* sp. nov. (*Septobasidiaceae*) and *S. henningsii* new to China. Mycotaxon 117: 291–296. <http://dx.doi.org/10.5248/117.291>
- Couch JN. 1938. The genus *Septobasidium*. Univ. of North Carolina Press, Chapel Hill. 480 p.
- Kirschner R, Chen CJ. 2007. New reports of two hypophyllous *Septobasidium* species from Taiwan. Fung. Sci. 22(1,2): 39–46.
- Lu CX, Guo L. 2009a. *Septobasidium maesae* sp. nov. (*Septobasidiaceae*) from China. Mycotaxon 109: 103–106. <http://dx.doi.org/10.5248/109.103>
- Lu CX, Guo L. 2009b. Two new species of *Septobasidium* (*Septobasidiaceae*) from China. Mycotaxon 109: 477–482. <http://dx.doi.org/10.5248/109.477>
- Lu CX, Guo L. 2009c. *Septobasidium annulatum* sp. nov. (*Septobasidiaceae*) and *Septobasidium kameii* new to China. Mycotaxon 110: 239–245. <http://dx.doi.org/10.5248/110.239>
- Lu CX, Guo L. 2010a. Three new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. Mycotaxon 112: 143–151. <http://dx.doi.org/10.5248/112.143>
- Lu CX, Guo L. 2010b. Two new species of *Septobasidium* (*Septobasidiaceae*) and *S. pallidum* new to China. Mycotaxon 113: 87–93. <http://dx.doi.org/10.5248/113.87>
- Lu CX, Guo L. 2010c. Two new species of *Septobasidium* (*Septobasidiaceae*) from Hainan province in China. Mycotaxon 114: 217–223. <http://dx.doi.org/10.5248/114.217>
- Lu CX, Guo L. 2011. Two new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. Mycotaxon 116: 395–400. <http://dx.doi.org/10.5248/116.395>
- Lu CX, Guo L, Wei JG, Li JB. 2010. Two new species of *Septobasidium* (*Septobasidiaceae*) from southern China. Mycotaxon 111: 269–274. <http://dx.doi.org/10.5248/111.269>
- Sawada K. 1933. Descriptive catalogue of the Formosan fungi. Part VI. Rep. Dept. Agric. Govt. Res. Inst. Formosa 61: 1–99.
- Tai FL. 1979. Sylloge Fungorum Sinicorum. Science Press, Beijing. 1527 p.
- Teng SC. 1963. Fungi of China. Science Press, Beijing. 808 p.