

MYCOTAXON

<http://dx.doi.org/10.5248/120.1>

Volume 120, pp. 1–9

April–June 2012

***Glomus truffemii* (Glomeromycetes), a new sporocarpic species from Brazilian sand dunes**BRUNO TOMIO GOTO¹, JOMAR GOMES JARDIM¹, GLADSTONE ALVES DA SILVA²,
EDUARDO FURRAZOLA³, YAMIR TORRES-ARIAS³ & FRITZ OEHL⁴¹*Departamento de Botânica, Ecologia e Zoologia, CB, Universidade Federal do Rio Grande do Norte, Campus Universitário, 59072-970, Natal, RN, Brazil*²*Departamento de Micologia, CCB, Universidade Federal de Pernambuco,**Av. Prof. Nelson Chaves, S/N, CEP 50670-420, Cidade Universitária, Recife, PE, Brazil*³*Instituto de Ecologia e Sistemática, IES-CITMA, A.P. 11900, La Habana, Cuba*⁴*Federal Research Institute Agroscope Reckenholz-Tänikon ART, Organic Farming Systems, Reckenholzstrasse 191, CH-8046 Zürich, Switzerland**CORRESPONDENCE TO: brunogoto@hotmail.com

ABSTRACT — *Glomus truffemii*, which forms large aggregates ($\leq 850 \times 1200 \mu\text{m}$) in the rhizosphere of a herbaceous plant community in NE Brazilian sand dunes, is described as new. Its subglobose glomerospores measuring $72\text{--}92 \times 79\text{--}105 \mu\text{m}$ have two spore wall layers: an evanescent hyaline $0.3\text{--}0.8 \mu\text{m}$ thick outer layer and a laminate orange brown to dark red brown $7.4\text{--}15.5 \mu\text{m}$ thick inner layer. The pigmentation of the subtending hypha is similar but often much lighter than that of the laminate spore wall layer. The spore size and color, spore wall structure, and features of the spore base and subtending hyphae separate this species from similar yellow-brown to brown spored species like *G. badium*, *G. glomerulatum*, and *G. brohultii*.

KEY WORDS — morphology, *Glomeromycota*, *Glomerales*, restinga

Introduction

The Tulasne brothers described the genus *Glomus* in 1844 for *G. macrocarpum* (the type species according to Clements & Shear 1931). Since then many other fungi with glomoid spores like those of *G. macrocarpum* have been named, so that now *Glomus* comprises >60 species. For many years *Glomus* was considered monophyletic based solely on morphological evidence. Morton & Redecker (2001), based on morphological and molecular data, erected the genera *Paraglomus* (*Paraglomeraceae*) and *Archaeospora* (*Archaeosporaceae*), whose members also formed glomoid spores. Schwarzott et al. (2001) inferred from phylogenetic SSU rDNA molecular analyses that *Glomus* is non-

monophyletic. Schüßler et al. (2001) subsequently erected a new phylum, *Glomeromycota*, in which *Glomus* was represented by three phylogenetically different groups: A, B, and C. Oehl & Sieverding (2004) transferred glomoid species with two spore walls (with the inner wall serving as a germinal wall) to their new genus *Pacispora*. Thereafter Walker & Schüßler (2004) transferred *Glomus spurcum* C.M. Pfeiff. et al. to the newly established genus *Diversispora* and order *Diversisporales*.

To date, glomoid glomerospores can be found in species representing *Archaeosporales*, *Diversisporales*, *Glomerales*, and *Paraglomerales*, making their identification difficult in the absence of molecular data. Morphological identification has been facilitated recently in a thorough review of glomoid spore-forming species (Oehl et al. 2011), which introduced helpful suggestions based on molecular data associated with morphological differences among the major phylogenetic clades in the *Glomeromycota*. Our description of *G. trufemii* is based exclusively on the new key morphological criteria, and the species fits the Oehl's et al. (2011) revised concept of *Glomus*.

Material & methods

Study area and sites

Samples were collected in the Parque Estadual das Dunas de Natal "Jornalista Luiz Maria Alves," the first Conservation Unit created in the Municipality of Natal, Rio Grande do Norte State and currently one of the largest urban conservation areas with dune vegetation in Brazil. The site is located at 5°46'S 35°12'W with soils analyzed as pH (H₂O) = 5.5–5.75, P = 2–4 mg kg⁻¹, 2,86 g dm³ of organic matter. The climate is tropical rainy (type Am of Köppen) with a short four-month dry period, with a 25.5 °C mean annual temperature and 1191 mm mean annual precipitation. In the sand dune ecosystem, the typical 'restinga' vegetation varies from herbaceous to shrubs and trees (Oliveira-Filho 1993, Oliveira-Filho & Carvalho 1993). The park is composed of mainly represented by tree species of the *Anacardiaceae*, *Bignoniaceae*, *Fabaceae*, *Myrtaceae*, and *Rubiaceae* (*Anacardium occidentale*, *Caesalpinia echinata*, *Campomanesia dichotoma*, *Chamaecrista ensiformis*, *Eugenia ligustrina*, *Guettarda platypoda*, *Myrcia guianensis*, *Myrciaria tenella*, *Tabebuia roseoalba*, *Tocoyena sellowiana*) and herbaceous species of *Poaceae*, *Cyperaceae*, *Asteraceae*, *Araceae*, and *Rubiaceae*.

Morphological analyses

Glomerospores and sporocarps were separated from the soil samples by sucrose gradient in plate dishes (Błaszczowski et al. 2006). Sporocarps and glomerospores were placed in dishes with water and separated under a dissecting microscope. Sporocarps were fragmented to study spore organization. Glomerospores were mounted on microscope slides either in water (to check unmodified characteristics of spore wall components and spore colour; Spain 1990) or permanently in PVLG (Omar et al. 1979) or PVLG with Melzer's reagent (Koske & Tessier 1983).

Terminology follows Oehl et al. (2011) and Furrázola et al. (2011). The spore denomination of Goto & Maia (2006) was used. Zeiss Axioskop compound microscopes with or without

Nomarski differential interference contrast (DIC) were used for detailed spore observations; digital images were taken with either an AxioCam camera and AxioVision (v. 3.1 software at 1300 × 1030 dpi) or Canon digital cameras.

Arbuscular mycorrhizal cultures

Bait cultures for the soils from sand dunes (Parque das Dunas de Natal) with spores and sporocarps were established in the greenhouse of the Universidade Federal do Rio Grande do Norte with *Zea mays* L. as host plant, using sterilized soil and 0.5 L pots.

Pure cultures of the new fungus were attempted by inoculating *Zea mays* seedlings germinated in autoclaved quartz sand with *G. trufemii* sporocarps isolated from sand dunes. After filling 0.5 L pots with sterilized sand dune soil, we formed a small hole in the center using a glass shaker and discharged spores and sporocarps into the hole using Pasteur pipettes. Then a *Z. mays* seedling was placed into the hole, covered with sand, and grown for one cycle of 3–4 months. The fungus did not produce spores in either bait cultures grown even for 9–12 months or pure cultures.

Taxonomy

Glomus trufemii B.T. Goto, G.A. Silva & Oehl, sp. nov.

FIGS. 1–9

MYCOBANK MB 561567

Differs from *G. brohultii* by more regularly shaped subtending hyphae that are pigmented further below the spore bases and darker colored orange to red brown spores with slightly thicker walls.


TYPE: Brazil, Rio Grande do Norte, near Natal, Parque das Dunas de “Natal”, sand dunes with herbaceous plant community on a convex hillside, 15.Feb.2010, B.T. Goto, in permanent slide (PVLG) (Holotype: UFRN1482; Isotype: UFRN1483, ZT Myc 15118).

ETYMOLOGY: *trufemii* = in honor of Sandra Farto Botelho Trufem, taxonomist, who long taught many young students and researchers in arbuscular mycorrhizal fungal taxonomy.

SPOROCARPS formed in aggregates (500–850 × 780–1200 μm), orange brown to dark red brown adherent to living or dead roots with hundreds to thousands of glomerospores; rarely singly in soils. Melzer’s reaction not observed. Peridium absent. Sporocarpic hyphae hyaline (3.8–)7.6–10.4(–12.8) μm without a septum and with interwoven arrangements.

GLOMEROSPORES formed terminally on hyphae, subglobose to elliptical (72–92 × 79–105 μm) or rarely globose (72–96 μm diam.), orange brown when young to dark red brown at maturity, perhaps darkening slightly to dark brown when ageing in soil.

SPORE WALL 7.4–15.5 μm thick overall, consisting of two layers (SWL1, SWL2); SWL1 thin (0.3–0.8 μm), hyaline and difficult to observe in mature spore, but generally visible in the subtending hyphal wall; SWL2 thick (7.4–15.5 μm), pigmented orange brown to dark red brown, laminated, smooth. Spore wall layers continuous with subtending hyphal wall layers. The pigmentation of SWL2 continues into the subtending hyphal wall, but this layer is often lighter



in color in the subtending hypha. swL1 and swL2 do not stain when exposed to Melzer's reagent.

SUBTENDING HYPHA (SH) generally present, single, straight or constricted, cylindrical to sharply curved, light yellow to dark yellow; 5.1–12.7 μm diam. (mean 8.8 μm) at the spore base, the hyphal wall 2.6–5.1 μm thick (mean 4.2 μm) near the spore base and tapering to approx. 1 μm at 15 μm from the base; occlusion formed by introverted swL2 thickening and often through an additional bridging septum arising from swL2, but sometimes the pore appearing partly open.

ABORTED GLOMEROSPORES frequently observed and forming terminally on hyphae, generally collapsed or subglobose to elliptical (35–56 \times 40–46 μm), light yellow to dark yellow, possibly darkening slightly to light brown when aging in soils. Spore wall layer (swL2) (<3.8 μm) not staining in Melzer's reagent.

GERMINATION structures unknown.

GLOMEROSPORE DEVELOPMENT was deduced from positively identified spores in aggregates found in different developmental stages. The hyaline hyphal wall layer differentiates into a hyaline, evanescent spore wall layer (swL1) and then a laminate layer (swL2) that becomes more pigmented with increasing numbers of developing sublaminae. After the glomerospores mature, their pore is closed by introverted thickening of swL2 and an additional bridging septum arising from the laminate wall layer.

ARBUSCULAR MYCORRHIZA FORMATION remains unknown.

DISTRIBUTION — So far, the new fungus was detected only in Brazil. Known in soil from sand dunes mainly harbouring herbaceous species of *Araceae*, *Asteraceae*, *Cyperaceae*, *Poaceae* and *Rubiaceae* (e.g.: *Acroceras zizanioides* (Kunth) Dandy, *Anthurium affine* Schott, *Aspilia procumbens* Baker, *Centratherum punctatum* Cass., *Cyperus maritimus* Poir., *C. meyenianus* Kunth, *Mitracarpus eichleri* K.Schum., *Raddia brasiliensis* Bertol., *Richardia grandiflora* (Cham. & Schltdl.) Steud.) growing in native, ecologically unstable restinga ecosystems in Parque das Dunas, Natal (Brazil).

Discussion

Glomus species are very common in sand dunes (Błaszowski 2010; Błaszowski et al. 2009a,b, 2010a,b) and several new species have been described recently (*G. africanum* Błasz. & Kovács, *G. achrum* Błasz. et al.,

FIGS. 1–9. *Glomus trufemii*. 1. Sporocarps in PVLG; note the numerous glomerospores in this part of sporocarp. 2–3. Fragment of sporocarp with healthy glomerospores and aborted spores. 4. Aborted spores with only one spore wall layer (swL2). 5. Septum formed by swL2. 6. Spores in different developmental stages. 7. Spore wall layers in mature spore (swL1&2). 8–9. Spore wall layers (swL1&2) continuous with subtending hyphal wall (SW) that form distinct pigmentation.

G. bistratum Błaszcz. et al., *G. indicum* Błaszcz. et al., *G. iranicum* Błaszcz. et al., *G. majewskii* Błaszcz., *G. perpusillum* Błaszcz. & Kovács). *Glomus trufemii* is readily distinguished from previously described sporocarpic *Glomus* species by glomerospore size and colour, spore wall structure, (including that at the spore base), and characters of the subtending hypha. Most of these species form colourless or pale-coloured spores singly or in loose aggregates (Błaszczowski et al. 2009a,b; 2010a,b) and thus are easy to distinguish from the sporocarpic dark-spored *G. trufemii*. Sporocarps of *G. trufemii* slightly resemble spore aggregates of *G. aggregatum* N.C. Schenck & G.S. Sm. and *G. fasciculatum* (Thaxt.) Gerd. & Trappe (Schenck & Smith 1982, Gerdemann & Trappe 1974). However, the *G. fasciculatum* swL2 stains in Melzer's reagent (vs. no reaction in *G. trufemii*). Spores of *G. aggregatum* generally are irregular in shape (vs. globose to subglobose) and have much thinner spore and subtending hyphal walls. The spore wall of *G. trufemii* is one of the thickest among the known glomeromycotan species with glomoid spores.


Several *Glomus* spp. form sporocarps without a peridium with similarly sized and coloured spores similarly distributed in a sporocarp: *G. ambisporum* G.S. Sm. & N.C. Schenck, *G. atrouva* McGee & Pattinson, *G. badium* Oehl et al., *G. boreale* (Thaxt.) Trappe & Gerd., *G. botryoides* F.M. Rothwell & Victor, *G. brohultii* Sieverd. & R.A. Herrera, *G. flavisporum* (M. Lange & E.M. Lund) Trappe & Gerd., *G. glomerulatum* Sieverd., *G. heterosporum* G.S. Sm. & N.C. Schenck, *G. invermaium* I.R. Hall, and *G. macrocarpum* Tul. & C. Tul. (Oehl et al. 2011). However, of these only *G. badium*, *G. brohultii*, and *G. glomerulatum* also form spores with subtending hyphae generally clearly lighter-coloured than the spore wall (Sieverd. 1987, Herrera-Peraza et al. 2003, Oehl et al. 2005, Oehl et al. 2011). In the other species, the wall color of both spore and subtending hyphae is similar or identical (Oehl et al. 2011).

Glomus glomerulatum produces intercalary yellow brown glomerospores having 2–4 subtending hyphae (vs only terminal spores with one subtending hypha in *G. trufemii*). Moreover, *G. trufemii* spores are darker (orange-brown to dark reddish brown) and have a significantly thicker wall.

Glomus badium forms irregularly sized 3-layered spores around an intrasporocarpic gleba, while *G. trufemii* generally produces regularly sized subglobose 2-layered spores in sporocarps lacking glebas.

Glomus trufemii and *G. brohultii* spores have a similar size and wall structure. Also, although *G. brohultii* was not originally described as forming sporocarps

FIGS. 10–18. *Glomus brohultii*. Bi-layered spores (swL1&2). 10. Glomerospores in PVLG; note the subtending hypha (SH) which is pigmented up to ca. 50 µm below the spore base before becoming subhyaline to hyaline. 11–12. Spores frequently have two subtending hyphae. 13. Mature spore with swL2 thickened at the spore base. 14. Irregular subtending hypha becoming hyaline in a rather short distance from the spore base (20–30 µm). 16. Hyaline subtending hypha with two layers (shL1&2). 17–18. Crushed spores with two spore wall layers (swL1&2).



or aggregates, sporocarps have been observed recently (Oehl, pers. obs.) during AMF diversity studies in cultivated white and yellow yam sites and adjacent natural savannas in West Africa (Tchabi et al. 2009), ruling out sporocarp or aggregate formation as a diagnostic character separating *G. trufemii* and *G. brohultii*. However, *G. brohultii* spores are paler and lack orange or red ones— yellow-brown to brown, not orange-brown to dark red-brown — and its spore walls are generally thinner. Furthermore, in *G. trufemii* the subtending hyphal pigmentation extends further below the spore base, both in spores formed in sporocarps or singly in the soil. Finally, the shape of the subtending hyphae is much more irregular in *G. brohultii* (Herrera-Peraza et al. 2003: this paper, Figs. 10–18).

Acknowledgements

The authors acknowledge, in special, Dr. Janusz Błaszowski (Department of Plant Protection, West Pomeranian University of Technology, Szczecin, Poland) and Dr. Ewald Sieverding (Institute for Plant Production and Agroecology in the Tropics and Subtropics, University of Hohenheim), for reviewing the manuscript and making helpful comments and suggestions and appreciate the corrections by Shaun Pennycook, Nomenclatural Editor, and suggestions by Lorelei L. Norvell, Editor-in-Chief. The study was supported by the Universidade Federal de Pernambuco which provided a grant to F. Oehl as ‘visiting professor’ and Programa de Pós-Graduação em Sistemática e Evolução that invited F. Oehl for a technical visit to Natal, RN.

Literature cited

- Błaszowski J. 2010. *Glomus majewskii*, a new species of arbuscular mycorrhizal fungi (*Glomeromycota*). Polish Botanical Journal 55(2): 265–270.
- Błaszowski J, Renker C, Buscot F. 2006. *Glomus drummondii* and *G. walkeri*, two new species of arbuscular mycorrhizal fungi (*Glomeromycota*). Mycol. Res. 110: 555–566. <http://dx.doi.org/10.1016/j.mycres.2006.02.006>
- Błaszowski J, Ryszka P, Oehl F, Koegel S, Wiemken A, Kovács GM, Redecker D. 2009a. *Glomus achrum* and *G. bistratum*, two new species of arbuscular mycorrhizal fungi (*Glomeromycota*) found in maritime sand dunes. Botany 87: 260–271. <http://dx.doi.org/10.1139/B08-138>
- Błaszowski J, Kovács GM, Balázs T. 2009b. *Glomus perpusillum*, a new arbuscular mycorrhizal fungus. Mycologia 101(2): 247–255. <http://dx.doi.org/10.3852/08-087>
- Błaszowski J, Wubet T, Harikumar VS, Ryszka P, Buscot F. 2010a. *Glomus indicum*, a new arbuscular mycorrhizal fungus. Botany 88: 132–143. <http://dx.doi.org/10.1139/B09-104>
- Błaszowski J, Kovács GM, Balázs T, Orłowska E, Sadravi M, Wubet T, Buscot F. 2010b. *Glomus africanum* and *G. iranicum*, two new species of arbuscular mycorrhizal fungi (*Glomeromycota*). Mycologia 102(6): 1450–1462. <http://dx.doi.org/10.3852/09-302>
- Clements FE, Shear CL. 1931. The genera of fungi. Hafner Publishing Co., New York, USA.
- Furrazola, E, Torres-Arias Y, Ferrer RL, Herrera RA, Berbara RLL, Goto BT. 2011. *Glomus crenatum*, a new ornamented species in the *Glomeromycetes* from Cuba. Mycotaxon 116: 143–149. <http://dx.doi.org/10.5248/116.143>
- Gerdemann JW, Trappe JM. 1974. The *Endogonaceae* in the Pacific Northwest. Mycologia Memoir No. 5. 76 p.

- Goto BT, Maia LC. 2006. Glomerospores, a new denomination for the spores of *Glomeromycota*, a group molecularly distinct from *Zygomycota*. *Mycotaxon* 96: 129–132.
- Herrera-Peraza RA, Ferrer RL, Sieverding E. 2003. *Glomus brohultii*: a new species in the arbuscular mycorrhiza forming *Glomerales*. *Journal of Applied Botany and Food Quality* 77: 37–40.
- Koske RE, Tessier B. 1983. A convenient, permanent slide mounting medium. *Mycol. Soc. Am. Newsl.* 34: 59.
- Morton JB, Redecker D. 2001. Two new families of *Glomales*, *Archaeosporaceae* and *Paraglomeraceae*, with two new genera *Archaeospora* and *Paraglomus*, based on concordant molecular and morphological characters. *Mycologia* 93(1): 181–195.
- Oehl F, Sieverding E. 2004. *Pacispora*, a new vesicular arbuscular mycorrhizal fungi genus in the *Glomeromycetes*. *Journal of Applied Botany* 78: 72–82.
- Oehl F, Redecker D, Sieverding E. 2005. *Glomus badium*, a new sporocarpic arbuscular mycorrhizal fungal species from European grasslands of higher soil pH. *J. Appl. Bot. Food Qual.* 79(1): 38–43.
- Oehl F, Silva GA, Goto BT, Sieverding E. 2011. *Glomeromycota*: three new genera, and glomoid species reorganized. *Mycotaxon* 116: 75–120. <http://dx.doi.org/10.5248/116.75>
- Oliveira-Filho AT. 1993. Gradient analysis of an area of coastal vegetation in the state of Paraíba, Northeastern Brazil. *Edinburgh Journal of Botany* 50(2): 217–236.
- Oliveira-Filho AT, Carvalho DA. 1993. Florística e fisionomia da vegetação no extremo norte do litoral da Paraíba. *Rev. Bras. bot.* 16(1): 115–130.
- Omar MB, Bolland L, Heather WA. 1979. A permanent mounting medium for fungi. *Bulletin of the British Mycological Society* 13: 31–32.
- Redecker D, Raab P, Oehl F, Camacho FJ, Courtecuisse R. 2007. A novel clade of sporocarp-forming species of glomeromycotan fungi in the *Diversisporales* lineage. *Mycological Progress* 6: 35–44. <http://dx.doi.org/10.1007/s11557-007-0524-2>
- Schenck NC, Smith GS. 1982. Additional new and unreported species of mycorrhizal fungi (*Endogonaceae*) from Florida. *Mycologia* 74: 77–92.
- Schüßler A, Schwarzott D, Walker C. 2001. A new fungal phylum, the *Glomeromycota*: phylogeny and evolution. *Micol. Res.* 105: 1413–1421. <http://dx.doi.org/10.1017/S0953756201005196>
- Schwarzott D, Walker C, Schüßler A. 2001. *Glomus*, the largest genus of the arbuscular mycorrhizal fungi (*Glomales*), is non-monophyletic. *Mol. Phylogenet. Evol.* 21: 190–197.
- Sieverding E. 1987. A VA mycorrhizal fungus, *Glomus glomerulatum* sp. nov., with two hyphal attachments and spores formed only in sporocarps. *Mycotaxon* 29: 73–79.
- Spain JL. 1990. Arguments for diagnoses based on unaltered wall structure. *Mycotaxon* 38: 71–76.
- Tchabi A, Burger S, Coyne D, Hountondji F, Lawouin L, Wiemken A, Oehl F. 2009. Promiscuous arbuscular mycorrhizal symbiosis of yam (*Dioscorea* spp.), a key staple crop in West Africa. *Mycorrhiza* 19: 375–392. <http://dx.doi.org/10.1007/s00572-009-0241-6>
- Tulasne LR, Tulasne C. 1844. Fungi nonnulli hypogaei, novi v. minus cognito act. *Gior. Bot. Ital.* 1(2(7–8)): 55–63.
- Walker C, Schüßler A. 2004. Nomenclatural clarifications and new taxa in *Glomeromycota*. *Micol. Res.* 108: 981–982. <http://dx.doi.org/10.1017/S0953756204231173>