

MYCOTAXON

<http://dx.doi.org/10.5248/120.231>

Volume 120, pp. 231–237

April–June 2012

A new species of *Postia* (*Basidiomycota*) from Northeast China

BAO-KAI CUI* & HAI-JIAO LI

Institute of Microbiology, P.O. Box 61, Beijing Forestry University,
Beijing 100083, China

*CORRESPONDENCE TO: baokaicui@yahoo.com.cn

ABSTRACT — A new polypore, *Postia subplacenta* sp. nov. from Northeast China, is described and illustrated. It is characterized by resupinate basidiocarps, angular pores (3–5 per mm) with a cream buff to pale cinnamon-buff pore surface, and cylindrical basidiospores (4.2–6.0 × 1.9–2.4 µm) that are negative in Melzer's reagent and Cotton Blue. A key to accepted species of Chinese *Postia* is supplied.

KEY WORDS — lignicolous, poroid, fungi, *Fomitopsidaceae*, taxonomy

Introduction

Postia Fr. (*Polyporales*, *Basidiomycota*) is an important genus of brown-rot fungi that are widely distributed in the northern hemisphere and mostly grow on gymnosperm wood. Some species of the genus are economically important, such as *Postia guttulata* (Peck) Jülich and *P. lactea* (Fr.) P. Karst., which have been used as medicine in China (Dai et al. 2009a).

Some mycologists consider *Postia* a synonym of *Oligoporus* Bref. (Gilbertson & Ryvarden 1987, Núñez & Ryvarden 2001, Ryvarden & Gilbertson 1994), while others regard it as an independent genus (Dai & Hattori 2007, Larsen & Lombard 1986, Niemelä et al. 2004, Renvall 1992, Wei & Dai 2006). The genus is characterized by an annual growth habit, a monomitic hyphal system with clamp connections, and thin-walled, allantoid to cylindrical or ellipsoid basidiospores. Twenty-eight species have previously been known from China (Cui et al. 2008, Dai 2009, Dai et al. 2003, 2004, 2007a,b, 2009b, Dai & Penttilä 2006, Li et al. 2007, 2008, Wang et al. 2009, 2011, Wei & Dai 2006, Wei & Qin 2010, Yuan & Dai 2008, Yuan et al. 2010).

During a study on the taxonomy and diversity of wood-rotting fungi in Northeast China, two *Postia* specimens were found that represented one previously undescribed species, which we propose as *P. subplacenta*. In addition, we provide an identification key to the *Postia* species thus far reported from China.

Materials & methods

The studied specimens were deposited in the herbarium of Beijing Forestry University (BJFC). The microscopic procedure follows Dai (2010) and Cui et al. (2011). In presenting the variation in the size of the spores, we exclude 5% of measurements (given in parentheses) from each end of the range. The following abbreviations are used: IKI = Melzer's reagent, IKI- = negative in Melzer's reagent, KOH = 5% potassium hydroxide, CB = Cotton Blue, CB- = acyanophilous, L = mean spore length (arithmetic average of all spores), W = mean spore width (arithmetic average of all spores), Q = variation in the L/W ratios between the specimens studied, n = number of spores measured from given number of specimens. Sections were studied at magnifications up to $\times 1000$ using a Nikon Eclipse E 80i microscope and phase contrast illumination. Drawings were made with the aid of a drawing tube. Special colour terms follow Petersen (1996).

Taxonomy

Postia subplacenta B.K. Cui, sp. nov.

FIG. 1

MYCOBANK MB 563784

Differs from *Postia placenta* in the cream-buff to pale cinnamon-buff pore surface, wider subcircular hyphae, smaller basidiospores, and absence of gloeopleuroous hyphae.

TYPE. — China. Jilin Province, Antu County, Changbaishan Nature Reserve, on fallen trunk of *Pinus*, 8.VIII.2011, Cui 10001 (holotype, BJFC).

ETYMOLOGY — *subplacenta* (Lat.) denotes a similarity with *Postia placenta*.

FRUITBODY — Basidiocarps annual, resupinate, corky when fresh, corky to fragile when dry, up to 5 cm long, 3 cm wide, and 2.5 mm thick at the center. Pore surface white to cream-buff when fresh, cream-buff to pale cinnamon-buff when dry; pores angular, 3–5 per mm; dissepiments thin, entire to lacerate. Margin indistinct, narrow to almost lacking. Subiculum cream to cream-buff, corky when dry, ≤ 0.3 mm thick. Tubes paler than pore surface, pale cream-buff, fragile when dry, ≤ 2.2 mm long.

HYPHAL STRUCTURE — Hyphal system monomitic; generative hyphae bearing clamp connections, IKI-, CB-; tissues unchanged in KOH.

SUBICULUM — Generative hyphae hyaline, thin- to mostly slightly thick-walled, occasionally branched, interwoven, 3–7 μm in diam.

TUBES — Generative hyphae hyaline, slightly thick- to distinctly thick-walled, with a wide to narrow lumen, occasionally branched, interwoven, 2–5 μm in diam. Cystidia and cystidioles absent. Basidia clavate with four sterigmata and a basal clamp connection, $12–18 \times 4–5 \mu\text{m}$; basidioles in shape similar to basidia, but slightly smaller.

SPORES — Basidiospores cylindrical, hyaline, thin-walled, smooth, IKI-, CB-, $(4.0–)4.2–6.0(–6.9) \times 1.9–2.4(–2.5) \mu\text{m}$, L = 5.07 μm , W = 2.11 μm , Q = 2.37–2.45 (n = 60/2).

TYPE OF ROT — Brown rot.

ADDITIONAL SPECIMEN EXAMINED — CHINA. HEILONGJIANG PROVINCE, YICHUN, Fenglin Nature Reserve, on stump of *Pinus*, 1.VIII.2011 Cui 9818 (BJFC).




FIG. 1. *Postia subplacenta* (holotype), microscopic structures.

a: Basidiospores. b: Basidia and basidioles. c: Hyphae from trama. d: Hyphae from subiculum.

REMARKS—*Postia subplacenta* resembles *P. placenta* (Fr.) M.J. Larsen & Lombard in having resupinate basidiocarps, similar pores, and cylindrical basidiospores; however, *P. placenta* has a salmon-pink pore surface, gloeopleurous hyphae, narrow subicular hyphae, and larger basidiospores ($5.5\text{--}7 \times 2\text{--}2.5 \mu\text{m}$; Gilbertson & Ryvarden 1987, Ryvarden & Gilbertson 1994).

Postia subplacenta is similar to *P. rancida* (Bres.) M.J. Larsen & Lombard, which differs in having rancid taste and larger basidiospores ($6\text{--}8 \times 2\text{--}3 \mu\text{m}$; Gilbertson & Ryvarden 1987).

Postia obliqua Y.L. Wei & W.M. Qin, recently described from Southwest China (Wei & Qin 2010), may be confused with *P. subplacenta* in its resupinate

basidiocarps and similar pores and basidiospores ($4.8\text{--}6.3 \times 2.0\text{--}2.5 \mu\text{m}$) but differs in its very large basidiocarps with oblique tubes that usually form imbricate pseudopilei and the presence of gloeopleurous-like hyphae.

Macroscopically, *Postia subplacenta* looks somewhat like *Oligoporus rennyi* (Berk. & Broome) Donk, which, however, has chlamydospores and produces basidiospores that are oblong-ellipsoid and thin- to slightly thick-walled (Ryvarden & Gilbertson 1994).

Key to species of *Postia* in China

1. Basidiocarps effused-reflexed, pileate or stipitate	2
1. Basidiocarps resupinate	23
2. Basidiocarps stipitate or substipitate	3
2. Basidiocarps effused-reflexed or pileate	4
3. Pores 1–3 per mm	<i>P. subundosa</i> Y.L. Wei & Y.C. Dai
3. Pores 3–5 per mm	<i>P. ceriflua</i> (Berk. & M.A. Curtis) Jülich
4. Basidiocarps with distinct grey to bluish tints	5
4. Basidiocarps white, cream, yellowish or brown	7
5. On angiosperm	<i>P. alni</i> Niemelä & Vampola
5. On gymnosperm	6
6. Basidiospores $<1.8 \mu\text{m}$ wide	<i>P. caesia</i> (Schrad.) P. Karst.
6. Basidiospores $>1.8 \mu\text{m}$ wide	<i>P. luteoceaesia</i> (A. David) Jülich
7. Basidiocarps becoming brown when bruised or when dry	8
7. Basidiocarps unchanged when bruised or dry	10
8. Gloeocystidia present	<i>P. leucomallella</i> (Murrill) Jülich
8. Gloeocystidia absent	9
9. Basidiospores $<1.6 \mu\text{m}$ wide	<i>P. lateritia</i> Renvall
9. Basidiospores $>1.7 \mu\text{m}$ wide	<i>P. fragilis</i> (Fr.) Jülich
10. Basidiocarps chalky when dry	<i>P. calcarea</i> Y.L. Wei & Y.C. Dai
10. Basidiocarps fragile, corky or woody hard when dry	11
11. Cystidia present	12
11. Cystidia absent	14
12. Cystidia thin-walled	<i>P. amurensis</i> Y.C. Dai & Penttilä
12. Cystidia thick-walled	13
13. Cystidia amyloid in Melzer's reagent	<i>P. pileata</i> (Parmasto) Y.C. Dai & Renvall
13. Cystidia inamyloid in Melzer's reagent	<i>P. balsamea</i> (Peck) Jülich
14. Pores bigger, 1–3 per mm	<i>P. undosa</i> (Peck) Jülich
14. Pores smaller, >3 per mm	15

15. Gloeocystidia present	16
15. Gloeocystidia absent	17
16. Hyphal pegs abundant	<i>P. gloeocystidiata</i> Y.L. Wei & Y.C. Dai
16. Hyphal pegs absent	<i>P. qinensis</i> Y.C. Dai & Y.L. Wei
17. Pileal surface more or less pink when fresh	18
17. Pileal surface never pink when fresh	19
18. Basidiospores >1.5 µm wide	<i>P. persicina</i> Niemelä & Y.C. Dai
18. Basidiospores <1.5 µm wide	<i>P. cana</i> H.S. Yuan & Y.C. Dai
19. Pileal surface zonate	<i>P. zebra</i> Y.L. Wei & Y.C. Dai
19. Pileal surface azonate	20
20. Basidiocarps mild, upper surface greyish brown	<i>P. tephroleuca</i> (Fr.) Jülich
20. Basidiocarps bitter, upper surface cream or yellow-brownish	21
21. Basidiospores >2 µm wide	<i>P. guttulata</i>
21. Basidiospores <2 µm wide	22
22. Basidiocarps woody hard when dry	<i>P. stipitica</i> (Pers.) Jülich
22. Basidiocarps fragile when dry	<i>P. lactea</i>
23. Basidiocarps becoming reddish to rusty brown when touched	<i>P. lateritia</i>
23. Basidiocarps unchanged when touched	24
24. Basidiospores mostly >2 µm wide, cystidia absent	25
24. Basidiospores mostly <2 µm wide, cystidia present	28
25. Basidiocarps salmon pink; basidiospores 2–3 µm wide	<i>P. placenta</i>
25. Basidiocarps white or cream; basidiospores 2–2.5 µm wide	26
26. Basidiocarps with rancid smell	<i>P. rancida</i>
26. Basidiocarps without rancid smell	27
27. Basidiocarps very large, gloeopleurous hyphae present	<i>P. obliqua</i>
27. Basidiocarps small, gloeopleurous hyphae absent	<i>P. subplacenta</i>
28. Cystidia thick-walled	<i>P. pileata</i>
28. Cystidia thin-walled	29
29. Pores 5–6 per mm; basidiospores mostly <1 µm wide	<i>P. simanii</i> (Pilát) Jülich
29. Pores 3–4 per mm; basidiospores mostly >1 µm wide	<i>P. hibernica</i> (Berk. & Broome) Jülich

Acknowledgements

Special thanks are due to Mr. Chang-Lin Zhao (China) for help in field collection. We express our gratitude to Drs. Michal Tomšovský (Czech Republic) and Yu-Lian Wei (China) who reviewed the manuscript. The research was financed by the National Natural Science Foundation of China (Project No. 31170018) and Program for New Century Excellent Talents in University (NCET-11-0585).

Literature cited

- Cui BK, Yuan HS, Dai YC. 2008. Wood-rotting fungi in eastern China 1. Polypores from Wuyi Mountains, Fujian Province. *Sydowia* 60: 25–40.
- Cui BK, Li HJ, Dai YC. 2011. Wood-rotting fungi in eastern China 6. Two new species of *Antrodia* (*Basidiomycota*) from Mt. Huangshan, Anhui Province. *Mycotaxon* 116: 13–20. <http://dx.doi.org/10.5248/116.13>
- Dai YC. 2009. A checklist of polypores in China. *Mycosistema* 28: 315–327.
- Dai YC. 2010. *Hymenochaetaceae* (*Basidiomycota*) in China. *Fungal Diversity* 45: 131–343. <http://dx.doi.org/10.1007/s13225-010-0066-9>
- Dai YC, Hattori T. 2007. *Postia japonica* (*Basidiomycota*), a new polypore from Japan. *Mycotaxon* 102: 113–118.
- Dai YC, Penttilä R. 2006. Polypore diversity of Fenglin Nature Reserve, northeastern China. *Annales Botanici Fennici* 43: 81–96.
- Dai YC, Häkkinen M, Niemelä T. 2003. Wood-inhabiting fungi in southern China 1. Polypores from Hunan Province. *Annales Botanici Fennici* 40: 381–393.
- Dai YC, Wei YL, Wang Z. 2004. Wood-inhabiting fungi in southern China 2. Polypores from Sichuan Province. *Annales Botanici Fennici* 41: 319–329.
- Dai YC, Cui BK, Huang MY. 2007a. Polypores from eastern Inner Mongolia, northeastern China. *Nova Hedwigia* 84: 513–520. <http://dx.doi.org/10.1127/0029-5035/2007/0084-0513>
- Dai YC, Cui BK, Yuan HS. 2007b. Notes on polypores from Gansu and Qinghai Province, northwest China. *Cryptogamie Mycologie* 28: 177–187.
- Dai YC, Yang ZL, Cui BK, Yu CJ, Zhou LW. 2009a. Species diversity and utilization of medicinal mushrooms and fungi in China (Review). *International Journal of Medicinal Mushrooms* 11: 287–302. <http://dx.doi.org/10.1615/IntJMedMushr.v11.i3.80>
- Dai YC, Yuan HS, Wang HC, Yang F, Wei YL. 2009b. Polypores (*Basidiomycota*) from Qin Mts. in Shaanxi Province, central China. *Annales Botanici Fennici* 46: 54–61.
- Gilbertson RL, Ryvarden L. 1987. North American polypores 2. *Fungiflora*, Oslo. 434–885 pp.
- Larsen MJ, Lombard FF. 1986. New combinations in the genus *Postia* Fr. (*Polyporaceae*). *Mycotaxon*. 26: 271–273.
- Li J, Xiong HX, Zhou XS, Dai YC. 2007. Polypores (*Basidiomycetes*) from Henan Province in central China. *Sydowia* 59: 125–137.
- Li J, Xiong HX, Dai YC. 2008. Polypores from Shennongjia Nature Reserve in Hubei Province, Central China. *Cryptogamie Mycologie* 29: 267–277.
- Niemelä T, Dai YC, Kinnunen J, Schigel DS. 2004. New and in North Europe rare polypore species (*Basidiomycota*) with annual, monomitic basidiocarps. *Karstenia* 44: 67–77.
- Núñez M, Ryvarden L. 2001. East Asian polypores 2. *Polyporaceae* s. lato. *Synopsis Fungorum* 14: 165–522.
- Petersen JH. 1996. Farvekort. The Danish Mycological Society's colour-chart. Foreningen til Svampekundskabens Fremme, Greve. 6 p.
- Renvall P. 1992. *Basidiomycetes* at the timberline in Lapland 4. *Postia lateritia* n. sp. and its rust-coloured relatives. *Karstenia* 32: 43–60.
- Ryvarden L, Gilbertson RL. 1994. European polypores 2. *Synopsis Fungorum* 7: 394–743.
- Wang B, Dai YC, Cui BK, Du P, Li HJ. 2009. Wood-rotting fungi in eastern China 4. Polypores from Dagang Mountains, Jiangxi Province. *Cryptogamie Mycologie* 30: 233–241.
- Wang B, Cui BK, Li HJ, Du P, Jia BS. 2011. Wood-rotting fungi in eastern China 5. Polypore diversity in Jiangxi Province. *Annales Botanici Fennici* 48: 237–246.

- Wei YL, Dai YC. 2006. Three new species of *Postia* (*Aphyllophorales, Basidiomycota*) from China. *Fungal Diversity* 23: 391–402.
- Wei YL, Qin WM. 2010. Two new species of *Postia* from China. *Sydowia* 62: 165–170.
- Yuan HS, Dai YC. 2008. Polypores from northern and central Yunnan Province, southwestern China. *Sydowia* 60: 147–159.
- Yuan HS, Dai YC, Wei YL. 2010. *Postia cana* sp. nov. (*Basidiomycota, Polyporales*) from Shanxi Province, northern China. *Nordic Journal of Botany* 28: 629–631.
<http://dx.doi.org/10.1111/j.1756-1051.2010.00849.x>