

---

# MYCOTAXON

<http://dx.doi.org/10.5248/120.239>

Volume 120, pp. 239–246

April–June 2012

---

## Three new *Caeoma* species on *Rosa* spp. from Pakistan

N.S. AFSHAN<sup>1\*</sup>, A.N. KHALID<sup>2</sup> & A.R. NIAZI<sup>2</sup>

<sup>1</sup>\*Centre for Undergraduate Studies & <sup>2</sup>Department of Botany, University of the Punjab,  
Quaid-e-Azam Campus, Lahore, 54590, Pakistan

\*CORRESPONDENCE TO: [pakrust@gmail.com](mailto:pakrust@gmail.com)

**ABSTRACT** — Three representatives of the anamorphic genus *Caeoma* — *C. ahmadii* on *Rosa microphylla*; *C. khanspurens* and *C. rosicola* on *Rosa webbiana*— are described as new rust species from Pakistan. This first report of *Caeoma* raises the number of known anamorphic rust genera from the country to five.

**KEY WORDS** — Khanspur, Mansehra, *Phragmidium*

### Introduction


The genus *Caeoma* Link is traditionally used for species having sori that lack obvious bounding structures and that produce catenulate spores with intercalary cells. This contrasts with the genus *Aecidium* Pers., which has a cup-shaped sorus with a well-developed peridium. Similar sori are found in the aecial state of *Melampsora* Castagne and the uredinia of *Chrysomyxa* Unger, *Coleosporium* Lév., and other genera (Cummins & Hiratsuka 2003). The aecia of *Phragmidium* are (usually) *Caeoma*-type with catenulate spores or (less often) *Uredo*-type (Petrova & Denchev 2004) with verrucose or echinulate aeciospores. Cummins & Hiratsuka (2003) refer to the anamorphic genus *Lecythea* Lévillé the *Phragmidium* species with aecia corresponding to *Caeoma* III of Sato & Sato (1985). Because Hennen et al. (2005) regard *Lecythea* as confusing and not in use, we consider *Caeoma* the appropriate anamorph for accommodating species with *Phragmidium* aecia.

So far, 22 genera and 417 species of rust fungi have been described or reported from Pakistan (Afshan et al. 2008a,b,c,d, 2009a,b, 2010a,b, 2011a,b; Sultan et al. 2008; Afshan & Khalid 2009; Khalid & Afshan 2009; Iqbal et al. 2008, 2009), including four genera of anamorphic rust fungi, *Aecidium*, *Monosporidium* Barclay, *Peridermium* (Link) J.C. Schmidt & Kunze, and *Uredo* Pers. Although no representatives of *Caeoma* have been reported previously from Pakistan, three new *Caeoma* species occurring on *Rosa* are described here.

### Materials & methods

Freehand sections of infected tissue and spores were mounted in lactophenol and gently heated to boiling point. The preparations were observed under a NIKON YS 100 microscope and photographed with a digipro-Labomed and a JSM5910 scanning electron microscope. Drawings of spores and paraphyses were made using a Camera Lucida (Ernst Leitz Wetzlar, Germany). An ocular micrometer was used for spore dimensions, with at least 25 spores measured for each spore stage. The rusted specimens have been deposited in the Botany Department herbarium, University of the Punjab, Lahore (LAH).

### Taxonomy



Figs. A–C: *Caeoma ahmadii* (holotype), SEM micrographs. A: Aecidium containing aeciospores and paraphyses. B: Aeciospores. C: Detail of an aeciospore showing verrucose wall ornamentation (paraphyses visible in background).

*Caeoma ahmadii* Afshan, Khalid & Niazi, sp. nov.

Figs. A–E


MYCOBANK MB 564295

Differs from *Phragmidium tuberculatum* in larger aeciospores that are hyaline with orange-yellow contents.

TYPE: PAKISTAN, KHYBER-PAKHTUNKHAWAH (KPK), Ayubia National Park, at 2135 m a.s.l., on *Rosa microphylla* Desf., stage I, 18 June, 2008, N.S. Afshan & A.N. Khalid. NSA #151A (Holotype, LAH NSA1114).

ETYMOLOGY: Named after the world renowned mycologist, Sultan Ahmad.

SPERMOGONIA, UREDINIA and TELIA not found. AECIA hypophyllous, on leaves, bright yellow to orangish yellow, rounded or oblong, scattered, 0.1–0.3 × 0.09–0.10 mm. PARAPHYSES numerous, erect to suberect, clavate, hyaline with light yellow granules, 50–70 µm long and 8–12 µm wide. AECIOSPORES globose to subglobose or ellipsoid to broadly ellipsoid, (23–)26–31 × 28–36 µm; wall 1–1.5 µm thick, hyaline with orange–yellow contents, densely verrucose, with 5–8 scattered pores.




Figs. D–E: *Caeoma ahmadii* (holotype), lucida drawings.

D: Aeciospores showing wall ornamentation. Scale bar = 9 µm.

E: Apices of paraphyses. Scale bar = 12 µm.

COMMENTS: Aeciospores of *C. ahmadii* are larger than those of *Phragmidium tuberculatum* J.B. Müll. (18–24 × 20–30 µm), *Ph. mucronatum* (Pers.) Schtdl. (17–21 × 20–28 µm), *Ph. montivagum* Arthur (16–19 × 21–26 µm), *Ph. rosae-arkansanae* Dietel (16–23 × 23–27 µm), *Ph. speciosum* (Fr.) Burrill (16–24 ×

24–35  $\mu\text{m}$ ), and *Ph. rosae-pimpinellifoliae* Dietel (15–20  $\times$  18–27  $\mu\text{m}$ ). They also differ in color and wall ornamentation from those of *Ph. tuberculatum*.



FIGS. F–G: *Caecoma khanspurensis* (holotype), lucida drawings.

F: Apices of clavate paraphyses. Scale bar = 20  $\mu\text{m}$ . G: Mature aeciospores. Scale bar = 10  $\mu\text{m}$ .

*Caecoma khanspurensis* Khalid, Afshan & Niazi, sp. nov.

FIGS. F–G

MYCOBANK MB 564317

Differs from *Phragmidium mucronatum* and *Ph. tuberculatum* in larger aeciospores with verrucose wall ornamentation.

TYPE: PAKISTAN, KHYBER-PAKHTUNKHAWAH (KPK), Helipad, Khanspur, at 2135 m a.s.l., on *Rosa webbiana* Wall. ex Royle, stage I, 23 May 2006, N.S. Afshan & A.N. Khalid. NSA #03 (Holotype, LAH NSA1116).

ETYMOLOGY: Named after the type locality, Khanspur.

SPERMOGONIA, UREDINIA and TELIA not found. AECIA hypophyllous or caulicolous, on the veins, stems and petioles, scattered, rounded or oblong, 0.3–0.9  $\times$  0.09–0.10 mm, orange yellow when fresh, surrounded by clavate paraphyses. AECIOSPORES globose, subglobose or ellipsoid to obovoid, pale yellow to orange-yellow, verrucose, occasionally with a plateau-shaped base, 23–32  $\times$  23–41  $\mu\text{m}$ , borne on one or two basal cells. PARAPHYSES clavate, hyaline, 70–105  $\mu\text{m}$  long, 8–12  $\mu\text{m}$  wide, with uniformly thin walls.

COMMENTS: Aeciospores of *C. khanspurensis* are larger than those of *Phragmidium mucronatum* (17–21  $\times$  20–28  $\mu\text{m}$ ) and *Ph. tuberculatum* (18–24

× 20–30 µm); their verrucose ornamentation further distinguishes them from the aeciospores of *Ph. mucronatum*.

The broader aeciospores of *C. khanspurens* differ from those of *C. rosicola* (reported on the same host), characterized by verrucose to echinulate, 19–28 µm broad aeciospores.

Aeciospores with a verrucose wall ornamentation with a plateau-shaped base distinguish *C. khanspurens* from *Phragmidium rosae-pimpinellifoliae* with smaller aeciospores (15–20 × 18–27 µm).




FIG. H: *Caeoma rosicola*, lucida drawing. Aeciospores. Scale bar = 10 µm.

***Caeoma rosicola*** Afshan, Niazi & Khalid, sp. nov.

FIGS. H–J

MYCOBANK MB 564318

Differs from other *Caeoma* species in larger aeciospores and lack of paraphyses.


TYPE: PAKISTAN, KHYBER-PAKHTUNKHAWAH (KPK), Helipad, Khanspur, at 2135 m a.s.l., on *Rosa webbiana* Wall. ex Royle, stage I, 23 May 2006, N.S. Afshan & A.N. Khalid. NSA #02 (Holotype, LAH NSA1117).

ETYMOLOGY: Named after the host genus, *Rosa*.

SPERMOGONIA, UREDINIA and TELIA not found. AECIA hypophyllous or petiocolous, mostly on the veins, petioles, branches and fruits, causing malformations, yellowish orange to bright orange, scattered. PARAPHYSES not observed. AECIOSPORES globose to subglobose or ellipsoid to ovoid; wall hyaline with orange yellow contents, echinulate to verrucose, 19–28 × 21–41 µm, germ pores 1–4, mostly equatorial.

COMMENTS: Larger aeciospores and the lack of paraphyses differentiate *C. rosicola* from other rust species reported on *Rosaceae*: *Phragmidium rosae-pimpinellifoliae* (15–20 × 18–27 µm), *Ph. rosae-arkansanae* (16–23 × 23–27 µm),





FIGS. I–J: *Caeoma rosicola* (holotype), SEM micrographs. I: Aeciospores. J: Detail of aeciospore showing echinulate wall ornamentation.

*Ph. mucronatum* (17–21 × 20–28 µm), and *Ph. montivagum* (16–19 × 21–26 µm). *Phragmidium montivagum* is further distinguished by aeciospores with 8 scattered germ pores.

#### Acknowledgements

We are highly obliged to Higher Education Commission (HEC) of Pakistan for providing financial support. We sincerely thank Dr. Marcin Piątek (W. Szafer Institute of Botany, Polish Academy of Sciences, Poland) and Dr. Omar Paino Perdomo (Dominican Society of Mycology Santo Domingo, Dominican Republic) for their valuable suggestions to improve the manuscript and acting as presubmission reviewers.

#### Literature cited

- Afshan NS, Khalid AN. 2009. New records of *Puccinia* and *Pucciniastrum* from Pakistan. *Mycotaxon* 108: 137–146. <http://dx.doi.org/10.5248/108.137>
- Afshan NS, Berndt R, Khalid AN, Niazi AR. 2008a. New graminicolous rust fungi from Pakistan. *Mycotaxon* 104: 123–130.
- Afshan NS, Khalid AN, Javed H. 2008b. Further additions to the rust flora of Pakistan. *Pakistan Journal of Botany* 40(3): 1285–1289.
- Afshan NS, Khalid AN, Niazi AR. 2008c. New records of graminicolous rust fungi from Pakistan. *Pakistan Journal of Botany* 40(3): 1279–1283.
- Afshan NS, Khalid AN, Niazi AR. 2008d. New records and distribution of rust fungi from Pakistan. *Mycotaxon* 105: 257–267.
- Afshan NS, Iqbal SH, Khalid AN, Niazi AR. 2009a. A new anamorphic rust fungus with a new record of *Uredinales* from Azad Kashmir, Pakistan. *Mycotaxon* 112: 451–456. <http://dx.doi.org/10.5248/112.451>
- Afshan NS, Khalid AN, Iqbal SH, Niazi AR, Sultan A. 2009b. *Puccinia subepidermalis* sp. nov. and new records of rust fungi from Pakistan. *Mycotaxon* 110: 173–182. <http://dx.doi.org/10.5248/110.173>
- Afshan NS, Khalid AN, Niazi AR. 2010a. Three new species of rust fungi from Pakistan. *Mycological Progress* <http://dx.doi.org/10.1007/s11557-010-0655-8>
- Afshan NS, Khalid AN, Iqbal SH, Niazi AR, Sultan A. 2010b. *Puccinia anaphalidis-virgatae*, a new species and a new variety of rust fungi from Fairy Meadows, Northern Pakistan, *Mycotaxon* 112: 483–490. <http://dx.doi.org/10.5248/112.483>
- Afshan NS, Khalid AN, Iqbal SH, Niazi AR. 2011a. *Puccinia* species new to Azad Jammu & Kashmir, Pakistan, *Mycotaxon*: 116: 175–182.
- Afshan NS, Khalid AN, Niazi AR, Iqbal SH. 2011b. New records of *Uredinales* from Fairy Meadows, Pakistan. *Mycotaxon*: 115: 203–213. <http://dx.doi.org/10.5248/115.203>
- Cummins GB, Hiratsuka Y. 2003. Illustrated genera of rust fungi. Third ed. The American Phytopathological Society. APS Press, St. Paul, MN.
- Hennen JF, Figueiredo MB, Carvalho AA, Hennen PG. 2005. Catalogue of the species of plant rust fungi (*Uredinales*) of Brazil. FAPESP, CNPq, FAPERJ, NSF, USDA.
- Iqbal SH, Khalid AN, Afshan NS, Niazi AR. 2008. Rust Fungi on *Saccharum* species from Pakistan. *Mycotaxon* 106: 219–226.
- Iqbal SH, Afshan NS, Khalid AN, Niazi AR, Sultan A. 2009. Additions to the rust fungi of Fairy Meadows, Northern Areas of Pakistan. *Mycotaxon* 109: 1–7. <http://dx.doi.org/10.5248/109.1>

- Khalid AN, Afshan NS. 2009. Additions to the graminicolous rust fungi of Pakistan. *Mycotaxon* 108: 175–183. <http://dx.doi.org/10.5248/108.175>
- Petrova RD, Denchev CM. 2004. A taxonomic study of *Phragmidiaceae* (*Uredinales*) in Bulgaria. *Mycologia Balcanica* 1: 95–115.
- Sato T, Sato S. 1985. Morphology of the aecia of the rust fungi. *Trans. Br. Mycol. Soc.* 85: 223–238.
- Sultan MA, Haq I, Khalid AN, Mukhtar H. 2008. Two new anamorphic rust fungi from northern areas of Pakistan. *Mycotaxon* 105: 23–27.