
MYCOTAXON

<http://dx.doi.org/10.5248/121.153>

Volume 121, pp. 153–157

July–September 2012

***Pholiota virescens*, a new species from China**

EN-JING TIAN & TOLGOR BAU*

Engineering Research Center of Chinese Ministry of Education for Edible and Medicinal Fungi,
Jilin Agricultural University, Changchun 130118, China

*CORRESPONDENCE TO: junwusuo@126.com

ABSTRACT — *Pholiota virescens*, a new species of *Pholiota* is described from China. It is characterized by its pileus with the green tone at maturity, and its prominent and conspicuously projecting pleurocystidia with thin wall.

KEY WORDS — *Agaricales*, *Strophariaceae*, taxonomy

Introduction

Pholiota (Fr.) P. Kumm. (*Agaricales*, *Strophariaceae*), established by Kummer in 1871, is a widespread genus (especially in the North Temperate Zone) with about 150 species worldwide (Kirk et al. 2008; Smith & Hesler 1968; Holec 2001; Noordeloos 2011).

There have been 55 taxa of *Pholiota* recorded in China (Tian & Bau 2004, 2005, 2011a,b; Bau et al. 2005), among which 20 species are edible (Dai et al. 2010) and 9 species are medicinal (Dai & Yang 2008). Recently, an additional new species was discovered during the research on the genus based on morphological examinations of collections.

Materials & methods

Specimens were examined by traditional taxonomic methods (Singer 1975). 5% KOH solution and Melzer's reagent were used as the mountants when examining the microstructure. Morphological characteristics of the species were described and illustrated according to the observations on the materials. Colour description for the new species refer to Kornerup & Wanscher (1978). The specimen studied is now deposited in the Herbarium of Mycology of Jilin Agricultural University (HMJAU).

Taxonomy

Pholiota virescens T. Bau & E.J. Tian, sp. nov.

PLATE 1, FIGURE 1

MYCOBANK MB563114

Differs from *Pholiota velaglutinosa* by the green tone of its mature pileus and by the thin-walled pleurocystidia.



PLATE 1 *Pholiota virescens*.
Habit.

TYPE: China, Jilin Province, Hunchun City, Shengli Village, 5 km to the east, on decayed hardwood, 10 July 2010, Tolgor Bau & En-Jing Tian (Holotype, HMJAU 22498).

ETYMOLOGY: The species epithet *virescens* refers to the pileus color becoming green with maturity.

PILEUS 2.8–6.4 cm broad, convex, then becoming plane, even at the margin, viscid, dull vinaceous-brown over the disc at first, but becoming dull green towards the margin, glabrous except for brown floccose veil-remnants adhering along the margin. LAMELLAE adnate, near avellaneous, becoming dull cinnamon-brown to dull greenish-brown at maturity, dense, medium broad, edges even. STIPE 2–4.5 cm long, 0.4–0.5 cm thick, equal, solid, white above, becoming pale vinaceous to brownish below, dingy brown at base, with scattered patches of fibrils to fibrillose-squamulose below; Veil brown, fibrillose, evanescent, leaving an evanescent ring. CONTEXT thin, brownish, odor and taste mild. SPORE PRINT dull cinnamon-brown.

SPORES $6.2\text{--}7.2(-7.7) \times 3.8\text{--}4.3(-5.3) \mu\text{m}$, ellipsoid, oblong to ovate in face view, phaseoliform in side view, smooth, pale rusty to cinnamon-brown in KOH, slightly paler in Melzer's reagent, germ pore minute. BASIDIA $17\text{--}20 \times 5\text{--}6 \mu\text{m}$, (2-)4-spored, narrowly clavate, hyaline in KOH. CHRYSOCYSTIDIA absent. PLEUROCYSTIDIA $50\text{--}80 \times 11\text{--}15 \mu\text{m}$, fusoid-ventricose with obtuse to slightly acute apex, upper part elongated (up to $44 \mu\text{m}$), wall thin, smooth, content homogeneous to collapsed, hyaline to tawny in KOH. CHEILOCYSTIDIA $24\text{--}44 \times 8.5\text{--}10 \mu\text{m}$, clavate, irregularly clavate to fusoid-ventricose, thin-walled, smooth, content homogeneous, hyaline to pale yellow in KOH. GILL TRAMA of parallel to subparallel hyphae hyaline in KOH and with smooth




FIGURE 1 *Pholiota virescens*.

a, Pleurocystidia; b, Basidia; c, Basidiospores; d, Cheilocystidia; e, Pileipellis.

Bars = 10 µm.

walls, the cells inflated, up to 25 µm in diam.; subhymenium of gelatinized, interwoven hyphae. PILEUS CUTIS an ixocutis of hyphae 2.5–5 µm in diam., thin-walled, smooth to slightly encrusted, yellowish to pale tawny in KOH.

CONTEXT hyphae thin-walled, smooth, hyaline to yellowish in KOH. CLAMP CONNECTIONS present.

HABIT—Scattered on dead wood of hardwoods in summer.

COMMENTS—This species belongs to the subgenus *Flammuloides* (Smith & Hesler 1968) based on its viscid pileus, prominent and conspicuously projecting cheilo- and pleurocystidia, and typically gelatinous subhymenium.

Pholiota virescens is similar to *P. velaglutinosa* A.H. Sm. & Hesler in young pileus color and basidiospore size and shape (Smith & Hesler 1968), but differs in the green tone in the mature pileus and the thin-walled pleurocystidia. It is also close to *P. vinaceobrunnea* A.H. Sm. & Hesler (Smith & Hesler 1968; Tian & Bau 2011a), which differs by its thick-walled pleurocystidia. *Pholiota lubrica* (Pers.) Singer is also similar, but its pileus is not dull vinaceous-brown but bright orange-brown (Noordeloos 1999; Holec 2001; Jacobsson 1990). *Pholiota velaglutinosa*, *P. vinaceobrunnea*, and *P. lubrica* are all known from China (Tian & Bau 2004, 2011a).

Acknowledgments

This study was supported by the National Natural Science Foundation of China (No. 31070013). We thank Dr. Takahito Kobayashi of Hokkaido University Museum, Japan, Dr. Jan Holec of National Museum, Mycological Department, Czech Republic and Dr. T.H. Li of Guangdong Institute of Microbiology, China, for revising the manuscript patiently and meticulously.

Literature cited

- Bau T, Tian EJ, Wang H. 2005. *Strophariaceae* of China (I) *Pholiota*. Journal of Fungal Research 3(3): 1–50 (in Chinese).
- Dai YC, Yang ZL. 2010. A revised checklist of medicinal fungi in China. Mycosystema 27: 801–824 (in Chinese).
- Dai YC, Zhou LW, Yang ZL. 2010. A revised checklist of edible fungi in China. Mycosystema 29: 1–21 (in Chinese).
- Holec J. 2001. The genus *Pholiota* in central and western Europe. Libri botanici 20: 1–220.
- Jacobsson S. 1990. *Pholiota* in northern Europe. Windahlia 19: 1–86.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. Ainsworth & Bisby's dictionary of the fungi. 10th edition. CAB International, Wallingford (UK). 771 p.
- Kornerup A, Wanscher JH. 1978. Methuen handbook of colour, third edition. Eyre Methuen Ltd., London. 252 p.
- Noordeloos ME. 1999. *Pholiota* (Fr.) Kumm., Führ. Pilzk.: 84. 1871. 80–107, in: C Bas et al. (eds). Flora Agaricina Neerlandica, vol. 4. Balkema, Rotterdam & Brookfield.
- Noordeloos ME. 2011. *Strophariaceae* s.l. Fungi Europaei, vol. 13. Edizioni Candusso, Alassio. 648 p.
- Singer R. 1975. The *Agaricales* in modern taxonomy (3rd ed.). J.Cramer, Vaduz. 912 p.
- Smith AH, Hesler LR. 1968. The North American species of *Pholiota*. Hafner Publishing Company, New York. 402 p.

- Tian EJ, Bau T. 2004. Known species of *Pholiota* (s. l.) from China and their distribution. *Journal of Fungal Research* 2(1): 25–34 (in Chinese).
- Tian EJ, Bau T. 2005. A new record of *Pholiota* from China. *Mycosystema* 24(2): 310–311 (in Chinese).
- Tian EJ, Bau T. 2011a. New records of *Pholiota* collected from Changbai Mt. in China. *Mycosystema* 30(3) : 408–413 (in Chinese).
- Tian EJ, Bau T. 2011b. New Chinese records of *Pholiota* collected from Inner Mongolia and Heilongjiang. *Mycosystema* 30(4) : 649–652 (in Chinese).