

---

# MYCOTAXON

<http://dx.doi.org/10.5248/121.327>

Volume 121, pp. 327–331

July–September 2012

---

## ***Phytophthora castaneae*, the correct name for *P. katsurae* nom. nov. superfl.**

SHAUN R. PENNYCOOK

*Manaaki Whenua Landcare Research, Private Bag 92 170, Auckland, New Zealand*

CORRESPONDENCE TO: [PennycookS@LandcareResearch.co.nz](mailto:PennycookS@LandcareResearch.co.nz)

**ABSTRACT** — *Phytophthora katsurae* was proposed as a nom. nov. for *P. castaneae* on the assumption that the replaced name was an illegitimate later homonym. This assumption was based on the invalid nomen nudum publication of “*Phytophthora castaneae*” in a host-pathogen index as a synonym of *Mycelophagus castaneae*, and an erroneous interpretation of Clements & Shear’s listing of the type of the genus *Mycelophagus*. There is no valid earlier homonym. *Phytophthora castaneae* is the correct legitimate name for the taxon causing trunk rot of *Castanea crenata*, and *P. katsurae* is an illegitimate superfluous name.

**KEY WORDS** — chestnut disease, ink disease, nomenclature, taxonomy

### **Introduction**

In 1976, Katsura & Uchida described a new *Phytophthora* species, *P. castaneae*, causing trunk rot of *Castanea crenata* in Japan (Katsura 1976). Subsequently, Ko & Chang (1979) claimed that *P. castaneae* was an illegitimate later homonym, and published a replacement name, *P. katsurae*, which has been used uncritically in taxonomic and phytopathological literature since that time (Ko & Arakawa 1980, Ho 1981, Ooka & Uchida 1984, Stamps 1985, Chang & Shu 1988, Liyanage & Wheeler 1989, Stamps et al. 1990, Oudemans & Coffey 1991, Uchida et al. 1992, Erwin & Ribeiro 1996, Elliott et al. 2004, Ristaino 2011). In this paper, I survey the relevant literature and demonstrate that Ko & Chang were mistaken — there is no earlier homonym and the correct name for this taxon is the legitimate original name, *P. castaneae*.

### **Ko & Chang’s major error**

Ko & Chang (1979: 841) presented a plausible case for the illegitimacy of *P. castaneae* Katsura & K. Uchida as a later homonym, but the basis of their case was the assumption that “The name of the type species [of *Mycelophagus*] was

consequently changed to *Phytophthora castaneae* (Mangin) Clements et Shear (7).” The referral of this citation to their LITERATURE CITED (7) [= MacFarlane (1968)] is puzzling, since MacFarlane (1968: 683) used the name “*Phytophthora castaneae*” in a host–pathogen index as a nomen nudum cross-reference to *Mycelophagus castaneae*, with no mention of Clements & Shear, or any other nomenclatural author. Moreover, Ko & Chang’s attribution of the combination to Clements & Shear is a misreading or misunderstanding of Clements & Shear (1931: 241), where a list of synonyms of *Phytophthora* includes the monotypic genus *Mycelophagus* L. Mangin (Mangin 1903), with its type listed as “*M. castaneae* Mang.” Clements & Shear did not make an explicit combination of the epithet in *Phytophthora*; and their citation of “*M. castaneae*” does not constitute a definite association of the epithet with the genus name *Phytophthora* or its abbreviation, as required for a valid combination (McNeill et al. 2006: Art. 33.1). The combination “*Phytophthora castaneae* (Mangin) Clements et Shear” is neither illegitimate (as Ko & Chang 1979 asserted, without explanation) nor invalid — it is non-existent!

### The correct nomenclature

***Phytophthora castaneae*** Katsura & K. Uchida, Trans. Mycol. Soc. Japan 17: 241. 1976.  
 = *Phytophthora katsurae* W.H. Ko & H.S. Chang, Mycologia 71:  
 841. 1979, nom. nov., nom. illegit. [superfluous].

In the absence of any earlier homonym, the name *P. castaneae* Katsura & K. Uchida is legitimate and the earliest available name for the taxon. Consequently, the nomen novum *P. katsurae* proposed by Ko & Chang (1979) is superfluous and therefore illegitimate (McNeill 2006: Art. 52.1).

Very few publications have avoided Ko & Chang’s error and its erroneous nomenclatural consequences. Uchida (1976) and Newhook et al. (1978: 7) both presented the taxon as *P. castaneae*, but this predated the 1979 publication of *P. katsurae*. Index of Fungi 4(20): 514. 1980 recognised (in entries for *P. castaneae* and *P. katsurae*) that the *Phytophthora* combination based on *M. castaneae* was not proposed in Clements & Shear (1931), and stated that it was first proposed by MacFarlane (1968: 683) as an invalid combination lacking any reference to the author and place of publication of the basionym (McNeill et al. 2006: Art. 33.4). Miyata & Odagiri (1982a,b) used the name *P. castaneae*, and Dick (2001: 498, 499) listed *P. katsurae* as a synonym, cross-referenced to the main entry of *P. castaneae* [misspelt and misattributed as “*P. castanae* [sic] Katsura & S. [sic] Uchida”]. Index of Fungi 6(16): 868. 1998 listed *P. katsurae* as a superfluous synonym of *P. castaneae*.

### The taxonomy and nomenclature of *Mycelophagus castaneae*

*Mycelophagus castaneae* L. Mangin, C.R. Hebd. Séanc. Acad. Sci., Paris 136: 472. 1903.

“*Phytophthora castaneae*” H.H. MacFarl., Rev. App. Mycol., Pl. Host-Path. Index vv. 1–40 (1922–61): 683. 1968, nom. inval. [no basionym author and reference].

“*Phytophthora castaneae*” W.H. Ko & H.S. Chang, Mycologia 71: 841. 1979, nom. inval. [accepted only as an illegitimate synonym of *M. castaneae*].

The French protologue of *M. castaneae* was reproduced and translated into English in Waterhouse (1970: 54–55) under the heading “Names not taken into *Phytophthora*.” Waterhouse (1963: 18) concluded that the genus and species were “most probably *Phytophthora* sp. but impossible to tell from the description.” Although its association with “ink disease of chestnut” in Europe (Blin 1922, Schell 1922, Waterhouse 1963) might suggest a possible relationship with either *P. cambivora* or *P. cinnamomi* (Erwin & Ribeiro 1996: 260, Vannini & Vettraiño 2001, Vettraiño et al. 2005), it is preferable to accept Waterhouse’s expert opinion that *Mycelophagus castaneae* should be treated as a name of dubious application.

Index of Fungi 6(16): 868. 1998 interpreted the publication by Ko & Chang (1979) of the fictitious combination “*Phytophthora castaneae* (Mangin) Clements et Shear” as constituting an unintentional but valid combination by Ko & Chang, which would be an illegitimate later homonym of *P. castaneae* Katsura & K. Uchida. The Ko & Chang paper certainly appears to contain all the elements required for a valid comb. nov., although the full and direct basionym reference is made via a LITERATURE CITED reference, contrary to an ICBN recommendation (McNeill et al. 2006: Rec. 33A.1). However, Ko & Chang explicitly accepted the binomial only as an illegitimate synonym of *Mycelophagus castaneae*, which invalidates any putative new combination (McNeill et al. 2006: Art. 34.1(c)).

In a further concatenation of misinformation, Ristaino (2011) stated that *P. katsurae* was “a replacement name for *P. castaneae* (L. Mangin) Clements & Shear (1931) and for *P. castaneae* Katsura & Uchida (1976), which are both considered illegitimate names.” Neither Ko & Chang (1979) nor any other author has ever suggested any taxonomic connection between Mangin’s and Katsura & Uchida’s species, the putative Clements & Shear combination is non-existent, and *P. castaneae* Katsura & K. Uchida is a legitimate name.

### Acknowledgments

This paper is dedicated to the memory of my late colleague, Ross Beever, who first posed the question, and who subjected the arguments to his characteristically intense critical analysis and questioning. I thank Stan Bellgard and Bevan Weir (Landcare

Research, Auckland) for assistance with literature and for their comments and criticisms of an earlier draft, and Joost Stalpers (CBS, Utrecht) and Scott Redhead (Systematic Mycology and Botany, Agriculture and Agri-Food Canada, Ottawa) for acting as pre-submission referees.

#### Literature cited

- Blin H. 1922. La maladie dite "de l'encre" des châtaigniers. *La Nature* 50(2534): 282–284. [Abstract in: *Review of Applied Mycology* 2: 188–189. 1923.]
- Chang HS, Shu IM. 1988. Oospore germination of homothallic *Phytophthora* species and the identity of *Phytophthora heveae* isolates from Taiwan. *Botanical Bulletin of Academia Sinica, Taiwan* 29: 15–21.
- Clements FE, Shear CL. 1931. *The genera of fungi*. 2nd edition. New York, H.W. Wilson Company. 496 p.
- Dick MW. 2001. *Straminipilous fungi*. Dordrecht, Kluwer Academic Publishers. 670 p.
- Elliott ML, Broschat TK, Uchida JY, Simone GW. 2004. *Compendium of ornamental palm diseases and disorders*. St. Paul MN, APS Press. 69 p.
- Erwin DC, Ribeiro OK. 1996. *Phytophthora* diseases worldwide. St. Paul MN, APS Press. 562 p.
- Ho HH. 1981. Synoptic keys to the species of *Phytophthora*. *Mycologia* 73: 705–714. <http://dx.doi.org/10.2307/3759497>
- Katsura K. 1976. Two new species of *Phytophthora* causing damping-off of cucumber and trunk rot of chestnut. *Transactions of the Mycological Society of Japan* 17: 238–242.
- Ko WH, Arakawa CK. 1980. Oospore germination of *Phytophthora katsurae*. *Transactions of the Mycological Society of Japan* 21: 215–219.
- Ko WH, Chang HS. 1979. *Phytophthora katsurae*, a new name for *P. castaneae*. *Mycologia* 71: 840–844. <http://dx.doi.org/10.2307/3759198>
- Liyanage NIS, Wheeler BEJ. 1989. *Phytophthora katsurae* from cocoa. *Plant Pathology* 38: 627–629. <http://dx.doi.org/10.1111/j.1365-3059.1989.tb01463.x>
- MacFarlane HH (compiler). 1968. *Review of applied mycology: plant host–pathogen index to volumes 1–40 (1922–1961)*. Kew, Commonwealth Mycological Institute. 820 p.
- McNeill J, Barrie FR, Burdet HM, Demoulin V, Hawksworth DL, Marhold K, Nicolson DH, Prado J, Silva PC, Skog JE, Wiersema JH, Turland NJ. 2006. *International Code of Botanical Nomenclature (Vienna Code) adopted by the Seventeenth International Botanical Congress, Vienna, Austria, July 2005*. *Regnum Vegetabile* 146. 568 p.
- Mangin L. 1903. Sur la maladie du châtaignier causée par le *Mycelophagus castaneae*. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences* 136: 470–473.
- Miyata Y, Odagiri Y. 1982a. [A culture method for the mass production of *Phytophthora castaneae* oospores.] *Scientific Reports of the Kyoto Prefectural University, Agriculture* 34: 28–34. [Abstract in: *Review of Plant Pathology* 62: 380. 1983.]
- Miyata Y, Odagiri Y. 1982b. [Fine structural comparison of *Phytophthora castaneae* oogonia and oospores with or without sterol.] *Scientific Reports of the Kyoto Prefectural University, Agriculture* 34: 35–45. [Abstract in: *Review of Plant Pathology* 62: 380. 1983.]
- Newhook FJ, Waterhouse GM, Stamps DJ. 1978. *Tabular key to the species of Phytophthora de Bary*. *Mycological Papers* 143. 20 p.
- Ooka JJ, Uchida JY. 1984. *Phytophthora katsurae* fruit rot of coconut in Hawaii. *Phytopathology* 74: 816.

- Oudemans P, Coffey MD. 1991. A revised systematics of twelve papillate *Phytophthora* species based on isozyme analysis. *Mycological Research* 95: 1025–1046.  
[http://dx.doi.org/10.1016/S0953-7562\(09\)80543-1](http://dx.doi.org/10.1016/S0953-7562(09)80543-1)
- Ristaino JB. 2011. Fact sheet – *Phytophthora katsurae*. Key for Identification of Common Phytophthora Species. St. Paul MN, APS Press. CD.
- Schell E. 1922. Diseases of the French chestnut tree—particularly the ‘ink-malady’. *Journal of the American Leather Chemists Association* 17: 353–358. [Abstract in: *Review of Applied Mycology* 3: 5–7. 1924.]
- Stamps DJ. 1985. *Phytophthora katsurae*. C.M.I. Descriptions of Pathogenic Fungi and Bacteria No. 837. 2 p.
- Stamps DJ, Waterhouse GM, Newhook FJ, Hall GS. 1990. Revised tabular key to the species of *Phytophthora*. Second edition. *Mycological Papers* 162. 28 p.
- Uchida K. 1976. [Studies on the trunk rot of Japanese chestnut trees caused by *Phytophthora castaneae* Katsura & Uchida.] *Bulletin of the Ibaraki-ken Horticultural Experiment Station, Special Issue No. 3*. 48 p. [Abstract in: *Review of Plant Pathology* 58: 247–248. 1979.]
- Uchida JY, Aragaki M, Ooka JJ, Nagata NM. 1992. *Phytophthora* fruit and heart rots of coconut in Hawaii. *Plant Disease* 76: 925–927. <http://dx.doi.org/10.1094/PD-76-0925>
- Vannini A, Vettraino AM. 2001. Ink disease in chestnuts: impact on the European chestnut. *Forest Snow and Landscape Research* 76: 345–350.
- Vettraino AM, Morel O, Perlerou C, Robin C, Diamandis S, Vannini A. 2005. Occurrence and distribution of *Phytophthora* species in European chestnut stands, and their association with ink disease and crown decline. *European Journal of Plant Pathology* 111: 169–180.  
<http://dx.doi.org/10.1007/s10658-004-1882-0>
- Waterhouse GM. 1963. Key to the species of *Phytophthora* de Bary. *Mycological Papers* 92. 22 p.
- Waterhouse GM. 1970. The genus *Phytophthora* de Bary. Second edition. *Mycological Papers* 122. 59 p.