

## MYCOTAXON

<http://dx.doi.org/10.5248/121.375>

Volume 121, pp. 375–383

July–September 2012

**Three new species of *Septobasidium* (*Septobasidiaceae*) from southern and southwestern China**SUZHEN CHEN<sup>1,2</sup> & LIN GUO<sup>1\*</sup><sup>1</sup>*State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China*<sup>2</sup>*Ocean University of China, Qingdao 266003, China*\* CORRESPONDENCE TO: [guol@im.ac.cn](mailto:guol@im.ac.cn)

ABSTRACT — Three new species, *Septobasidium symploci* on *Symplocos* sp. associated with *Aulacaspis* sp., *Septobasidium euonymi* on *Euonymus japonicus* associated with *Aonidiella* sp., and *Septobasidium cotoneastri* on *Cotoneaster rubens* associated with *Aulacaspis* sp., are described. They were collected from Hainan and Yunnan Provinces and Xizang Autonomous Region, China.

KEY WORDS — *Pucciniomycetes*, *Septobasidiales*, taxonomy

Southern and southwestern China has a varied and rich mycota. Several mycological investigations dealing with many new species and new Chinese records from these areas have been published recently (Dai et al. 2011, Zhang et al. 2012). An additional three new species of *Septobasidium* are reported as follows:

***Septobasidium symploci* S.Z. Chen & L. Guo, sp. nov.**

FIGS 1–7

FUNGAL NAME FN 570030

Differs from *Septobasidium thwaitesii* by its smaller basidiomata and basidia.TYPE: China, Hainan Province, Bawangling Natural Reserve, alt. 950 m, on *Symplocos* sp. (*Symplocaceae*), associated with *Aulacaspis* sp. (*Diaspididae*), 13.IV.2011, L. Guo 11603, (HMAS 242888, **holotype**).

ETYMOLOGY: The epithet refers to the substrate plant genus.

Basidiomata on trunks and branches, punctiform, 0.1–2 cm long, 0.1–0.3 cm wide, grey-brown or cinnamon-brown, forming small isolated or confluent patches, irregular growth; margin indeterminate, surface smooth. In section 500–1050 µm thick. Subiculum brownish, 30–50 µm thick. Pillars brown,




FIG. 1. *Septobasidium symploci* (HMAS 242888, holotype). Probasidia and basidia.

50–80  $\mu\text{m}$  high, 40–50  $\mu\text{m}$  wide. Hyphae often forming 3–4 inconspicuous strata, 400–900  $\mu\text{m}$  high, brown. Hymenium hyaline or brown, 110–135  $\mu\text{m}$  thick. Probasidia subglobose, obovoid or ellipsoidal, 12–18  $\times$  7–14  $\mu\text{m}$ , hyaline or brown. Basidia cylindrical, curved, 4-celled, 32–42  $\times$  8–9  $\mu\text{m}$ , brown, with persisting probasidial cell. Sterigmata conical, 5–8  $\times$  2–3  $\mu\text{m}$ , hyaline. Basidiospores ovoid, 10–20  $\times$  5–6  $\mu\text{m}$ . Haustoria consisting of hyphae.

COMMENTS: *Septobasidium symploci* is similar to *S. thwaitesii* (Berk. & Broome) Pat., but the latter has larger basidiomata, 15–20 cm long, with larger fissions on the surface, slightly larger and wider basidia, 37–46  $\times$  12–13.5  $\mu\text{m}$ , and two conspicuous horizontal layers in the section (Couch 1938).

***Septobasidium cotoneastri* S.Z. Chen & L. Guo, sp. nov.**

Figs 8–14

FUNGAL NAME FN 570031






Differs from *Septobasidium patouillardii* by its thicker section and smaller basidia.

TYPE: China, Xizang, Nyingchi, Gadinggou, alt. 2980 m, on *Cotoneaster rubens* W.W. Sm. (*Rosaceae*), associated with *Aulacaspis* sp., 25.IX.2010, S.H. He XZ11 (HMAS 251318, holotype).

ETYMOLOGY: The epithet refers to the substrate plant genus.

---

Figs. 2–7 *Septobasidium symploci* (HMAS 242888, holotype). 2. Basidiomata on trunk. 3–4. Sections of basidiomata. 5–6. Probasidia and basidia (arrows). 7. Haustoria.



8








FIG. 8. *Septobasidium cotoneastri* (HMAS 251318, holotype). Basidia.

Basidiomata on branches, 0.2–10 cm long, 0.1–4.5 cm wide, brown or chestnut-brown, forming small isolated or larger patches, irregular growth, sometimes growing along the subiculum, up to 5000  $\mu\text{m}$  high, emerging chestnut-brown hyphal layers; margin determinate, surface smooth, cracked at maturity. In section 600–1750(–5000)  $\mu\text{m}$  thick. Subiculum brown, 30–50  $\mu\text{m}$  thick. Pillars brown, 150–200  $\mu\text{m}$  high, 30–70  $\mu\text{m}$  wide. Hyphal layer 200–1100(–4500)  $\mu\text{m}$  high, brown. Hymenium hyaline or brown, 80–120  $\mu\text{m}$  thick, with vertical thin-walled, ca. 0.5  $\mu\text{m}$  thick paraphyses. Basidia arising directly from the hyphae without a probasidial cell, clavate, straight, 2-celled, 15–19  $\times$  4–6.5  $\mu\text{m}$ , hyaline or brown; wall 0.5  $\mu\text{m}$ . Basidiospores not seen. Haustoria consisting of regularly coiled hyphae.

ADDITIONAL SPECIMENS EXAMINED: CHINA, XIZANG, NYINGCHI, Gadinggou, alt. 2980 m, on *Cotoneaster obscurus* Rehder & E.H. Wilson (*Rosaceae*), 25.IX.2010, S.H. He XZ14, (HMAS 251320); on *Sorbus rufopilosa* C.K. Schneid. (*Rosaceae*), 25.IX.2010, S.H. He XZ13, (HMAS 251319); 25.IX.2010, S.H. He XZ19, (HMAS 251328).

Figs. 9–14. *Septobasidium cotoneastri*. 9. Basidiomata on branch (HMAS 251318, holotype). 10. Basidiomata on branches (HMAS 251319, paratype). 11. Section of basidioma (HMAS 251318, holotype). 12–13. Basidia (arrows) (HMAS 251318, holotype). 14. Haustoria (HMAS 251318, holotype).





COMMENTS: *Septobasidium cotoneastri* is similar to *S. patouillardii* Burt, but the latter has a thinner section (300–460  $\mu\text{m}$  high), larger basidia (18–23  $\times$  5.2–5.6  $\mu\text{m}$ ) and thick-walled (ca. 3.2  $\mu\text{m}$  thick) paraphyses in the hymenium (Couch 1938).

*Septobasidium euonymi* S.Z. Chen & L. Guo, sp. nov.

FIGS 15–21

FUNGAL NAME FN 570032

Differs from *Septobasidium patouillardii* by its thinner section and transverse, thin-walled threads in the hymenium.

TYPE: China, Yunnan Province, Lijiang, Yushuizhai, alt. 2200 m, on *Euonymus japonicus* Thunb. (Celastraceae), associated with *Aonidiella* sp. (Diaspididae), 25.XI.2011, S.H. He YN01 (HMAS 251324, holotype).

ETYMOLOGY: The epithet refers to the substrate plant genus.

Basidiomata on branches, resupinate, 2–8 cm long, 1.5–4 cm wide, cinnamon-brown, brown or chestnut-brown; margin white, determinate, surface smooth, cracked at maturity. In section 180–320  $\mu\text{m}$  thick. Subiculum brown, 20–40  $\mu\text{m}$  thick. Pillars brown, 60–110  $\mu\text{m}$  high, 20–130(–300)  $\mu\text{m}$  wide. Hyphal layer 20–80  $\mu\text{m}$  high, brown. Hymenium hyaline or brown, 50–60  $\mu\text{m}$  thick,








FIG. 15. *Septobasidium euonymi* (HMAS 251324, holotype). Basidia.

FIGS. 16–21. *Septobasidium euonymi* (HMAS 251324, holotype). 16. Basidiomata on branch. 17–18. Sections of basidiomata. 19–20. Basidia (arrows). 21. Haustoria.



with transverse, thin-walled threads. Basidia arising directly from the hyphae without a probasidial cell, clavate, straight, 2-celled,  $17\text{--}25 \times 6.5\text{--}8 \mu\text{m}$ , hyaline or brownish; wall ca.  $1 \mu\text{m}$ . Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae and globose cells.

COMMENTS: *Septobasidium euonymi* is similar to *S. patouillardii*, but the latter has a thicker section ( $300\text{--}460 \mu\text{m}$  high) and vertical, thick-walled (ca.  $3.2 \mu\text{m}$  thick) paraphyses in the hymenium (Couch 1938).

Including the three new species reported in this paper, 49 *Septobasidium* species have now been reported in China (Sawada 1933, Couch 1938, Teng 1963, Tai 1979, Kirschner & Chen 2007, Lu & Guo 2009a,b,c, 2010a,b,c, 2011, Lu et al. 2010, Chen & Guo 2011a,b,c,d, 2012a,b).

#### Acknowledgements

The authors would like to express their deep thanks to Dr. Eric H.C. McKenzie (Auckland, New Zealand) for serving as pre-submission reviewer, to Dr. Shuanghui He (Beijing Forestry University) for serving as pre-submission reviewer and collecting specimens, to Dr. Shaun Pennycook (Auckland, New Zealand) for nomenclatural review, to Prof. Zhenyu Li and Mr. Ziyu Cao (Institute of Botany, Chinese Academy of Sciences) for identifying the host plants, to Prof. Sanan Wu (Beijing Forestry University) for identifying the scale insects, and to Mrs. Xiangfei Zhu for inking in line drawings. This study was supported by the foundation of Ministry of Science and Technology of the People's Republic of China (No. 2006FY110500-5).

#### Literature cited

- Chen SZ, Guo L. 2011a. *Septobasidium sichuanense* sp. nov. (*Septobasidiaceae*) from China. Mycotaxon 115: 481–484. <http://dx.doi.org/10.5248/115.481>
- Chen SZ, Guo L. 2011b. *Septobasidium atalantiae* sp. nov. (*Septobasidiaceae*) and *S. henningsii* new to China. Mycotaxon 117: 291–296. <http://dx.doi.org/10.5248/117.291>
- Chen SZ, Guo L. 2011c. *Septobasidium saurauiae* sp. nov. (*Septobasidiaceae*) and *S. pseudopedicellatum* new to China. Mycotaxon 118: 283–288. <http://dx.doi.org/10.5248/118.283>
- Chen SZ, Guo L. 2011d. *Septobasidium glycosmidis* and *S. albiziae* spp. nov. (*Septobasidiaceae*) from Hainan Province. Mycosystema 30: 861–864.
- Chen SZ, Guo L. 2012a. Three new species and three new Chinese records of *Septobasidium* (*Septobasidiaceae*). Mycosystema 31: 651–655.
- Chen SZ, Guo L. 2012b. Three new species of *Septobasidium* (*Septobasidiaceae*) from Hainan Province in China. Mycotaxon 120: 269–276. <http://dx.doi.org/10.5248/120.269>
- Couch JN. 1938. The genus *Septobasidium*. Univ. of North Carolina Press, Chapel Hill. 480 p.
- Dai YC, Cui BK, Yuan HS, He SH, Wei YL, Qin WM, Zhou LW, Li HJ. 2011. Wood-inhabiting fungi in southern China 4. Polypores from Hainan Province. Annales Bot. Fennici 48: 219–231.
- Kirschner R, Chen CJ. 2007. New reports of two hypophyllous *Septobasidium* species from Taiwan. Fung. Sci. 22(1,2): 39–46.
- Lu CX, Guo L. 2009a. *Septobasidium maesae* sp. nov. (*Septobasidiaceae*) from China. Mycotaxon 109: 103–106. <http://dx.doi.org/10.5248/109.103>



- Lu CX, Guo L. 2009b. Two new species of *Septobasidium* (*Septobasidiaceae*) from China. *Mycotaxon* 109: 477–482. <http://dx.doi.org/10.5248/109.477>
- Lu CX, Guo L. 2009c. *Septobasidium annulatum* sp. nov. (*Septobasidiaceae*) and *Septobasidium kameii* new to China. *Mycotaxon* 110: 239–245. <http://dx.doi.org/10.5248/110.239>
- Lu CX, Guo L. 2010a. Three new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. *Mycotaxon* 112: 143–151. <http://dx.doi.org/10.5248/112.143>
- Lu CX, Guo L. 2010b. Two new species of *Septobasidium* (*Septobasidiaceae*) and *S. pallidum* new to China. *Mycotaxon* 113: 87–93. <http://dx.doi.org/10.5248/113.87>
- Lu CX, Guo L. 2010c. Two new species of *Septobasidium* (*Septobasidiaceae*) from Hainan province in China. *Mycotaxon* 114: 217–223. <http://dx.doi.org/10.5248/114.217>
- Lu CX, Guo L. 2011. Two new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. *Mycotaxon* 116: 395–400. <http://dx.doi.org/10.5248/116.395>
- Lu CX, Guo L, Wei JG, Li JB. 2010. Two new species of *Septobasidium* (*Septobasidiaceae*) from southern China. *Mycotaxon* 111: 269–274. <http://dx.doi.org/10.5248/111.269>
- Sawada K. 1933. Descriptive catalogue of the Formosan fungi. Part VI. Rep. Dept. Agric. Govt. Res. Inst. Formosa 61: 1–99.
- Tai FL. 1979. Sylloge Fungorum Sinicorum. Science Press, Beijing. 1527 p.
- Teng SC. 1963. Fungi of China. Science Press, Beijing. 808 p.
- Zhang YD, Ma J, Ma LG, Zhang XG. 2012. Two new species of *Taeniolina* from southern China. *Mycol. Progr.* 11: 71–74. <http://dx.doi.org/10.1007/s11557-010-0729-7>