
MYCOTAXON

<http://dx.doi.org/10.5248/121.413>

Volume 121, pp. 413–417

July–September 2012

***Heteroconium bannaense* sp. nov. and a new record of the genus from China**

JI-WEN XIA, SHOU-CAI REN, LI-GUO MA & XIU-GUO ZHANG*

Department of Plant Pathology, Shandong Agricultural University, Taian, 271018, China

*CORRESPONDENCE TO: zhxg@sdau.edu.cn, sdau613@163.com

ABSTRACT — Two species of the anamorphic genus *Heteroconium* were discovered from tropical forests in southern China. *Heteroconium bannaense* sp. nov. is described and illustrated based on a specimen collected on dead stems of *Phragmites communis*. *Heteroconium arundicum* is newly recorded from China.

KEY WORDS — conidial fungi, taxonomy

Introduction

Heteroconium was established by Petrak (1949) with *H. citharexyli* Petr. as the type species. The genus is characterized by distinct, single conidiophores with monoblastic, terminal, determinate or percurrently extending conidiogenous cells that produce catenate, euseptate conidia. The conidiophores are unbranched or with a secondary branch originating after conidial secession or near a percurrent proliferation (Petrak 1949, Castañeda et al. 1999, 2008, Taylor et al. 2001). The criteria used for species delimitation in *Heteroconium* are primarily based on conidial morphology and size. Castañeda et al. (2008) compared *Heteroconium* with several other similar genera, including *Cladophialophora* Borelli, *Lylea* Morgan-Jones, *Phaeoblastophora* Partr. & Morgan-Jones, *Septonema* Corda, *Taeniolella* S. Hughes, and *Xenoheteroconium* Bhat et al. To date, 24 legitimate names have been included under *Heteroconium* (Mycobank 2012, Ma et al. 2012a,b, Ren et al. 2012). However, *Heteroconium chaetospira* (Grove) M.B. Ellis, *H. solaninum* (Sacc. & P. Syd.) M.B. Ellis, and *H. tetracoilum* (Corda) M.B. Ellis have been transferred to other hyphomycete genera (Holubová-Jechová 1978, Crous et al. 2007, Hughes 2007), and *H. queenslandicum* Matsush. is considered probably to belong in the genus *Parapleurotheciopsis* P.M. Kirk (Castañeda et al. 2008). Thus, *Heteroconium* currently has 20 accepted taxa. Six species, *H. annesleae* S.C. Ren & X.G. Zhang, *H. fici* L.G. Ma & X.G. Zhang, *H. neolitseae* S.C. Ren & X.G. Zhang,




FIG. 1. *Heteroconium bannaense*.

A. Colonies on natural substratum. B. Conidiophores with conidia. C, D. Conidia.

H. phellodendri J.Ma & X.G. Zhang, *H. schimae* Y.D. Zhang & X.G. Zhang, and *H. tsoongiodendri* L.G. Ma & X.G. Zhang have been previously described from China (Zhang et al. 2010, Ma et al. 2012a,b, Ren et al. 2012).

During our ongoing survey of anamorphic fungi associated with woody debris in tropical forests of southern China, two species with morphological characteristics of genus *Heteroconium* were collected on decaying twigs and dead stems. One species is proposed herein as a new species, while the other is a new record for China.

***Heteroconium bannaense* J.W. Xia & X.G. Zhang, sp. nov.**

FIG. 1

MYCOBANK MB 801029

Differs from all other *Heteroconium* species in its typical obclavate conidia with both eusepta and distosepta.

TYPE: China, Yunnan Province: Xishuangbanna, on dead stems of *Phragmites communis* Trin. (*Poaceae*), 2 Nov 2011, J.W. Xia (**Holotype**, HSAUP H6035; **isotype**, HMAS 243411).

ETYMOLOGY: in reference to the type locality.

Colonies on natural substrate effuse, dark brown, hairy. Mycelium partly superficial, partly immersed in the substratum, composed of septate, pale brown, smooth, 1–2 μm wide hyphae. Conidiophores macronematous, mononematous, unbranched, erect, straight or slightly flexuous, cylindrical, smooth, thick-walled, brown, 6–11-septate, $76.5\text{--}197 \times 3\text{--}7 \mu\text{m}$. Conidiogenous cells monoblastic, integrated, terminal, cylindrical, determinate or percurrent, brown, smooth, truncate at the apex, $6.5\text{--}8 \times 2.5\text{--}3.5 \mu\text{m}$. Conidial secession schizolytic. Conidia holoblastic, acrogenous, blastocatenate, in chains of up to 9, straight or slightly curved, obclavate, brown, smooth, 14–22-septate (6–8-euseptate, 8–14-distoseptate), $58\text{--}93.5 \mu\text{m}$ long, $5\text{--}8 \mu\text{m}$ wide, truncate at the base.

COMMENTS – Among the known species, *Heteroconium bannaense* bears some resemblance in conidial shape to *H. tropicale* R.F. Castañeda & W.B. Kendrick (Castañeda & Kendrick 1990). However, the conidia of *H. bannaense* are larger than those of *H. tropicale* (conidia $26\text{--}52 \times 4\text{--}6 \mu\text{m}$, 3–6-euseptate), and have more septa. In addition, *H. bannaense* is unique within the genus in its typical obclavate conidia with both eusepta and distosepta.

***Heteroconium arundicum* Chowdhry, Indian Phytopath. 33: 361, 1981 [“1980”].**

FIG. 2

Colonies on natural substrate effuse, brown to blackish brown, hairy. Mycelium mostly immersed in the substratum. Conidiophores macronematous, mononematous, solitary or in groups, sometimes branched, erect, straight to flexuous, cylindrical, septate, brown, smooth, up to $180 \times 5.5\text{--}7.5 \mu\text{m}$. Conidial secession schizolytic. Conidia holoblastic, acrogenous, blastocatenate, straight or slightly curved, cylindrical to fusiform, brown, smooth, 3–8-euseptate, $56\text{--}110 \mu\text{m}$ long, $7.5\text{--}11 \mu\text{m}$ wide, truncate at base.




FIG. 2. *Heteroconium arundicum*.
Conidiophores and conidia.

SPECIMEN EXAMINED: CHINA, HAINAN PROVINCE: Lingao, on decaying twigs of unidentified broad-leaved tree, 10 April 2011, J. Ma (HSAUP H5493-2, HMAS 243412).

COMMENTS – *Heteroconium arundicum* is reported for the first time from China. Compared with the type material described by Chowdhry (1981), the conidiophores of Chinese specimen are occasionally branched, but other features of this taxon closely match those of the original description. *Heteroconium arundicum* is similar to *H. citharexylis* (Petraek 1949) in conidial shape, but differs in having larger conidia ($56\text{--}110 \times 7.5\text{--}11 \mu\text{m}$ vs. $10\text{--}40 \times 3\text{--}7 \mu\text{m}$).

Acknowledgments

The authors express gratitude to Dr. Eric H.C. McKenzie and Dr. Bryce Kendrick for serving as pre-submission reviewers and for their valuable comments and suggestions. This project was supported by the National Natural Science Foundation of China (Nos. 31093440, 30499340, 30770015) and the Ministry of Science and Technology of the People's Republic of China (Nos. 2006FY120100, 2006FY110500–5).

Literature cited

- Castañeda Ruíz RF, Kendrick B. 1990. Conidial fungi from Cuba: II. Univ. Waterloo Biol. Ser. 33: 1–61.
- Castañeda Ruíz RF, Saikawa M, Guarro J. 1999. A new species of *Heteroconium* from a tropical rainforest. Mycotaxon 71: 295–300.
- Castañeda Ruíz RF, Iturriaga T, Heredia Abarca G, Minter DW, Gené J, Stadler M, Saikawa M, Silvera-Simón C. 2008. Notes on *Heteroconium* and a new species from Venezuela. Mycotaxon 105: 175–184.
- Chowdhry PN. 1981 [“1980”]. A new species of *Heteroconium* from India. Indian Phytopath. 33: 361–362.
- Crous PW, Mohammed C, Glen M, Verkley GJM, Groenewald JZ. 2007. Eucalyptus microfungi known from culture. 3. *Eucasphaeria* and *Sympoventuria* genera nova, and new species of *Furcaspora*, *Harknessia*, *Heteroconium* and *Phacidiella*. Fungal Divers. 25: 19–36.
- Holubová-Jechová V. 1978. Lignicolous hyphomycetes from Czechoslovakia 5. *Septonema*, *Hormiactella*, and *Lylea*. Folia Geobot. Phytotax. 13: 421–442.
- Hughes SJ. 2007. *Heteroconium* and *Pirozynskiella* n. gen., with comments on conidium transeptation. Mycologia 99: 628–638. <http://dx.doi.org/10.3852/mycologia.99.4.628>
- Ma LG, Ma J, Zhang YD, Castañeda Ruíz RF, Zhang XG. 2012a. New species and records of *Heteroconium* (anamorphic fungi) from southern China. Mycoscience. <http://dx.doi.org/10.1007/s10267-012-0190-3>
- Ma J, Ma LG, Zhang YD, Castañeda Ruíz RF, Zhang XG. 2012b. New species or records of *Endophragmiella* and *Heteroconium* from southern China. Cryptog. Mycol. 33: 127–135.
- Mycobank. 2012. MycoSearch database search. International Mycological Association, [Accessed: 24 July 2012] <http://www.mycobank.org/Biolomics.aspx?Table=Mycobank&Page=200&ViewMode=Basic>.
- Petrak F. 1949. Neue Hyphomyzeten-Gattungen aus Ekuador. Sydowia 3: 259–266.
- Ren SC, Ma J, Zhang XG. 2012. Two new *Heteroconium* species and two other forest microfungi newly recorded from China. Mycotaxon 119: 361–367. <http://dx.doi.org/10.5248/119.361>
- Taylor JE, Crous PW, Palm ME. 2001. Foliar and stem fungal pathogens of *Proteaceae* in Hawaii. Mycotaxon 78: 449–490.
- Zhang YD, Ma J, Ma LG, Zhang XG. 2010. A new species of *Heteroconium* from Fujian, China. Mycotaxon 114: 315–318. <http://dx.doi.org/10.5248/114.315>