
MYCOTAXON

<http://dx.doi.org/10.5248/122.157>

Volume 122, pp. 157–160

October–December 2012

Myxomycetes from China 16: *Arcyodes incarnata* and *Licea retiformis*, newly recorded for China

BO ZHANG¹ & YU LI^{1*}

¹Engineering Research Center of Chinese Ministry of Education for Edible and Medicinal Fungi,
Jilin Agricultural University, 2888 Xincheng Street, Changchun City, P. R. China

* CORRESPONDENCE TO: yuli966@126.com

ABSTRACT — Two new records for China, *Arcyodes incarnata* and *Licea retiformis*, are reported in this paper. Materials collected from Yunnan and Fujian province are described and illustrated by scanning electron micrographs. Those specimens are deposited in the Herbarium of Mycological Institute of Jilin Agricultural University (HMJAU), Changchun, China.

KEY WORDS — taxonomy, slime molds, SEM

Introduction

Our examination of previous myxomycete collections has revealed two new records for China. One is in *Arcyodes*, which is currently accepted as a monotypic genus. The other is in *Licea*, which now comprises about 71 species (Kirk et al. 2008, Lado 2001, 2012–14), of which 13 are reported for mainland China (Chou 1937; Li & Li 1989, 1994; Li 2005) and 11 for Taiwan (Liu et al. 2002).

Materials & methods

Fruiting bodies and microscopic structures were examined using light and scanning electron microscopes (Martin & Alexopoulos 1969; Li et al. 1993, Li & Li 1995). Permanent slides were mounted in Hoyer's reagent (Martin & Alexopoulos 1969). Colored slides were prepared according to Robbrecht (1974) by spreading the capillitium in a drop of 94% alcohol, determining color after one minute, and then mounting in Hoyer's reagent. Color terms are given according to the Flora of British Fungi (1969). The specimens are deposited in the Herbarium of the Mycological Institute, Jilin Agricultural University (HMJAU).

Taxonomy




PLATE 1. *Arcyodes incarnata*: A, fruiting bodies. B, inner surface of peridium. C, D, part of the capillitium and spores. E, spore. Scale bars: A = 1 mm; B, E = 2 μ m; C = 5 μ m; D = 10 μ m.

Arcyodes incarnata (Alb. & Schwein.) O.F. Cook, Science, N.Y. 15: 651, 1902. PL. 1

SPOROCARPS aggregated, sessile, crowded and heaped, 0.5–0.7 mm in diam., fawn, fading to dull ochraceous. HYPOTHALLUS inconspicuous. PERIDIUM single, membranous, somewhat opalescent, persistent, irregularly dehiscent above, yellow to colourless by transmitted light, inner surface with many prominent warts, low irregular ridges, sometime forming a net with small protuberances. COLUMELLA absent. CAPILLITIUM tubular, elastic, branched and anastomosed, pale yellow by transmitted light, mostly 4 μ m in diam., with numerous inflations, decorated with many cogs and irregular edges, sometimes with a faint reticulation. SPORES free, pale pink in mass, yellowish pale to colourless by transmitted light, 7–9.5 μ m in diam., globose to subglobose, densely warted, with scattered groups of more prominent warts.

SPECIMEN EXAMINED: CHINA, Yunnan Province, Puer, on the bark surface of a dead log, 28 August 2011, Zhang Bo1101 (HMJAU10301).

COMMENTS: *Arcyodes incarnata*, which has been recorded from North America, Europe, and Africa (Martin & Alexopoulos 1969, Lado 1994), is apparently rare in Asia. The Yunnan specimen has slightly larger spores than the type specimen

(6–8 μm in diam.). However, the Yunnan and type specimen are similar in habitat on dead logs, a pale pink spore mass, faintly reticulate capillitium, and pale copper-colored sporocarps.




PLATE 2. *Licea retiformis*: A, B, fruiting bodies. C, part of peridium. D, spore. Scale bar: A = 1 mm.

Licea retiformis Nawawi, Trans. Br. mycol. Soc. 60(1): 153, 1973.

PL. 2

FRUCTIFICATIONS plasmodiocarps, gregarious, reticulation 0.6–3 mm in extension, pale yellowish brown, fading to pale brown. PLASMODIOCARPS 90–100 μm in diameter, appearing smooth on the surface. HYPOTHALLUS inconspicuous. PERIDIUM single, membranous, firm, translucent. DEHISCENCE by longitudinal slit, decorated with loose small mesh. COLUMELLA absent. CAPILLITIUM absent. SPORES free, pale ochraceous in mass, yellowish pale to colourless by transmitted light, subglobose to ovate, 9–11 \times 8.5–12.5 μm , densely warted.

SPECIMENS EXAMINED: CHINA, Fujian Province, Fuzhou, on the bark surface of a dead log, 9 July 2010, Li Ming0031 (HMJAU10302); Li Ming0042 (HMJAU10303).

COMMENTS: This is the only *Licea* species with reticulate plasmodiocarps. *Licea retiformis*, which has been reported from Japan (Yamamoto 1998), Malaysia (Nawawi 1973), and Taiwan (Liu et al. 2002), is here newly recorded from mainland China. The two Fujian specimens have slightly shorter spores than the type specimen (9.3–13.5 \times 8.5–12.5 μm , Nawawi 1973).

Acknowledgments

We thank Professors. Guozhong Lü and A.J.S. Whalley for their valuable revisions and kind help. This study was supported by funds from the Ministry of Agriculture of China.

Literature cited

- Chou ZH. 1937. Notes on myxomycetes from China. Bull. Fan. Men. Inst. Biol. 7: 257–278.
- Cook OF. 1902. Types and synonyms. Science 15: 646–656.
<http://dx.doi.org/10.1126/science.15.382.646>
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. Ainsworth & Bisby's dictionary of the fungi. 10th ed. CAB International, Wallingford. 771 p.
- Lado C. 1994. A checklist of myxomycetes of the Mediterranean countries. Mycotaxon 52(1): 117–185.
- Lado C. 2001. Nomenmyx: a nomenclatural taxa base of myxomycetes. Cuadernos de Trabajo de Flora Micológica Ibérica 16. 224 p.
- Lado C. 2012–14. An online nomenclatural information system of *Eumycetozoa*.
<http://www.nomen.eumycetozoa.com>
- Li Y. 2005. Liceales in China. Chinese Science Press. 126 p.
- Li Y, Li HZ. 1989. Myxomycetes from China I: A checklist of myxomycetes from China. Mycotaxon 35(2): 429–436.
- Li Y, Li HZ. 1994. Myxomycetes from China XII: A new species of *Licea*. Mycosystema 7: 133–135.
- Li Y, Li HZ. 1995. Myxomycetes from China III: Description of a new species, *Cribraria media*, and discussion of the relationship between *Cribraria* and *Dictydium*. Mycotaxon 53: 69–80.
- Li Y, Chen SL, Li HZ. 1993. Myxomycetes from China X: Additions and notes to *Trichiaceae* from China. Mycosystema 6: 107–112.
- Liu CH, Chen YF, Chang JH, Yang FH. 2002. Myxomycetes of Taiwan. Three new records. Taiwania 47(3): 179–185.
- Martin GM, Alexopoulos CJ. 1969. The myxomycetes. University of Iowa Press. Iowa. 561 p.
<http://dx.doi.org/10.2307/1218569>
- Nawawi A. 1973. New species of *Licea* from Malaysia. Tans. Br. Myc. Soc. 60: 153–155.
[http://dx.doi.org/10.1016/S0007-1536\(73\)80072-5](http://dx.doi.org/10.1016/S0007-1536(73)80072-5)
- Robbrecht E. 1974. The genus *Arcyria* Wiggers in Belgium. Bull. Jard. Bot. Nat. Belg. 44: 303–353.
<http://dx.doi.org/10.2307/3667676>
- Royal Botanic Garden, Edinburgh. 1969. Flora of British Fungi: colour identification chart. Edinburgh, H.M. Stationery Office. 6 p.
- Yamamoto Y. 1998. The Myxomycetes biota of Japan. Toyo Shorin Publishing Co., Ltd., Tokyo, Japan. 700 p.