

---

# MYCOTAXON

<http://dx.doi.org/10.5248/123.261>

Volume 123, pp. 261–264

January–March 2013

---

## ***Yunchangia*, a new genus of smut fungi (*Ustilaginaceae*) from China**

LIN GUO<sup>1\*</sup> & BIAO XU<sup>2</sup>

<sup>1</sup>State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences,  
Beijing 100101, China

<sup>2</sup>Tarim University, Xinjiang, Aral 843300, China

\* CORRESPONDENCE TO: [guol@im.ac.cn](mailto:guol@im.ac.cn)

**ABSTRACT** — *Yunchangia puccinelliae* gen. et sp. nov. on *Puccinellia glauca*, is described. The fungus was collected from Xinjiang Uygur Autonomous Region, China.

**KEY WORDS** — *Ustilaginomycetidae*, *Ustilaginales*, taxonomy

### **Introduction**

The Taklimakan desert located in the southern Xinjiang Uygur Autonomous Region is the second largest shifting sand desert in the world, covering 337,600 km<sup>2</sup>. During a mycological expedition to the margin of the western desert in 2010, a smut species on *Puccinellia glauca* was collected. The sorus of the smut causes a short, longitudinal leaf streak. It is covered by the epidermis and simultaneously contains fungal hyphae and single ustilospores. The fungal hyphae lie longitudinally along the sides of the sorus. Attempts to germinate ustilospores from *Puccinellia glauca* were carried out several times at 24°C on PDA (potato dextrose agar), but failed. To date 269 species of smut fungi have been reported in China (Guo 2000, 2011, He et al. 2011).

### **Material & methods**

Ustilospores and hyphae were mounted in lactophenol and examined by light microscopy (LM), after heating to boiling point. For scanning electron microscopy (SEM), dried ustilospores and hyphae were dusted onto double-sided adhesive tape, fixed on specimen stubs, sputter coated with gold, ca. 20 nm thick, and studied with a FEI Quanta 200 electron microscope.

## Taxonomy

### *Yunchangia* L. Guo & B. Xu, gen. nov.

FUNGAL NAME FN570049

Differs from *Ustilago* by its sorus simultaneously containing fungal hyphae and ustilospores.

TYPE SPECIES: *Yunchangia puccinelliae* L. Guo & B. Xu

ETYMOLOGY: The genus is dedicated to Professor Wang Yunchang, an outstanding Chinese mycologist on taxonomy of smut and rust fungi.

Sori on host plant in the *Poaceae* forming short streaks mainly on leaves, covered by the epidermis. Ustilospores single, pigmented. The fungal hyphae lie longitudinally along the sides of the sorus. Hyphae are composed of pigmented cells.

COMMENTS: *Yunchangia* is similar to *Ustilago* (Pers.) Roussel, which differs by absence of fungal hyphae in the sorus (Vánky 2002). Its characters demonstrate that the genus *Yunchangia* is a member of the family *Ustilaginaceae*.

### *Yunchangia puccinelliae* L. Guo & B. Xu, sp. nov.

FIGS 1–6

FUNGAL NAME FN 570050

Differs from *Ustilago striiformis* by its slightly larger ustilospores and with fungal hyphae in sorus.

TYPE: China, Xinjiang, Wuqia, Baykurt, alt. 2600 m, on *Puccinellia glauca* (Regel) V.I.Krecz. ex Drobow (*Poaceae*), 30.VIII.2010, Y.F. Zhu & L. Guo 231 (HMAS 243467, holotype; HMUT 1228, isotype).


ETYMOLOGY: The epithet refers to the host plant genus.

Sori in leaves as short streaks, sparsely in pedicels and inflorescences, 0.5–3.5 mm long, covered by the epidermis which later ruptures exposing powdery, dark brown spore mass. Ustilospores subglobose, ellipsoidal, ovoid or irregular, 9–19 × 7–12.5 µm, reddish brown; wall 0.5–0.8 µm thick, echinulate as seen by LM, finely verruculose between the spines as seen by SEM. Fungal hyphae are composed of cells; cells cylindrical, ovoid or irregular, 5–10.5 × 4–8 µm, reddish brown; wall 0.5–0.8 µm thick, smooth as seen by LM, rough or verruculose as seen by SEM.

COMMENTS: *Yunchangia puccinelliae* is similar to *Ustilago striiformis* (Westend.) Niessl, which differs by its slightly smaller ustilospores [9–15(–16) × 8–12(–13) µm] and the absence of fungal hyphae in the sorus (Vánky 1994).

---

FIGS. 1–6. *Yunchangia puccinelliae* (HMAS 243467, holotype). 1. Sori in leaves. 2–4. Ustilospores and fungal hyphae (LM). 5–6. Ustilospores and fungal hyphae (SEM).



### Acknowledgements

The authors would like to express their deep thanks to Dr. K. Vánky (Tübingen, Germany) for suggestion to describe the new genus, to Drs. Eric H.C. McKenzie (Auckland, New Zealand) and Cvetomir M. Denchev (Sofia, Bulgaria) for serving as pre-submission reviewers, to Dr. Shaun Pennycook (Auckland, New Zealand) for nomenclatural review, and to Dr. Wenli Chen (Institute of Botany, Chinese Academy of Sciences) for identifying the host plant. This study was supported by the National Natural Science Foundation of China (No. 30960019).

### Literature cited

- Guo L. 2000. Flora fungorum sinicorum vol. 12 *Ustilaginaceae*. Science Press, Beijing. 124p.
- Guo L. 2011. Flora fungorum sinicorum vol. 39 *Tilletiales Urocystales* etc. Science Press, Beijing. 124p.
- He SQ, Li HX, Chang XY, Wang SR. 2011. *Anthracoidea maquensis*, a new species of smut fungi from China. *Mycol Progr.* 10:53–55. <http://dx.doi.org/10.1007/s11557-010-0672-7>
- Vánky K. 1994. European smut fungi. Gustav Fischer Verlag Stuttgart, Jena, New York. 570p.
- Vánky K. 2002. Illustrated genera of smut fungi. APS Press, St. Paul, Minnesota. 238 p.