
MYCOTAXON

<http://dx.doi.org/10.5248/123.363>

Volume 123, pp. 363–367

January–March 2013

A new species of *Miriquidica* from China

XIN ZHAO, LU-LU ZHANG & ZUN-TIAN ZHAO*

College of Life Sciences, Shandong Normal University, Jinan, 250014, P. R. China

* CORRESPONDENCE TO: ztzhao@sohu.com

ABSTRACT — A new lichen species, *Miriquidica yunnanensis*, is described from China. Conspicuous features are its yellowish white to grayish white thallus, black-green epihymenium, colorless to light brown hypothecium, and a unique chemistry. A key to the seven similar *Miriquidica* species is also provided.

KEY WORDS — Yunnan, *Lecanorales*, saxicolous, taxonomy

Introduction

The generic concept of *Miriquidica* (*Lecanoraceae*) was first formulated by Hertel & Rambold (1987). It is characterized by lecideine apothecia, Lecanora-type asci, simple ascospores without a perispore, filiform conidia, and the occurrence of miriquidic acid in most species. This widely distributed genus presently includes 26 species worldwide and usually occurs on acidic rocks (Hertel & Rambold 1987; Rambold & Schwab 1990; Timdal 1993; Owe-Larsson & Rambold 2001; Hertel & Andreev 2003; Andreev 2004; Nash et al. 2004; Fryday 2008; Fryday & Coppins 2008; Coppins & Aptroot 2008; Lendemer & Knudsen 2008; Øvstedal et al. 2009). In China, only one species, *M. complanata*, has been reported (Wei 1991).

During our study of the lichen flora of Mt. Laojun in Yunnan province, China, a species of *Miriquidica* new to science was found. Here we present a brief diagnosis, an extended description, and a key to seven similar species in the genus.

Materials & methods

The specimen was collected from Yunnan, China, and is preserved in the Lichen Section of Botanical Herbarium, Shandong Normal University (SDNU). The morphological and anatomical characters of the specimen were examined using a stereo microscope (NIKON SMZ 745T) and a polarizing microscope (OLYMPUS CX41). Lichen substances in the specimen cited were identified using the standardized thin

layer chromatography techniques (TLC) with C system (Orange et al. 2010). The morphological and anatomical photos of the specie were taken under an OLYMPUS SZX16 stereomicroscope and an OLYMPUS BX61 compound microscope with DP72 respectively.

Taxonomic description

Miriquidica yunnanensis Lu L. Zhang & X. Zhao, sp. nov.

FIGS 1–2

MYCOBANK MB 802482

Differs from all other *Miriquidica* spp. by its yellowish white to grayish white thallus, blackish-green epihymenium, colorless to light brown hypothecium, small ascospores, and unique combination of chemical characters.

TYPE: China. Yunnan province, Mt. Laojun, alt. 4000 m, on rock, 7 Nov. 2009, H.Y. Wang 20121132 (Holotype, SDNU).

ETYMOLOGY: The specific epithet “yunnanensis” refers to Yunnan province, where this species was found.

THALLUS crustose, areolate, dispersed on a conspicuous black prothallus; AREOLES roundish, slightly convex to convex, small, 0.2–0.6(–0.8) mm; UPPER SURFACE yellowish white to grayish white, due to pigmented hyphae




FIGURE 1. *Miriquidica yunnanensis* (Holotype: SDNU (Wang 20121132)). A, B. Thallus; C. TLC of chemical components (At = atranorin; M = miriquidic acid; N = norstictic acid; U = unknown substance).




FIGURE 2. *Miriquidica yunnanensis* (Holotype: SDNU (Wang 20121132)). A. Section of apothecium; B. Paraphyses, after treatment with K; C. Ascus, after treatment with K/I.

of a lichenicolous fungus appearing dark gray, esorediate; MEDULLA I–, with some granules; PHOTOBIONT chlorococcoid; algal cells of 7–12.5(–15) µm in diam. APOTHECIA: sessile, 0.4–0.9 mm in diam.; DISC black, flat to convex, usually immarginate; EXCIPLE blackish green to black; EPIHYMENIUM blackish green; HYMENIUM hyaline, 60–70 µm; PARAPHYSES occasionally branched and anastomosing, 1.5–3 µm, apically 2–3.5 µm thick; HYPOTHECIUM hyaline to pale brown; ASCI of *Lecanora*-type s.l., 8-spored; ASCOSPORES hyaline, simple, ellipsoid to broadly ellipsoid, 8–11 × 5–6 µm. PYCNIDIA immersed, blackish; CONIDIA filiform, 10–15 µm long.

CHEMISTRY — Atranorin, miriquidic acid, norstictic acid, and unknown substances present.

DISTRIBUTION AND SUBSTRATE — Known only from Laojun Mountain, growing on siliceous rock at ca. 4000 m alt.

COMMENTS — Atranorin is not common in *Miriquidica*; only *M. atrofulva* (±) and *M. pycnocarpa* contain it in addition to *M. yunnanensis*. However, *M. atrofulva* is distinguished by a rusty colored thallus with dark blue-gray to blackish soralia and pale brown to olive-brown epihymenium and contains

stictic acid as a major compound; *M. pycnocarpa* differs in its I+ violet medulla, small apothecia (0.15–0.3 mm in diam.), and dark brown hypothecium and contains only atranorin.

Because *Miriquidica yunnanensis* is similar in morphology and anatomy to *M. disjecta*, *M. griseoatra*, *M. leucophaeoides*, *M. limitata*, and *M. paanaensis*, and *M. complanata* has been reported in China, we provide a key to these seven species.

Key to seven similar *Miriquidica* species

- 1a. Containing fumarprotocetraric acid. *M. limitata*
- 1b. Lacking fumarprotocetraric acid 2
- 2a. Containing atranorin *M. yunnanensis*
- 2b. Lacking atranorin 3
- 3a. Containing norstictic acid 4
- 3b. Lacking norstictic acid 5
- 4a. Thallus white, yellowish, or pale brown, glossy. *M. leucophaeoides*
- 4b. Thallus pale gray to gray, tinged yellowish *M. paanaensis*
- 5a. Thallus white to purely white. *M. disjecta*
- 5b. Thallus gray, gray-brown or brownish yellow to dark bluish gray, matt 6
- 6a. Apothecia 0.2–0.7 mm diam., immersed. *M. complanata*
- 6b. Apothecia 0.3–1.2 mm diam., sessile *M. griseoatra*

Acknowledgments

The project was financially supported by the National Natural Science Foundation of China (31070010, 31170187) and the Program for Scientific Research Innovation Team in Colleges and Universities of Shandong Province. The authors thank Dr. A. Aptroot (ABL Herbarium, Soest, The Netherlands), Prof. Shou-yu Guo (Chinese Academy of Sciences, Beijing, China) and Prof. G. Rambold (Dept. Of Mycology, University of Bayreuth, Germany) for presubmission reviews.

Literature cited

- Andreev MP. 2004. Notes on the lichen genus *Miriquidica* (*Lecanorales*, *Lecanoraceae*) in Russia. *Bibliotheca Lichenologica* 88: 15–42.
- Coppins BJ, Aptroot A. 2008. New species and combinations in the lichens of the British Isles. *Lichenologist* 40(5): 363–374. <http://dx.doi.org/10.1017/S0024282908008165>
- Fryday AM. 2006. New and interesting North American lichen records from the alpine and sub-alpine zones of Mt. Katahdin, Maine. *Bryologist* 109(4): 570–578. [http://dx.doi.org/10.1639/0007-2745\(2006\)109\[570:NAINAL\]2.0.CO;2](http://dx.doi.org/10.1639/0007-2745(2006)109[570:NAINAL]2.0.CO;2)
- Fryday AM. 2008. Three new species of lichenized fungi with cephalodia from the southern New Zealand shelf islands (Campbell Plateau). *Lichenologist* 40(4): 283–294. <http://dx.doi.org/10.1017/S002428290800769X>

- Fryday AM, Coppins BJ. 2008. *Ameliella*, a new genus of lichen-forming fungi from north-west Europe and western Canada. *Lichenologist* 40(5): 387–397.
<http://dx.doi.org/10.1017/S0024282908007913>
- Hertel H, Andreev MP. 2003. On some saxicolous lecideoid lichens of the Beringian Region and adjacent areas of Eastern Siberia and the Russian Far East. *Bryologist* 106(4): 539–551.
[http://dx.doi.org/10.1639/0007-2745\(2003\)106\[539:OSSLLO\]2.0.CO;2](http://dx.doi.org/10.1639/0007-2745(2003)106[539:OSSLLO]2.0.CO;2)
- Hertel H, Rambold G. 1987. *Miriquidica* genus novum Lecanoracearum (Ascomycetes lichenisati). *Mitt. Bot. Staatssamml. München* 23: 377–392.
- Lendemer JC, Knudsen K. 2008. Studies in lichens and lichenicolous fungi: further notes on North American taxa. *Mycotaxon* 103: 75–86.
- Nash TH, Ryan BD, Diederich P, Gries C, Bungartz F. 2004. Lichen flora of the Greater Sonoran Desert Region, vol. 2. 742 p.
- Orange A, James PW, White FJ. 2010. *Microchemical methods for the identification of lichens*. 2nd edition. London: British Lichen Society.
- Øvstedal DO, Tønsberg T, Elvebakk A. 2009. The lichen flora of Svalbard. *Sommerfeltia* 33: 1–393.
- Owe-Larsson B, Rambold G 2001. The sorediate species of the lichen genus *Miriquidica* (*Lecanorales*, *Lecanoraceae*). *Bibliotheca Lichenologica* 78: 335–364.
- Rambold G, Schwab AJ. 1990. Rusty coloured species of the lichen genus *Miriquidica* (*Lecanoraceae*). *Nord. J. Bot.* 10: 117–121. <http://dx.doi.org/10.1111/j.1756-1051.1990.tb01757.x>
- Timdal 1993. *Miriquidica ventosa* comb. nov., a rediscovered lichen. *Bryologist* 94(4): 616–618.
- Wei JC. 1991. An enumeration of lichens in China. International Academic Publishers, Beijing. 313 p.