

---

# MYCOTAXON

<http://dx.doi.org/10.5248/128.127>

Volume 128, pp. 127–130

April–June 2014

---

## A new species of *Arachnophora* from submerged wood in the Amazon rainforest, Brazil

JOSIANE SANTANA MONTEIRO<sup>1</sup>, LUIS FERNANDO PASCHOLATI GUSMÃO<sup>2\*</sup>,  
& RAFAEL F. CASTAÑEDA-RUIZ<sup>3</sup>

<sup>1</sup>*Departamento de Micologia, Universidade Federal de Pernambuco, 50670-420, Recife, Brazil*

<sup>2</sup>*Departamento de Ciências Biológicas, Laboratório de Micologia, Universidade Estadual de Feira de Santana, BR116 KM03, 44031-460, Feira de Santana, Brazil*

<sup>3</sup>*Instituto de Investigaciones Fundamentales en Agricultura Tropical 'Alejandro de Humboldt' (INIFAT), Académico Titular de la Academia de Ciencias de Cuba, Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200*

\*CORRESPONDENCE TO: [lgusmao@uefs.br](mailto:lgusmao@uefs.br)

**ABSTRACT** — *Arachnophora combuensis* sp. nov. is described and illustrated from specimens collected on submerged wood from Combu Island (Belém municipality, Pará, Brazil) in the Amazon rainforest. The fungus is characterized by simple pale brown conidiophores, monoblastic percurrently extending conidiogenous cells, and staurosporous conidia with a pale brown basal cell and 2–3 dark brown central cells that bear 1–3 conical pale brown arms. The new species is compared with morphologically similar taxa.

**KEY WORDS** — taxonomy, freshwater fungi, tropics

### Introduction

During a mycological survey of conidial fungi occurring on submerged decaying plant materials from the Amazon rainforest in Brazil, an interesting fungus was collected on submerged wood. Its conidiogenesis and conidial features clearly suggest placement within the genus *Arachnophora* Hennebert (Hennebert 1963). However, the fungus shows remarkable differences from all previously described *Arachnophora* species and therefore is described as new to science.

### Material & methods

Samples of submerged litter were placed in paper and plastic bags, taken to the laboratory, and treated according to Castañeda-Ruiz (2005). Mounts were prepared in PVL (polyvinyl alcohol, lactic acid, and phenol) and measurements were taken at  $\times 1000$ .

Micrographs were obtained with an Olympus microscope (model BX51) equipped with bright field and Nomarski interference optics. The type specimens are deposited in the Herbarium of Universidade Estadual de Feira de Santana (HUEFS).

### Taxonomy

*Arachnophora combuensis* J.S. Monteiro, R.F. Castañeda & Gusmão, sp. nov. FIG. 1

MYCOBANK MB 807635

Differs from *Arachnophora crassa* by its wider conidial central body and fewer, larger conidial arms and from *A. pulneyensis* by its wider conidial central body separated from the conidiogenous cell by a basal cell.

TYPE: Brazil, Pará State, Belém, Área de Proteção Ambiental Ilha do Combu, 1°25'S 48°27' W, on submerged wood, 10 Jan. 2013, coll. J.S. Monteiro (Holotype: HUEFS 196432).

ETYMOLOGY: Latin *combuensis* refers to the island where the type specimen was collected.

COLONIES on the natural substrate effuse, hairy, dark brown. Mycelium superficial and immersed. Hyphae branched, septate, smooth, pale brown, 2–4 µm wide. CONIDIOPHORES macronematous, mononematous, simple or with one branch, erect, straight or slightly flexuous, cylindrical, 3–4-septate, smooth, pale brown, 87.5–225 × 5–7.5 µm; branches cylindrical, 1–2-septate, smooth, pale brown, 27.5–60 × 6–7.5 µm. CONIDIEXOGENOUS CELLS monoblastic, integrated, terminal, cylindrical, with enteroblastic percurrent extensions, smooth, pale brown. Conidial secession rhexolytic. CONIDIA acrogenous, solitary, staurosporous, 42–50(–60) × 13–16 µm, composed of a basal cell that is conical, smooth, pale brown, 7–14 × 5–11 µm, with a tapering base bearing remains of the conidiogenous cells, 2–3 µm wide; 1–2(–3) central cells that are dark brown, smooth, 7–21 × 9–16 µm; 1–3(–4) arms that are conical, 1-septate, 12–30 × 5–11 µm, with a pale brown basal cell, 6–15 × 5–11 µm and a pale brown distal cell, 7–19 × 3–6 µm. Synanamorph *Selenosporella*-like, present at the tip of conidial arms, producing conidia sympodially, aseptate, fusiform, straight, hyaline, 3–5 × 0.5–1 µm.

ADDITIONAL SPECIMEN EXAMINED: BRAZIL, PARÁ, Área de Proteção Ambiental Ilha do Combu, 1°25'S 48°27'W, on submerged wood, 10 Jan. 2013, coll. J.S. Monteiro (HUEFS 196433).

NOTE: *Arachnophora* is typified by *A. fagicola* Hennebert, which was collected from decaying cupules of *Fagus sylvatica* in Belgium (Hennebert 1963). The genus comprises nine species characterized by monoblastic integrated percurrently extending conidiogenous cells that produce staurosporous conidia with dark central cells and paler septate incurved arms (Révay & Gönczöl 1989, Castañeda-Ruiz et al. 1996, 1997, Castañeda-Ruiz & Guarro 1998, Pratibha et al. 2011, Leão-Ferreira et al. 2013, Ma et al. 2013). Among the described




FIG. 1. *Arachnophora combuensis* (holotype, HUEFS 196432). A–B. Conidiophores. C. Conidiogenous cell attached to a conidium. D–P. Conidia. Scale bars: A–B = 30  $\mu$ m; C–P = 20  $\mu$ m.

species only *Arachnophora crassa* Révay & J. Gönczöl and *A. pulneyensis* (Subram. & Bhat) R.F. Castañeda are morphologically similar to *A. combuensis*. *Arachnophora crassa* is distinguished by conidia with 5–7 arms measuring 9–14.5  $\times$  5–6.5  $\mu$ m and a two-celled central body of 16–22.5  $\times$  8–9.5  $\mu$ m (Révay & Gönczöl 1989). *Arachnophora pulneyensis* differs from *A. combuensis* by the

sizes of the central body (16–24 × 6.5–8 µm) and conidial arms (14 × 3–5 µm) and the absence of a basal cell (Castañeda-Ruiz et al. 1996).

#### Acknowledgments

The authors express their sincere gratitude to Dr. De-Wei Li and Dr. Xiu-Guo Zhang for their critical review of the manuscript. The authors are grateful to the Coordination for the Improvement of Higher Education Personnel (CAPES) for financial support and the 'Programa de Pós-graduação em Biologia de Fungos – PPGBF/UFPE'. The authors thank the support provided by 'Programa Ciência sem Fronteiras.' RFCR is grateful to Cuban Ministry of Agriculture and "Programa de Salud Animal y Vegetal," project P131LH003033, Cuban Ministry of Science, Technology and Environment for facilities.

#### Literature cited

- Castañeda-Ruiz RF. 2005. Metodología en el estudio de los hongos anamorfos. Anais do V Congresso Latino Americano de Micologia. Brasília: 182–183.
- Castañeda-Ruiz RF, Guarro J. 1998. Two new hyphomycetes from rainforest of Cuba. Can. J. Bot. 76: 1584–1588. <http://dx.doi.org/10.1139/b98-095>
- Castañeda-Ruiz RF, Guarro J, Cano J. 1996. Notes on conidial fungi. X. A new species of *Ceratosporella* and some new combinations. Mycotaxon 60: 275–281.
- Castañeda-Ruiz RF, Gams W, Saikawa M. 1997. Three new conidial fungi (hyphomycetes) from Cuba. Nova Hedwigia 64: 473–483.
- Hennebert GL. 1963. Un hyphomycète nouveau *Arachnophora fagicola* gen. nov. spec. nov. Can. J. Bot. 41(8): 1165–1169. <http://dx.doi.org/10.1139/b63-097>
- Leão-Ferreira SM, Gusmão LFP, Castañeda-Ruiz RF. 2013. Conidial fungi from the semi-arid Caatinga biome of Brazil. Three new species and new records. Nova Hedwigia 96: 479–494. <http://dx.doi.org/10.1127/0029-5035/2013/0084>
- Ma J, Xia JW, Zhang XG, Castañeda-Ruiz RF. 2013 ["2014"]. *Arachnophora dinghuensis* sp. nov. and *Websteromyces inaequale* sp. nov., and two new records of anamorphic fungi from dead branches of broad-leaved trees in China. Mycoscience 55(5): 329–335. <http://dx.doi.org/10.1016/j.myc.2013.11.007>
- Pratibha J, Bhat S, Raghukumar. 2011. Four anamorphic fungi (with two new species) from forests of Western Ghats, India. Mycotaxon 117: 269–278. <http://dx.doi.org/10.5248/117.269>
- Révay Á, Gönczöl J. 1989. Some dematiaceous hyphomycetes from woody-litter in Hungary. Nova Hedwigia 48: 237–245.