
MYCOTAXON

<http://dx.doi.org/10.5248/129.403>

Volume 129(2), pp. 403–406

October–December 2014

Two new species of *Myrothecium* from the Qinghai-Tibet Plateau Area, China

YUE-MING WU^{1,2^}, YU-LAN JIANG^{3^}, YA-NAN MA^{1,2} & TIAN-YU ZHANG^{1,2*}

¹ Department of Plant Pathology, Shandong Agricultural University, Taian, 271018, China

² Key Laboratory of Agricultural Microbiology, Shandong Province, Taian, 271018, China

³ Agriculture College, Guizhou University, Guiyang, Guizhou, 550025, China

*CORRESPONDENCE TO: tyzhang1937@yahoo.com.cn

ABSTRACT — Two new species from the Qinghai-Tibet Plateau Area (China), *Myrothecium variabile* and *M. xigazense*, are described and illustrated. Specimens (dried cultures) and living cultures are deposited in the Herbarium of Shandong Agricultural University, Plant Pathology (HSAUP), and the Herbarium of Institute of Microbiology, Academia Sinica (HMAS).

KEY WORDS — *Ascomycota*, dematiaceous hyphomycetes, soil fungi, *Stachybotriaceae*, taxonomy

Introduction

During a survey of dematiaceous soil hyphomycetes in China, several unusual *Myrothecium* specimens were collected, of which two represent new species. They are described and illustrated from cultures grown on potato dextrose agar (PDA) (Domsch et al. 2007).

Myrothecium Tode [type species: *M. inundatum* Tode (Tode 1790)] is characterized by conspicuous sporodochia that are cupulate, pustulate or stalked, synnema-like, formed of densely compacted conidiophores arising from a developed stroma and bearing a mass of slimy, green to black conidia. Sporodochia are surrounded by differentiated, often curling, marginal hairs. Conidiophores are hyaline or slightly olivaceous, and branch repeatedly to the ultimate branches, the phialides. Phialides are compacted in a dense parallel layer, cylindrical with conically tapering tips and usually undifferentiated collarettes. Conidia are 1-celled, subhyaline to pale greenish brown but dark olivaceous in mass. Seifert et al. (2011) listed 30 taxa for the genus.

[^]Yue-Ming Wu and Yu-Lan Jiang contributed equally to this work.




FIG. 1. *Myrothecium variabile* (ex holotype HSAUP II₀₇1083).
Conidia, conidiophores, and conidiogenous cells. Scale bars = 25 μ m.

Myrothecium variabile Y.M. Wu & T.Y. Zhang, sp. nov.

FIG. 1

MYCOBANK MB 808618

Differs from *Myrothecium cinctum* and *M. striatisporum* by its longer conidia.

TYPE: China, Tibet: Xigaze, altitude 3100 m, from a forest soil, 11 Sept. 2007, Y.M. Wu (Holotype HSAUP II₀₇1083; isotype HMAS 196284).

ETYMOLOGY: The epithet refers to the variable shape of the conidia.

COLONIES ON PDA effuse, flocculent, pale yellow at initial stage, later green to black. Mycelium superficial or immersed, hyphae branched, septate, smooth, hyaline, 2–3 μ m wide. SPOROCHIA usually not more than 1.0 mm diameter, often confluent, at first green, later black with a white margin, without setae. CONIDIOPHORES straight or flexuous, branched, smooth, with apical branches arranged penicillately. CONIDIOGENOUS CELLS phialides colourless, smooth, 10–25 \times 2–2.5 μ m. CONIDIA unicellular, cylindrical, clavate, fusiform to navicular (L/W=6–11.25), often slightly protuberant and truncate at the base, smooth, pale greenish brown, in mass greenish brown to black, with distinct longitudinal or oblique striations, 12.5–16.25 \times 2.5–3.5 μ m.

COMMENTS: Morphologically, *Myrothecium variabile* resembles *M. cinctum* (Corda) Sacc. (Saccardo 1886) and *M. striatisporum* N.C. Preston (Preston 1948) its longitudinally or obliquely striate conidia. However, *M. cinctum* ($6.5\text{--}14 \times 2.5\text{--}4.5 \mu\text{m}$) and *M. striatisporum* ($7\text{--}12 \times 2.5\text{--}3.5 \mu\text{m}$) have smaller conidia and verrucose conidiophores.




FIG. 2. *Myrothecium xigazense* (ex holotype HSAUP II₀₇0969).
Conidia, conidiophores, setae, and conidiogenous cells. Scale bars = 25 μm .

Myrothecium xigazense Y.M. Wu & T.Y. Zhang, sp. nov.

FIG. 2

MYCOBANK MB 808619

Differs from *Myrothecium carmichaelii* and *M. jollymanii* by its longer conidia.

TYPE: China, Tibet: Xigaze, altitude 3900 m, from a grassland soil, 9 Sept. 2007, Y.M. Wu (Holotype HSAUP II₀₇0969; isotype HMAS 196285).

ETYMOLOGY: in reference to the type locality.

COLONIES on PDA effuse, flocculent, white at initial stage, later black, Mycelium mostly superficial, hyphae branched, septate, smooth, hyaline, 2–3 µm wide. SPOROCHIA usually not more than 1.2 mm diameter, often confluent, at first green, later black with a white margin. SETAE pale brown, smooth, septate, 220–250 µm long, 4–6 µm thick. CONIDIOPHORES straight or flexuous, branched, with apical branches arranged penicillately. CONIDIOGENOUS CELLS phialides colourless, smooth, 8–15×2–2.5 µm. CONIDIA unicellular, cylindrical or clavate, often slightly protuberant and truncate at the base, smooth, pale greenish brown, dark greenish in mass, 12.5–15 × 2–3 µm.

COMMENTS: Morphologically, *Myrothecium xigazense* resembles *M. carmichaelii* Grev. (Greville 1825) and *M. jollymanii* N.C. Preston (Preston 1948) in conidial colour and shape. However, both *M. carmichaelii* (10–11 × 1–1.3 µm) and *M. jollymanii* (10–12 × 2–2.5 µm) have shorter conidia. Additionally, *M. carmichaelii* lacks setae while the setae in *M. jollymanii* are hyaline, not pale brown as in *M. xigazense*.

Acknowledgments

The authors are grateful for pre-submission comments and suggestions provided by Dr. Eric McKenzie, Dr. Yong Wang, and Dr. Shaun Pennycook. This project was supported by the National Science Foundations of China (no. 30970011 & 30499340).

Literature cited

- Domsch KH, Gams W, Anderson TH. 2007. Compendium of soil fungi, ed. 2. IHW-Verlag Eching. 672 p. http://dx.doi.org/10.1111/j.1365-2389.2008.01052_1.x
- Greville RK. 1825. Scottish Cryptogamic Flora 3: pl. 121–180.
- Preston NC. 1948. Observations on the genus *Myrothecium*. II. *Myrothecium gramineum* Lib. and two new species. Transactions of the British Mycological Society 31(3–4): 271–276. [http://dx.doi.org/10.1016/S0007-1536\(48\)80010-0](http://dx.doi.org/10.1016/S0007-1536(48)80010-0)
- Saccardo PA. 1886. *Tubercularia dematiea*, *Amerospora*, *Myrothecium*. Sylloge Fungorum 4: 750–752.
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. 997 p. <http://dx.doi.org/10.3767/003158511X617435>
- Tode HJ. 1790. Fungi mecklenburgensis selecti. 1. 47 p.