

A checklist of xylophilous basidiomycetes (*Basidiomycota*) in mangroves

JULIANO MARCON BALTAZAR, LARISSA TRIERVEILER-PEREIRA
& CLARICE LOGUERCIO-LEITE

jm-baltazar@hotmail.com, lt_pereira@yahoo.com.br, cllite@ccb.ufsc.br

*Laboratório de Micologia, Depto. Botânica, CCB,
Universidade Federal de Santa Catarina*

Campus Universitário, 88090-040, Florianópolis, Santa Catarina, Brazil

Abstract — Based on intensive search of literature records of xylophilous basidiomycetes in mangroves, a list with 112 species is presented. These species are distributed in 63 genera, 27 families and 9 orders. *Polyporaceae* is the most represented family with 33 species; *Phellinus* is the genus with the highest number of species (7). Brazilian mangroves, with 55 species, are the best known areas. The most frequent host is *Rhizophora mangle* with 34 species recorded on it. For each species the localities and substrates are provided, when these data were available in the respective original article. The complete checklist is available on <http://www.mycotaxon.com/resources/weblists.html>.

Key words — *Agaricales, Aphyllophorales, Auriculariales, mycodiversity*

Introduction

Mangroves are transitional coastal ecosystems situated at the confluence of land and sea (Alongi 2002). Their distribution is closely related to basic features of the marine environment, mainly salinity (Chapman 1977). The atmospheric temperature also influences the distribution of mangroves and they are found mostly in the tropics; however under special climatic conditions they occur in subtropical regions, such as Japan and the State of Santa Catarina in Brazil (Cintrón & Schaeffer-Novelli 1980).

Although tropical forests typically have a high diversity of plant species, mangroves are low diversity ecosystems and there are roughly 70 species of mangroves plants (Alongi 2002). The most diverse regions border the Pacific Ocean and west Indian Ocean (Cintrón & Schaeffer-Novelli 1980).

Mangrove species diversity is well known for animals and plants, but poorly known for other organisms such as fungi (Macintosh & Ashton 2002). The study of mangrove fungi began in the 1950's in Australia; however fungal species were reported from these ecosystems earlier by authors who did not study particularly the manglicolous mycota (Schmit & Shearer 2003). The data



FIGURE 1. Mangrove Forest of Itacorubi, Santa Catarina, Southern Brazil (photo by L. Trieveiler-Pereira).

on mangrove fungi refer mainly to ‘marine fungi’, which grow and sporulate exclusively in marine or estuarine habitats (Kohlmeyer & Kohlmeyer 1979). In contrast, little is known about terrestrial fungi in mangrove forests (Hyde & Lee 1995).

The most extensive surveys of mangrove xylophilous basidiomycetes focus on corticioid fungi taxonomy (Maekawa et al. 2003, 2005) or polypore ecology (Gilbert & Sousa 2002, Gilbert et al. 2008). A remarkable number of species are recorded from Northern Brazilian mangroves due to the research of Campos et al. (2003) and Sotão et al. (1991, 2002, 2003). Other records correspond to isolated citations in several articles that have not focused exclusively on mangrove xylophilous basidiomycetes.

The objective of the present work is to compile data on mangrove xylophilous basidiomycetes and to present a list of species with the locality and substrate of each one.

Material and methods

This checklist has been compiled based on an intensive search of literature records of xylophilous basidiomycetes in mangroves. Nomenclature, taxonomic position, and author names followed the databases: Index Fungorum – IFS (<http://www.indexfungorum.org/Names/Names.asp>) and



FIGURE 2. Basidiomata of *Schizophyllum commune* growing on dead trunk of *Avicennia schaueriana* (photo by L. Trierveiler-Pereira).

the International Plant Names Index – IPNI (<http://www.ipni.org>). Genera and species are listed alphabetically within each family and order. Genera with taxonomic position not well established are placed in ‘incertae sedis’. Names not found either on the IFS database or in the literature were placed in a separated section named ‘Doubtful names cited from mangroves’. This checklist only includes records identified to the species level.

Although the compiled list was carefully revised, minor errors can occur. We plan to regularly update the internet version of the checklist, so we gratefully encourage any remarks concerning errors or omitted data.

Results

The 112 xylophilous basidiomycetes species reported from mangroves are distributed among 63 genera, 27 families, and 9 orders. The most represented family is *Polyporaceae* with 33 species, followed by *Hymenochaetaceae* with 15 species. The genus with highest number of species is *Phellinus* with seven species, followed by *Trametes* with six species and *Trichaptum* with five species.

The mangrove areas with the highest number of species recorded are located in Brazil (55 species), Micronesia (19), Japan (17) and Puerto Rico (15). Other localities are represented by eight or fewer reported species.

Rhizophora mangle with 34 species recorded is the most common host, followed by *Bruguiera gymnorhiza* (17) and *Sonneratia alba* (15).

The complete checklist is available on <http://www.mycotaxon.com/resources/weblists.html>.

Acknowledgements

We express our sincere gratitude to many people who kindly provided useful literature: Adriana Gugliotta, Andrea Romero, Ángel Nieves Rivera, Ricardo Drechsler Santos, Helen Sotão, Jan Kohlmeyer, Leif Ryvardeen, Tatiana Gibertoni and Vera Bononi. Many thanks are due to Dr. Mario Rajchenberg and Dr. John Paul Schmit for expert review of the manuscript and English improvements. This study was partly supported by PIBIC/CNPq/UFSC (Brazil).

Literature cited

- Alongi DM. 2002. Present state and the future of the world's mangroves forests. *Environmental Conservation* 29(3): 331–349.
- Campos EL, Sotão HMP, Cavalcanti MAQ, Luz AB. 2003. *Basidiomycetes* de manguezais da APA de Algodual-Maiandeuá, Pará, Brasil. *Bol. Mus. Para. Emílio Goeldi, sér. Ciências Naturais* 1(1): 97–102.
- Chapman VJ. 1977. Introduction. In: Chapman VJ (ed.). *Wet coastal ecosystems*. Oxford: Elsevier.
- Cintrón G, Schaeffer-Novelli Y. 1980. Introducción a la ecología del manglar. San Juan: Departament of Natural Resources and São Paulo: Instituto Oceanográfico, USP.
- Gilbert GS, Gorospe J, Ryvardeen L. 2008. Host and habitat preferences of polypore fungi in Micronesian tropical flooded forests. *Mycol. Res.* 112(6): 674–680.
- Gilbert GS, Sousa WP. 2002. Host specialization among wood-decay polypore fungi in a Caribbean mangrove forest. *Biotropica* 34(3): 396–404.
- Hyde KD, Lee SY. 1995. Ecology of mangrove fungi and their role in nutrient cycle: what gaps occur in our knowledge? *Hydrobiologia* 295: 107–118.
- Kohlmeyer J, Kohlmeyer E. 1979. *Marine mycology: the higher fungi*. New York: Academic Press.
- Macintosh DJ, Ashton EC. 2002. Review of mangrove biodiversity conservation and management. Final Report, University of Aarhus, Denmark.
- Maekawa N, Suhara H, Kinjo K, Kondo R. 2003. Corticioid fungi (*Basidiomycota*) in mangrove forests of the islands of Iriomote and Okinawa, Japan. *Mycoscience* 44: 403–409.
- Maekawa N, Suhara H, Kinjo K, Kondo R, Hoshi Y. 2005. *Haloaleurodiscus mangrovei* gen. sp. nov. (*Basidiomycota*) from mangrove forests in Japan. *Mycol. Res.* 109(7): 825–832.
- Schmit JP, Shearer CA. 2003. A checklist of mangrove-associated fungi, their geographical distribution and known host plants. *Mycotaxon* 85: 423–477.
- Sotão HMP, Bononi VLR, Figueiredo TS. 1991. *Basidiomycetes* de manguezais da Ilha de Maracá, Amapá, Brasil. *Bol. Mus. Para. Emílio Goeldi, ser. Bot.* 7(1): 109–114.
- Sotão HMP, Campos EL, Costa SPSE, Melo OA, Azevedo JC. 2002. *Basidiomycetes* macroscópicos de manguezais de Bragança, Pará, Brasil. *Hoehnea* 29(3): 215–224.
- Sotão HMP, Campos EL, Gugliotta AM, Costa SPSE. 2003. Fungos macroscópicos: *Basidiomycetes*. In: MEB Fernandes (org). *Os manguezais da costa norte brasileira*. Maranhão: Fundação Rio Bracanga.